

Notes: These are notes I took while studying for the Mathematics Subject GRE. There are likely a lot of errors and mistakes, please let me know if you find any!

Undergraduate Compendium

D. Zack Garza

D. Zack GarzaUniversity of Georgia
dzackgarza@gmail.com

 $Last\ updated \hbox{:}\ 2020\hbox{-}11\hbox{-}10$

1	Diffe	erential Calculus	1
	1.1	Big Theorems / Tools:	7
	1.2	Limits	0
	1.3	Tools for finding limits	0
	1.4	Asymptotes	1
	1.5	Recurrences	1
	1.6	Derivatives	1
	1.7	Implicit Differentiation	2
	1.8	Related Rates	2
2	Inte	gral Calculus	2
	2.1	Average Values	
	2.2	Area Between Curves	
	2.3	Solids of Revolution	
	2.4	Arc Lengths	
	2.5	Center of Mass	
	2.6	Big List of Integration Techniques	
	2.0	2.6.1 Integration by Parts:	
		2.6.2 Shoelace Method	
		2.6.3 Differentiating under the integral	
		2.6.4 Partial Fractions	
		2.6.5 Trigonometric Substitution	
	2.7	Optimization	
	2.1	Optimization	1
3	Vec	tor Calculus 1	8
	3.1	Plane Geometry	9
	3.2	Projections	9
	3.3	Lines	C
		3.3.1 Tangent Lines / Planes	C
		3.3.2 Normal Lines	1
	3.4	Planes	1
		3.4.1 Finding a Normal Vector	1
		3.4.2 Distance from origin to plane	2
		3.4.3 Distance from point to plane	2
	3.5	Curves	2
		3.5.1 Tangent line to a curve	2
		3.5.2 Normal line to a curve	2
	3.6	Minimal Distances	3
		3.6.1 Point to plane	
		3.6.2 Point to line	
		3.6.3 Point to curve	
		3.6.4 Line to line	
	3.7	Surfaces	
	5.1	3.7.1 Tangent plane to a surface	
		- or, in the term of the property of the prope	3

4	Mul	tivariable Calculus 2					
	4.1	Notation					
	4.2	Partial Derivatives					
	4.3	General Derivatives					
	4.4	The Chain Rule					
	4.5	Approximation					
	4.6	Optimization					
		4.6.1 Classifying Critical Points					
		4.6.2 Lagrange Multipliers					
	4.7	Change of Variables					
5	Voc	tor Calculus 3					
J	5.1	Notation					
	$5.1 \\ 5.2$	Big Theorems					
	3.2	-					
		5.2.1 Stokes' and Consequences					
	F 9	5.2.2 Directional Derivatives					
	5.3	Computing Integrals					
		5.3.1 Changing Coordinates					
		5.3.2 Line Integrals					
		5.3.3 Flux					
		5.3.4 Area					
	_ ,	5.3.5 Surface Integrals					
	5.4	Other Results					
6	Line	ear Algebra 3					
	6.1	Notation					
	6.2	Big Theorems					
	6.3	Big List of Equivalent Properties					
	6.4	Vector Spaces					
		6.4.1 Linear Transformations					
		6.4.2 Linear Independence					
		6.4.3 Bases					
		6.4.4 The Inner Product					
		6.4.5 Gram-Schmidt Process					
		6.4.6 The Fundamental Subspaces Theorem					
		6.4.7 Computing change of basis matrices					
	6.5	Matrices					
	0.0	6.5.1 Systems of Linear Equations					
		6.5.2 Determinants					
		6.5.3 Computing Determinants					
		6.5.4 Inverting a Matrix					
		6.5.5 Bases for Spaces of a Matrix					
		6.5.6 Eigenvalues and Eigenvectors					
		9					
	6.6	6.5.7 Useful Counterexamples					
	0.0	Dample I folicino					
7		ear Algebra: Advanced Topics 5					
	7 1	Changing Rasis					

	7.2	Orthogonal Matrices	7
	7.3	Projections	7
		7.3.1 Projection Onto a Vector	8
		7.3.2 Projection Onto a Subspace	8
			69
	7.4	Normal Forms	60
	7.5		60
		•	60
8	App	endix: Lists of things to know 6	1
	8.1	_	31
	8.2		$\tilde{52}$
	8.3		3
	8.4		64
	8.5		55
	0.0		,0
9	Ordi	, i	5
	9.1	1	55
	9.2		66
	9.3	Linear Homogeneous	66
	9.4	Linear Inhomogeneous	67
		9.4.1 Undetermined Coefficients	37
		9.4.2 Variation of Parameters	8
		9.4.3 Reduction of Order	8
	9.5	Systems of Differential Equations	8
	9.6		8
	9.7		70
		•	70
	9.8	•	1
	9.9		$\overline{2}$
10	Alge	shra	'2
10	_		2 2
			73
			3
	10.4		75
		- J	7
			7
	10.5	Ring Theory	7
11	Nun	nber Theory 7	8'
	11.1	Notation and Basic Definitions	78
	11.2	Big Theorems	9
			9
	11.4	Divisibility	9
		·	79
			20

	1.5 Modular Arithmetic	80
	11.5.1 Diophantine Equations	81
	11.5.2 Computations	81
	11.5.3 Invertibility	82
	1.6 The Totient Function	83
	1.7 Quadratic Residues	84
	1.8 Primality Tests	85
	1.9 Sequences in Metric Spaces	85
	_	
12	Sequences	85
	2.1 Known Examples	
	2.2 Convergence	
	12.2.1 Checklist	86
12	Sums ("Series")	87
IJ	3.1 Known Examples	
	13.1.1 Conditionally Convergent	
	13.1.2 Convergent	
	13.1.2 Convergent	
	3.2 Convergence	
	13.2.2 Checklist	
	3.3 Radius of Convergence	
	.g.g Itadius of Convergence	91
14	Real Analysis	91
	4.1 Notation	92
		92
		93
	4.4 Limits	93
	4.5 Commuting Limits	94
	4.6 Continuity	97
	14.6.1 Lipschitz Continuity	98
	4.7 Differentiability	98
	14.7.1 Properties, strongest to weakest	99
	4.8 Giant Table of Relations	99
	4.9 Integrability	100
	4.10List of Free Conclusions:	101
	4.11Convergence	101
	14.11.1 Sequences and Series of Functions	101
	14.11.2 Pointwise convergence	102
	14.11.3 Uniform Convergence	
	4.12Topology	
	4.13Counterexamples	
15	1 63	108
	F 1 D.C.::::	100

16	Prob	pability	11(
	16.1	Definitions	. 110
	16.2	Theory and Background	113
	16.3	Distributions	115
		16.3.1 Uniform	. 115
		16.3.2 Bernoulli	115
		16.3.3 Binomial	116
		16.3.4 Poisson	116
		16.3.5 Negative Binomial	116
		16.3.6 Geometric	
		16.3.7 Hypergeometric	. 117
		16.3.8 Normal	
	16.4	Table of Distributions	. 118
		Common Problems	
	16.6	Notes, Shortcuts, Misc	. 119
17		nbinatorics	119
	17.1	Notation	. 120
	17.2	The Important Numbers	
		17.2.1 Common Problems	. 122
		17.2.2 The Twelvefold Way	. 122
10	_		101
18	Com	nplex Analysis	123
		Useful Equations and Definitions	
	18.2	Complex Arithmetic and Calculus	
	100	18.2.1 Complex Differentiability	
		Complex Integrals	
		Definitions	
	18.5	Preliminary Notions	
		18.5.1 What is the Complex Derivative?	
		18.5.2 <i>n</i> th roots of a complex number	
	10.0	18.5.3 The Cauchy-Riemann Equations	
	18.6	Integration	
		18.6.1 The Residue Theorem	
		18.6.2 Computing Residues	
		18.6.3 Simple Poles	
		18.6.4 Rational Functions	
	10 =	18.6.5 Computing Integrals	
	18.7	Conformal Maps	
		18.7.1 Applications	
		18.7.2 Theorems	
	100	18.7.3 Some Useful Formulae	
	18.8	Questions	138
19	Com	nmon Mistakes	139
ეი	Λ	endix 1	139
4 0		Nest Tricks	130

2	20.2 Big Derivative / Integral Table	41
2	20.3 Useful Series and Sequences	42
2	20.4 Partial Fraction Decomposition	14
2	20.5 Properties of Norms	44
	20.6 Logic Identities	
2	20.7 Set Identities	45
2	20.8 Preimage Identities	45
	20.9 Pascal's Triangle:	
2	20.10Table of Small Factorials	47
2	20.11Primes Under 100:	48
2	20.12Checking Divisibility by Small Numbers	48
	20.13Hyperbolic Functions	
2	20.14Integral Tables	50
21 [Definitions 15	51
2	21.1 Set Theory	51
	21.2 Calculus	
	21.3 Analysis	
	21.4 Linear Algebra	
	21.5 Differential Equations	
	21.6 Algebra	
	21.7 Complex Analysis	
	21.8 Algebra	
	21.9 Geometry	
22 I	ndices 16	63

1 | Differential Calculus

1.1 Big Theorems / Tools:

Proposition 1.1.1 (Fundamental Theorem of Calculus I).

$$\frac{\partial}{\partial x} \int_{a}^{x} f(t)dt = f(x)$$

Proposition $1.1.2 (Generalized\ Fundamental\ Theorem\ of\ Calculus).$

$$\frac{\partial}{\partial x} \int_{a(x)}^{b(x)} f(x,t)dt - \int_{a(x)}^{b(x)} \frac{\partial}{\partial x} f(x,t)dt = f(x,t) \cdot \frac{\partial}{\partial x} (t) \Big|_{t=a(x)}^{t=b(x)}$$

$$= f(x,b(x)) \cdot b'(x) - f(x,a(x)) \cdot a'(x)$$

If f(x,t) = f(t) doesn't depend on x, then $\frac{\partial f}{\partial x} = 0$ and the second integral vanishes:

$$\frac{\partial}{\partial x} \int_{a(x)}^{b(x)} f(t)dt = f(b(x)) \cdot b'(x) - f(a(x)) \cdot a'(x)$$

Find examples

Remark 1.1.1: Note that you can recover the original FTC by taking

$$a(x) = c$$
$$b(x) = x$$
$$f(x,t) = f(t).$$

Corollary 1.1.1(?).

$$\frac{\partial}{\partial x} \int_{1}^{x} f(x,t)dt = \int_{1}^{x} \frac{\partial}{\partial x} f(x,t)dt + f(x,x)$$

 ${\bf Proposition} \ 1.1.3 ({\it Extreme \ Value \ Theorem}).$

Todo

Todo

Proposition 1.1.4 (Mean Value Theorem).

$$f \in C^0(I) \implies \exists p \in I : f(b) - f(a) = f'(p)(b - a)$$

 $\implies \exists p \in I : \int_a^b f(x) \ dx = f(p)(b - a).$

Proposition 1.1.5 (Rolle's Theorem).

todo

Proposition 1.1.6 (L'Hopital's Rule).

Ιf

• f(x) and g(x) are differentiable on $I - \{pt\}$, and

$$\lim_{x \to \{\text{pt}\}} f(x) = \lim_{x \to \{\text{pt}\}} g(x) \in \{0, \pm \infty\}, \qquad \forall x \in I, g'(x) \neq 0, \qquad \lim_{x \to \{\text{pt}\}} \frac{f'(x)}{g'(x)} \text{ exists},$$

Then it is necessarily the case that

$$\lim_{x \to \{\text{pt}\}} \frac{f(x)}{g(x)} = \lim_{x \to \{\text{pt}\}} \frac{f'(x)}{g'(x)}.$$

Remark 1.1.2: Note that this includes the following indeterminate forms:

$$\frac{0}{0}$$
, $\frac{\infty}{\infty}$, $0 \cdot \infty$, 0^0 , ∞^0 , 1^∞ , $\infty - \infty$.

For $0 \cdot \infty$, can rewrite as $\frac{0}{\frac{1}{\infty}} = \frac{0}{0}$, or alternatively $\frac{\infty}{\frac{1}{0}} = \frac{\infty}{\infty}$.

For 1^{∞} , ∞^0 , and 0^0 , set

$$L := \lim f^g \implies \ln L = \lim g \ln(f)$$

to recover $\infty \cdot 0, 0 \cdot \infty$, or $0 \cdot 0$.

Proposition 1.1.7 (Taylor Expansion).

$$T(a,x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

$$= f(a) + f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \frac{1}{6}f'''(a)(x-a)^3 + \frac{1}{24}f^{(4)}(a)(x-a)^4 + \cdots$$

There is a bound on the error:

$$|f(x) - T_k(a, x)| \le \left| \frac{f^{(k+1)}(a)}{(k+1)!} \right|$$

where $T_k(a,x) = \sum_{n=0}^k \frac{f^{(n)}(a)}{n!} (x-a)^n$ is the kth truncation.

Remark 1.1.3 : Approximating change: $\Delta y \approx f'(x)\Delta x$

1.2 Limits

1.3 Tools for finding limits

Examples

How to find $\lim_{x\to a} f(x)$ in order of difficulty:

- Plug in: if f is continuous, $\lim_{x\to a} f(x) = f(a)$.
- Check for indeterminate forms and apply L'Hopital's Rule.
- Algebraic rules
- Squeeze theorem
- Expand in Taylor series at a
- Monotonic + bounded
- One-sided limits: $\lim_{x\to a^-} f(x) = \lim_{\varepsilon\to 0} f(a-\varepsilon)$
- Limits at zero or infinity:

$$\lim_{x \to \infty} f(x) = \lim_{\frac{1}{x} \to 0} f\left(\frac{1}{x}\right) \text{ and } \lim_{x \to 0} f(x) = \lim_{x \to \infty} f\left(\frac{1}{x}\right)$$

– Also useful: if $p(x) = p_n x^n + \cdots$ and $q(x) = q_n x^m + \cdots$,

$$\lim_{x \to \infty} \frac{p(x)}{q(x)} = \begin{cases} 0 & \deg p < \deg q \\ \infty & \deg p > \deg q \\ \frac{p_n}{q_n} & \deg p = \deg q \end{cases}$$

 \triangle Warning 1.1: Be careful: limits may not exist!! Example: $\lim_{x\to 0} \frac{1}{x} \neq 0$.

1.4 Asymptotes

1.2 Limits 10

- Vertical asymptotes: at values x=p where $\lim_{x\to p}=\pm\infty$
- Horizontal asymptotes: given by points y=L where $L\lim_{x\to\pm\infty}f(x)<\infty$
- Oblique asymptotes: for rational functions, divide terms without denominators yield equation of asymptote (i.e. look at the asymptotic order or "limiting behavior").
 - Concretely:

$$f(x) = \frac{p(x)}{q(x)} = r(x) + \frac{s(x)}{t(x)} \sim r(x)$$

1.5 Recurrences

- Limit of a recurrence: $x_n = f(x_{n-1}, x_{n-2}, \cdots)$
 - If the limit exists, it is a solution to x = f(x)

1.6 Derivatives

Proposition 1.6.1 (Chain Rule).

$$\frac{\partial}{\partial x} (f \circ g) = (f' \circ g) \cdot g'$$

Proposition 1.6.2 (Product Rule).

$$\frac{\partial}{\partial x} f \cdot g = f' \cdot g + g' \cdot f$$

Proposition 1.6.3 (Quotient Rule).

$$\frac{\partial}{\partial x} \frac{f(x)}{g(x)} = \frac{f'g - g'f}{g^2}$$

Mnemonic: Low d-high minus high d-low

11

Proposition 1.6.4 (Inverse Rule).

$$\frac{\partial f^{-1}}{\partial x} \left(f(x_0) \right) = \left(\frac{\partial f}{\partial x} \right)^{-1} (x_0) = 1/f'(x_0)$$

1.7 Implicit Differentiation



$$(f(x))' = f'(x) dx, (f(y))' = f'(y) dy$$

- Often able to solve for $\frac{\partial y}{\partial x}$ this way.
 - Obtaining derivatives of inverse functions: if $y = f^{-1}(x)$ then write f(y) = x and implicitly differentiate.

1.8 Related Rates



General series of steps: want to know some unknown rate y_t

- Lay out known relation that involves y
- Take derivative implicitly (say w.r.t t) to obtain a relation between y_t and other stuff.
- Isolate $y_t = \text{known stuff}$

Example 1.8.1 (?): Example: ladder sliding down wall

• Setup: l, x_t and x(t) are known for a given t, want y_t .

$$x(t)^{2} + y(t)^{2} = l^{2} \implies 2xx_{t} + 2yy_{t} = 2ll_{t} = 0$$

(noting that l is constant)

$$- So y_t = -\frac{x(t)}{y(t)} x_t$$

– x(t) is known, so obtain $y(t) = \sqrt{l^2 - x(t)^2}$ and solve.

2 | Integral Calculus

2.1 Average Values



Proposition 2.1.1 (Integral formula for average value).

$$\mu_f = \frac{1}{b-a} \int_a^b f(t)dt$$

Proof (?). Apply MVT to F(x).

2.2 Area Between Curves

Area in polar coordinates:

$$A = \int_{r_1}^{r_2} \frac{1}{2} r^2(\theta) \ d\theta$$

2.3 Solids of Revolution

Disks

Cylinders

$$A = \int \pi r(t)^2 dt$$
$$A = \int 2\pi r(t)h(t) dt.$$

2.4 Arc Lengths

$$L = \int ds$$

$$= \int_{x_0}^{x_1} \sqrt{1 + \frac{\partial y}{\partial x}} dx$$


$$= \int_{y_0}^{y_1} \sqrt{\frac{\partial x}{\partial y} + 1} dy$$

$$ds = \sqrt{dx^2 + dy^2}$$

$$SA = \int 2\pi r(x) \ ds$$

2.1 Average Values

2.5 Center of Mass



$$x_i = \frac{\int_R x_i \rho(x) \ dx}{\int_R \rho(x) \ dx}$$

2.6 Big List of Integration Techniques

Given f(x), we want to find an antiderivative $F(x) = \int f$ satisfying $\frac{\partial}{\partial x} F(x) = f(x)$

- Guess and check: look for a function that differentiates to f.
- *u* substitution
 - More generally, any change of variables

$$x = g(u) \implies \int_{a}^{b} f(x) \ dx = \int_{q^{-1}(a)}^{g^{-1}(b)} (f \circ g)(x) \ g'(x) \ dx$$

2.6.1 Integration by Parts:

The standard form:

$$\int udv = uv - \int vdu$$

• A more general form for repeated applications: let $v^{-1} = \int v, v^{-2} = \int \int v$, etc.

$$\int_{a}^{b} uv = uv^{-1} \Big|_{a}^{b} - \int_{a}^{b} u^{1}v^{-1}$$

$$= uv^{-1} - u^{1}v^{-2} \Big|_{a}^{b} + \int_{a}^{b} u^{2}v^{-2}$$

$$= uv^{-1} - u^{1}v^{-2} + u^{2}v^{-3} \Big|_{a}^{b} - \int_{a}^{b} u^{3}v^{-3}$$

$$\vdots$$

$$\implies \int_{a}^{b} uv = \sum_{k=1}^{n} (-1)^{k} u^{k-1} v^{-k} \Big|_{a}^{b} + (-1)^{n} \int_{a}^{b} u^{n} v^{-n}$$

14

• Generally useful when one term's nth derivative is a constant.

2.5 Center of Mass

2.6.2 Shoelace Method

• Note: you can choose u or v equal to 1! Useful if you know the derivative of the integrand.

Derivatives	Integrals	Signs	Result
\overline{u}	v	NA	NA
u'	$\int v$	+	$u\int v$
u''	$\int \int v$	_	
:	:	:	:

Fill out until one column is zero (alternate signs). Get the result column by multiplying diagonally, then sum down the column.

2.6.3 Differentiating under the integral

$$\frac{\partial}{\partial x} \int_{a(x)}^{b(x)} f(x,t)dt - \int_{a(x)}^{b(x)} \frac{\partial}{\partial x} f(x,t)dt = f(x,\cdot) \frac{\partial}{\partial x} (\cdot) \Big|_{a(x)}^{b(x)}$$
$$= f(x,b(x)) \ b'(x) - f(x,a(x)) \ a'(x)$$

Proof (?).

Let F(x) be an antiderivative and compute F'(x) using the chain rule.

For constants, this should allow differentiating under the integral when f, f_x are "jointly continuous"

- LIPET: Log, Inverse trig, Polynomial, Exponential, Trig: generally let u be whichever one comes first.
- The ridiculous trig sub: for any integrand containing only trig terms
 - Transforms any such integrand into a rational function of x
 - Let $u = 2 \tan^{-1} x$, $du = \frac{2}{x^2 + 1}$, then

$$\int_{a}^{b} f(x) \ dx = \int_{\tan \frac{a}{2}}^{\tan \frac{b}{2}} f(u) \ du$$

Example 2.6.1 (?):

$$\int_0^{\pi/2} \frac{1}{\sin \theta} \ d\theta = 1/2$$

• Trigonometric Substitution

$$\sqrt{a^2 - x^2} \qquad \Rightarrow \qquad x = a\sin(\theta) \qquad dx = a\cos(\theta) \ d\theta$$

$$\sqrt{a^2 + x^2} \qquad \Rightarrow \qquad x = a\tan(\theta) \qquad dx = a\sec^2(\theta) \ d\theta$$

$$\sqrt{x^2 - a^2} \qquad \Rightarrow \qquad x = a\sec(\theta) \qquad dx = a\sec(\theta)\tan(\theta) \ d\theta$$

2.6.4 Partial Fractions

2.6.5 Trigonometric Substitution

Completing the square

• Trig Formulas

$$\sin^2(x) = \frac{1}{2}(1 - 2\cos x)$$

$$=$$

$$=$$

$$=$$

$$=$$

Trig functions, double angle formulas.

• Products of trig functions

- Setup:
$$\int \sin^{a}(x) \cos^{b}(x) dx$$

$$\Leftrightarrow \text{Both } a, b \text{ even: } \sin(x) \cos(x) = \frac{1}{2} \sin(x)$$

$$\Leftrightarrow a \text{ odd: } \sin^{2} = 1 - \cos^{2}, \ u = \cos(x)$$

$$\Leftrightarrow b \text{ odd: } \cos^{2} = 1 - \sin^{2}, \ u = \sin(x)$$
- Setup:
$$\int \tan^{a}(x) \sec^{b}(x) dx$$

$$\Leftrightarrow a \text{ odd: } \tan^{2} = \sec^{2} -1, \ u = \sec(x)$$

$$\Leftrightarrow b \text{ even: } \sec^{2} = \tan^{2} -1, u = \tan(x)$$

Other small but useful facts:

$$\int_0^{2\pi} \sin\theta \ d\theta = \int_0^{2\pi} \cos\theta \ d\theta = 0.$$

2.7 Optimization



- Critical points: boundary points and wherever f'(x) = 0
- Second derivative test:

$$-f''(p) > 0 \implies p \text{ is a min}$$

 $-f''(p) < 0 \implies p \text{ is a max}$

- Inflection points of h occur where the tangent of h' changes sign. (Note that this is where h' itself changes sign.)
- Inverse function theorem: The slope of the inverse is reciprocal of the original slope
- If two equations are equal at exactly one real point, they are tangent to each other there therefore their derivatives are equal. Find the x that satisfies this; it can be used in the original equation.
- Fundamental theorem of Calculus: If

$$\int f(x)dx = F(b) - F(a) \implies F'(x) = f(x).$$

- Min/maxing either derivatives of Lagranage multipliers!
- Distance from origin to plane: equation of a plane

$$P: ax + by + cz = d.$$

- You can always just read off the normal vector $\mathbf{n} = (a, b, c)$. So we have $\mathbf{n}\mathbf{x} = d$.
- Since $\lambda \mathbf{n}$ is normal to P for all λ , solve $\mathbf{n}\lambda \mathbf{n} = d$, which is $\lambda = \frac{d}{\|\mathbf{n}\|^2}$
- A plane can be constructed from a point p and a normal n by the equation np = 0.
- In a sine wave $f(x) = \sin(\omega x)$, the period is given by $2\pi/\omega$. If $\omega > 1$, then the wave makes exactly ω full oscillations in the interval $[0, 2\pi]$.
- The directional derivative is the gradient dotted against a *unit vector* in the direction of interest
- Related rates problems can often be solved via implicit differentiation of some constraint function
- The second derivative of a parametric equation is not exactly what you'd intuitively think!

• For the love of god, remember the FTC!

$$\frac{\partial}{\partial x} \int_0^x f(y) dy = f(x)$$

- Technique for asymptotic inequalities: WTS f < g, so show $f(x_0) < g(x_0)$ at a point and then show $\forall x > x_0, f'(x) < g'(x)$. Good for big-O style problems too.
- Inflection points of h occur where the tangent of h' changes sign. (Note that this is where h' itself changes sign.)
- Inverse function theorem: The slope of the inverse is reciprocal of the original slope
- If two equations are equal at exactly one real point, they are tangent to each other there therefore their derivatives are equal. Find the x that satisfies this; it can be used in the original equation.
- Fundamental theorem of Calculus: If

$$\int f(x)dx = F(b) - F(a) \implies F'(x) = f(x).$$

- Min/maxing either derivatives of Lagranage multipliers!
- Distance from origin to plane: equation of a plane

$$P: ax + by + cz = d.$$

3 | Vector Calculus

Need lots of pictures

Notation:

$$\mathbf{v}, \mathbf{a}, \cdots$$
 vectors in \mathbb{R}^n $\mathbf{R}, \mathbf{A}, \cdots$ matrices $\mathbf{r}(t)$ A parameterized curve $\mathbf{r}: \mathbb{R} \to \mathbb{R}^n$ $\widehat{\mathbf{v}}$

Vector Calculus 18

3.1 Plane Geometry



Proposition 3.1.1(Slope of a vector in \mathbb{R}^2).

$$\mathbf{v} = [x, y] \in \mathbb{R}^2 \implies m = \frac{y}{x}.$$

Proposition 3.1.2(Rotation matrices in \mathbb{R}^2).

$$\mathbf{R}_{\theta} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \implies \mathbf{R}_{\frac{\pi}{2}} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

Corollary 3.1.1(?).

$$\mathbf{R}_{\frac{\pi}{2}}\mathbf{x} \coloneqq \mathbf{R}_{\frac{\pi}{2}} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix} \in \mathbb{R}\mathbf{x}^{\perp}.$$

Thus if a planar line is defined by the span of [x,y] and a slope of m=y/x, a normal vector is given by the span of [-y,x] of slope $-\frac{1}{m}=-x/y$.

Example 3.1.1 (?): Given \mathbf{v} , the rotated vector $\mathbf{R}_{\frac{\pi}{2}}\mathbf{v}$ is orthogonal to \mathbf{v} , so this can be used to obtain normals and other orthogonal vectors in the plane.

Proposition 3.1.3.

There is a direct way to come up with one orthogonal vector to any given vector:

$$\mathbf{v} = [a,b,c] \implies \mathbf{y} \coloneqq \begin{cases} [-(b+c),a,a] & \mathbf{v} = [-1,-1,0], \\ [c,c,-(a+b)] & \text{else} \end{cases}$$

3.2 Projections



For a subspace given by a single vector **a**:

$$\operatorname{proj}_{\mathbf{a}}(\mathbf{x}) = \langle \mathbf{x}, \mathbf{a} \rangle \widehat{\mathbf{a}}$$
 $\operatorname{proj}_{\mathbf{a}}(\mathbf{x}) = \mathbf{x} - \operatorname{proj}_{\mathbf{a}}(\mathbf{x}) = \mathbf{x} - \langle \mathbf{x}, \mathbf{a} \rangle \widehat{\mathbf{a}}$

In general, for a subspace $colspace(A) = \{\mathbf{a}_1, \cdots \mathbf{a}_n\},\$

Vector Calculus

$$\operatorname{proj}_{A}(\mathbf{x}) = \sum_{i=1}^{n} \langle \mathbf{x}, \mathbf{a}_{i} \rangle \widehat{\mathbf{a}}_{i} = A(A^{T}A)^{-1}A^{T}\mathbf{x}$$

3.3 Lines

General Equation

$$Ax + By + C = 0$$

Parametric Equation

$$\mathbf{r}(t) = t\mathbf{x} + \mathbf{b}.$$

Characterized by an equation in inner products:

$$\mathbf{y} \in L \iff \langle \mathbf{y}, \mathbf{n} \rangle = 0$$

Proposition 3.3.1 (Equation for a line between two points).

Given $\mathbf{p}_0, \mathbf{p}_1$, take $\mathbf{x} = \mathbf{p}_1 - \mathbf{p}_0$ and $\mathbf{b} = \mathbf{p}_i$ for either i:

$$\mathbf{r}(t) = t(\mathbf{p}_1 - \mathbf{p}_0) + \mathbf{p}_0$$
 $= t\mathbf{p}_1 + (1 - t)\mathbf{p}_0.$

Proposition 3.3.2(Symmetric equation of a line).

If a line L is given by

$$\mathbf{r}(t) = t[x_1, x_2, x_3] + [p_1, p_2, p_3],$$

then

$$(x, y, z) \in L \iff \frac{x - p_1}{x_1} = \frac{y - p_2}{x_2} = \frac{z - p_3}{x_3}.$$

Example 3.3.1 (?): The symmetric equation of the line through [2,1,-3] and [1,4,-3] is given by

$$\frac{x-2}{1} = \frac{y+1}{-5} = \frac{z-3}{6}.$$

3.3.1 Tangent Lines / Planes

Key idea: just need a point and a normal vector, and the gradient is normal to level sets.

Theorem 3.3.1 (The Tangent Plane Equation).

For any locus $f(\mathbf{x}) = 0$, we have

$$\mathbf{x} \in T_f(\mathbf{p}) \implies \langle \nabla f(\mathbf{p}), \ \mathbf{x} - \mathbf{p} \rangle = 0.$$

3.3.2 Normal Lines

Key idea: the gradient is normal.

To find a normal line, you just need a single point \mathbf{p} and a normal vector \mathbf{n} ; then

$$L = \left\{ \mathbf{x} \mid \mathbf{x} = \mathbf{p} + t\mathbf{v} \right\}.$$

3.4 Planes

General Equation

$$Ax + By + Cz + D = 0$$

Parametric Equation

$$\mathbf{y}(t,s) = t\mathbf{x}_1 + s\mathbf{x}_2 + \mathbf{b}$$

Characterized by an equation in inner products:

$$\mathbf{y} \in P \iff \langle \mathbf{y} - \mathbf{p}_0, \mathbf{n} \rangle = 0$$

Proposition 3.4.1 (Writing equation from a point and a normal).

Determined by a point \mathbf{p}_0 and a normal vector \mathbf{n}

Proposition 3.4.2 (Writing equation from two vectors).

Given $\mathbf{v}_0, \mathbf{v}_1$, set $\mathbf{n} = \mathbf{v}_0 \times \mathbf{v}_1$.

3.4.1 Finding a Normal Vector

- Normal vector to a plane
 - Can read normal off of equation: $\mathbf{n} = [a, b, c]$

3 Vector Calculus

- Computing *D*:
 - $-D = \langle \mathbf{p}_0, \mathbf{n} \rangle = p_1 n_1 + p_2 n_2 + p_3 n_3$
 - Useful trick: once you have \mathbf{n} , you can let \mathbf{p}_0 be any point in the plane (don't necessarily need to use the one you started with, so pick any point that's convenient to calculate)

3.4.2 Distance from origin to plane

• Given by $D/\|\mathbf{n}\| = \langle \mathbf{p}_0, \ \hat{\mathbf{n}} \rangle$. Gives a signed distance.

Distance from origin to plane.

3.4.3 Distance from point to plane

- Given by $\langle \, \cdot \, , \, \, \widehat{\mathbf{n}} \rangle$
- Finding vectors in the plane
- Given $P = [A, B, C] \cdot [x, y, z] = 0$, can take $\left[-\frac{B}{A}, 1, 0 \right], \left[-\frac{C}{A}, 0, 1 \right]$

Distance from point to plane

3.5 Curves

$$\mathbf{r}(t) = [x(t), y(t), z(t)].$$

3.5.1 Tangent line to a curve

We have an equation for the tangent vector at each point:

$$\widehat{\mathbf{T}}(t) = \widehat{\mathbf{r}'}(t),$$

so we can write

$$\mathbf{L}_T(t) = \mathbf{r}(t_0) + t\widehat{\mathbf{T}}(t_0) := \mathbf{r}(t_0) + t\widehat{\mathbf{r}'}(t_0).$$

3.5.2 Normal line to a curve

• Use the fact that $\mathbf{r}''(t) \in \mathbb{R}\mathbf{r}'(t)^{\perp}$, so we have an equation for a normal vector at each point:

$$\widehat{\mathbf{N}}(t) = \widehat{\mathbf{r}''}(t).$$

Thus we can write

$$\mathbf{L}_N(t) = \mathbf{r}(t_0) + t\widehat{\mathbf{N}}(t_0) = \mathbf{r}(t_0) + t\widehat{\mathbf{r}''}(t_0).$$

Special case: planar graphs of functions Suppose y = f(x). Set g(x,y) = f(x) - y, then

$$\nabla g = [f_x(x), -1] \implies m = -\frac{1}{f_x(x)}$$

3.6 Minimal Distances

Fix a point **p**. Key idea: find a subspace and project onto it.

Key equations: projection and orthogonal projection of **b** onto **a**:

$$\operatorname{proj}_{\mathbf{a}}(\mathbf{b}) = \langle \mathbf{b}, \ \mathbf{a} \rangle \widehat{\mathbf{a}}$$

$$\operatorname{proj}_{\mathbf{a}}^{\perp}(\mathbf{b}) = \mathbf{b} - \operatorname{proj}_{\mathbf{a}}(\mathbf{b}) = \mathbf{b} - \langle \mathbf{b}, \ \mathbf{a} \rangle \widehat{\mathbf{a}}$$

3.6.1 Point to plane

• Given a point \mathbf{p} and a plane $S = \{\mathbf{x} \in \mathbb{R}^3 \mid n_0x + n_1y + n_2z = d\}$, let $\mathbf{n} = [n_1, n_2, n_3]$, find any point $\mathbf{q} \in S$, and project $\mathbf{q} - \mathbf{p}$ onto $S^{\perp} = \operatorname{Span}(\mathbf{n})$ using

$$d = \|\operatorname{proj}_{\mathbf{n}}(\mathbf{q} - \mathbf{p})\| = \|\langle \mathbf{q} - \mathbf{p}, \ \mathbf{n} \rangle \widehat{\mathbf{n}}\| = \langle \mathbf{q} - \mathbf{p}, \ \mathbf{n} \rangle.$$

• Given just two vectors \mathbf{u}, \mathbf{v} : manufacture a normal vector $\mathbf{n} = \mathbf{u} \times \mathbf{v}$ and continue as above.

Origin to plane Special case: if p = 0,

$$d = \|\operatorname{proj}_{\mathbf{n}}(\mathbf{q})\| = \|\langle \mathbf{p}, \ \mathbf{n} \rangle \widehat{\mathbf{n}}\| = \langle \mathbf{p}, \ \mathbf{n} \rangle..$$

3.6.2 Point to line

• Given a line $L: \mathbf{x}(t) = t\mathbf{v}$ for some fixed \mathbf{v} , use

$$d = \left\| \operatorname{proj}_{\mathbf{v}}^{\perp}(\mathbf{p}) \right\| = \left\| \mathbf{p} - \langle \mathbf{p}, \ \mathbf{v} \rangle \widehat{\mathbf{v}} \right\|.$$

• Given a line $L: \mathbf{x}(t) = \mathbf{w}_0 + t\mathbf{w}$, let $\mathbf{v} = \mathbf{x}(1) - \mathbf{x}(0)$ and proceed as above.

Wector Calculus

3.6.3 Point to curve

todo

3.6.4 Line to line

Given $\mathbf{r}_1(t) = \mathbf{p}_1 + t\mathbf{v}_2$ and $\mathbf{r}_2(t) = \mathbf{p}_2 + t\mathbf{v}_2$, let d be the desired distance.

- Let $\hat{\mathbf{n}} = \widehat{\mathbf{v}_1 \times \mathbf{v}_2}$, which is orthogonal to both lines.
- Then project the vector connecting the two fixed points \mathbf{p}_i onto this subspace and take its norm:

$$d = \|\operatorname{proj}_{\mathbf{n}}(\mathbf{p}_{2} - \mathbf{p}_{1})\|$$

$$= \|\langle \mathbf{p}_{2} - \mathbf{p}_{1}, \ \mathbf{n} \rangle \hat{\mathbf{n}}\|$$

$$= \langle \mathbf{p}_{2} - \mathbf{p}_{1}, \ \mathbf{n} \rangle$$

$$\coloneqq \langle \mathbf{p}_{2} - \mathbf{p}_{1}, \ \mathbf{v}_{1} \times \mathbf{v}_{2} \rangle.$$

3.7 Surfaces

$$S = \left\{ (x, y, z) \mid f(x, y, z) = 0 \right\}$$
 $z = f(x, y)$

3.7.1 Tangent plane to a surface

- Need a point \mathbf{p} and a normal \mathbf{n} . By cases:
- f(x, y, z) = 0
 - ∇f is a normal vector.
 - Write the tangent plane equation $\langle \mathbf{n}, \mathbf{x} \mathbf{p}_0 \rangle$, done.
- z = g(x, y):
 - Let f(x, y, z) = g(x, y) z, then $\mathbf{p} \in S \iff \mathbf{p}$ is in a level set of f.
 - $-\nabla f$ is normal to level sets (and thus the surface), so compute $\nabla f = [g_x, g_y, -1]$
 - Proceed as in previous case

3.7.2 Surfaces of revolution

• Given $f(x_1, x_2) = 0$, can be revolved around either the x_1 or x_2 axis.

- f(x,y) around the x axis yields $f(x,\pm\sqrt{y^2+z^2})=0$
- -f(x,y) around the y axis yields $f(\pm\sqrt{x^2+z^2},y)=0$
- Remaining cases proceed similarly leave the axis variable alone, replace other variable with square root involving missing axis.
- Equations of lines tangent to an intersection of surfaces f(x, y, z) = g(x, y, z):
 - Find two normal vectors and take their cross product, e.g. $n = \nabla f \times \nabla g$, then

$$L = \left\{ \mathbf{x} \mid \mathbf{x} = \mathbf{p} + t\mathbf{n} \right\}$$


- Level curves:
 - Given a surface f(x, y, z) = 0, the level curves are obtained by looking at e.g. f(x, y, c) = 0.

4 Multivariable Calculus

Theorem 4.0.1 (Key Theorem).

Given a function $f: \mathbb{R}^n \to \mathbb{R}$, let $S_k := \{ \mathbf{p} \in \mathbb{R}^n \mid f(\mathbf{p}) = k \}$ denote the level set for $k \in \mathbb{R}$. Then

$$\nabla f(\mathbf{p}) \in S_k^{\perp}$$
.





$$\mathbf{v} = [v_1, v_2, \cdots]$$

$$\mathbf{e}_i = [0, 0, \cdots, \overbrace{1}^{i \text{th term}}, \cdots, 0]$$

the ith standard basis vector

$$\varphi: \mathbb{R}^n \to \mathbb{R}$$

$$\varphi(x_1, x_2, \cdots) = \cdots$$

a functional on
$$\mathbb{R}^n$$

$$\mathbf{F}: \mathbb{R}^n \to \mathbb{R}^n$$

$$\mathbf{F}(x_1, x_2, \dots) = [\mathbf{F}_1(x_1, x_2, \dots), \mathbf{F}_2(x_1, x_2, \dots), \dots, \mathbf{F}_n(x_1, x_2, \dots)]$$

a multivariable function

Multivariable Calculus 25

4.2 Partial Derivatives

Definition 4.2.1 (Partial Derivative).

For a functional $f: \mathbb{R}^n \to \mathbb{R}$, the **partial derivative** of f with respect to x_i is

$$\frac{\partial f}{\partial x_i}(\mathbf{p}) \coloneqq \lim_{h \to 0} \frac{f(\mathbf{p} + h\mathbf{e}_i) - f(\mathbf{p})}{h}$$

Example 4.2.1 (n = 2):

$$f: \mathbb{R}^2 \to \mathbb{R}$$
$$\frac{\partial f}{\partial x}(x_0, y_0) = \lim_{h \to 0} \frac{f(x_0 + h, y_0) - f(x_0, y_0)}{h}$$

4.3 General Derivatives

Definition 4.3.1 (General definition of differentiability).

A function $f: \mathbb{R}^n \to \mathbb{R}^m$ is **differentiable** iff there exists a linear transformation $D_f: \mathbb{R}^n \to \mathbb{R}^m$ such that the following limit exists

$$\lim_{\mathbf{x}\to\mathbf{p}}\frac{\|f(\mathbf{x})-f(\mathbf{p})-D_f(\mathbf{x}-\mathbf{p})\|}{\|\mathbf{x}-\mathbf{p}\|}=0.$$

Remark 4.3.1: D_f is the "best linear approximation" to f.

Definition 4.3.2 (Jacobian).

When f is differentiable, D_f can be given in coordinates by

$$(D_f)_{ij} = \frac{\partial f_i}{\partial x_i}$$

This yields the **Jacobian** of f:

$$D_{f}(p) \begin{bmatrix} | & | & | \\ \nabla f_{1}(\mathbf{p}) & \nabla f_{2}(\mathbf{p}) & \cdots & \nabla f_{m}(\mathbf{p}) \\ | & | & | \end{bmatrix}^{T} = \begin{bmatrix} \frac{\partial f_{1}}{\partial x_{1}}(\mathbf{p}) & \frac{\partial f_{1}}{\partial x_{2}}(\mathbf{p}) & \cdots & \frac{\partial f_{1}}{\partial x_{n}}(\mathbf{p}) \\ \frac{\partial f_{2}}{\partial x_{1}}(\mathbf{p}) & \frac{\partial f_{2}}{\partial x_{2}}(\mathbf{p}) & \cdots & \frac{\partial f_{2}}{\partial x_{n}}(\mathbf{p}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_{m}}{\partial x_{1}}(\mathbf{p}) & \frac{\partial f_{m}}{\partial x_{2}}(\mathbf{p}) & \cdots & \frac{\partial f_{m}}{\partial x_{n}}(\mathbf{p}) \end{bmatrix}.$$

Remark 4.3.2 : This is equivalent to

4.2 Partial Derivatives 26

- Taking the gradient of each component f_i of f,
- Evaluating ∇f_i at \mathbf{p} ,
- Forming a matrix using these as the columns, and
- Transposing the resulting matrix.

Definition 4.3.3 (Hessian).

For a function $f: \mathbb{R}^n \to \mathbb{R}$, the **Hessian** is a generalization of the second derivative, and is given in coordinates by

$$(H_f)_{ij} = \frac{\partial^2 f}{\partial x_i x_j}$$


Explicitly, we have

$$H_{f}(\mathbf{p}) = \begin{bmatrix} | & | & | \\ D\nabla f_{1}(\mathbf{p}) & D\nabla f_{2}(\mathbf{p}) & \cdots & D\nabla f_{m}(\mathbf{p}) \end{bmatrix}^{T} = \begin{bmatrix} \frac{\partial^{2} f}{\partial x_{1} \partial x_{1}}(\mathbf{a}) & \cdots & \frac{\partial^{2} f}{\partial x_{1} \partial x_{n}}(\mathbf{a}) \\ \vdots & \ddots & \vdots \\ \frac{\partial^{2} f}{\partial x_{n} \partial x_{1}}(\mathbf{a}) & \cdots & \frac{\partial^{2} f}{\partial x_{n} \partial x_{n}}(\mathbf{a}) \end{bmatrix}.$$

Remark 4.3.3: Mnemonic: make matrix with ∇f as the columns, and then differentiate variables left to right.

4.4 The Chain Rule

Example 4.4.1 (How to expand a partial derivative): Write out tree of dependent variables:



4.4 The Chain Rule 27

Δ

Then sum each possible path.

Let subscripts denote which variables are held constant, then

$$\begin{split} \left(\frac{\partial z}{\partial x}\right)_y &= \left(\frac{\partial z}{\partial x}\right)_{u,y,v} \\ &+ \left(\frac{\partial z}{\partial v}\right)_{x,y,u} \left(\frac{\partial v}{\partial x}\right)_y \\ &+ \left(\frac{\partial z}{\partial u}\right)_{x,y,v} \left(\frac{\partial u}{\partial x}\right)_{v,y} \\ &+ \left(\frac{\partial z}{\partial u}\right)_{x,y,v} \left(\frac{\partial u}{\partial v}\right)_{x,y} \left(\frac{\partial v}{\partial x}\right)_y \end{split}$$

4.5 Approximation

Let z = f(x, y), then to approximate near $\mathbf{p}_0 = [x_0, y_0]$,

$$f(\mathbf{x}) \approx f(\mathbf{p}) + \nabla f(\mathbf{x} - \mathbf{p}_0)$$

$$\implies f(x, y) \approx f(\mathbf{p}) + f_x(\mathbf{p})(x - x_0) + f_y(\mathbf{p})(y - y_0)$$

4.6 Optimization

4.6.1 Classifying Critical Points

Definition 4.6.1 (Critical Points).

Critical points of f given by points \mathbf{p} such that the derivative vanishes:

$$\operatorname{crit}(f) = \left\{ \mathbf{p} \in \mathbb{R}^n \mid D_f(\mathbf{p}) = 0 \right\}$$

Proposition 4.6.1 (Second Derivative Test).

1. Compute

$$|H_f(\mathbf{p})| \coloneqq \left| egin{array}{cc} f_{xx} & f_{xy} \ f_{yx} & f_{yy} \end{array} \right| (\mathbf{p})$$

- 2. Check by cases:
 - $|H(\mathbf{p})| = 0$: No conclusion
 - $|H(\mathbf{p})| < 0$: Saddle point
 - $|H(\mathbf{p})| > 0$:

$$-f_{xx}(\mathbf{p}) > 0 \implies \text{local min}$$

 $-f_{xx}(\mathbf{p}) < 0 \implies \text{local max}$

Remark 4.6.1: What's really going on?

- Eigenvalues have same sign \iff positive definite or negative definite
 - Positive definite \implies convex \implies local min
 - Negative definite \implies concave \implies local max
- Extrema occur on boundaries, so parameterize each boundary to obtain a function in one less variable and apply standard optimization techniques to yield critical points. Test all critical points to find extrema.
- If possible, use constraint to just reduce equation to one dimension and optimze like single-variable case.

Add examples

4.6.2 Lagrange Multipliers

The setup:

Optimize
$$f(\mathbf{x})$$
 subject to $g(\mathbf{x}) = c$
 $\implies \nabla f = \lambda \nabla g$

- 1. Use this formula to obtain a system of equations in the components of x and the parameter λ .
 - 2. Use λ to obtain a relation involving only components of **x**.
 - 3. Substitute relations back into constraint to obtain a collection of critical points.
 - 4. Evaluate f at critical points to find max/min.

Add examples

4.7 Change of Variables



For any $f: \mathbb{R}^n \to \mathbb{R}^n$ and region R,

$$\int_{g(R)} f(\mathbf{x}) \ dV = \int_{R} (f \circ g)(\mathbf{x}) \cdot |D_g(\mathbf{x})| \ dV$$

4.7 Change of Variables

5 | Vector Calculus

5.1 Notation

R is a region, S is a surface, V is a solid.

$$\oint_{\partial S} \mathbf{F} \cdot d\mathbf{r} = \oint_{\partial S} [\mathbf{F}_1, \mathbf{F}_2, \mathbf{F}_3] \cdot [dx, dy, dz] = \oint_{\partial S} \mathbf{F}_1 dx + \mathbf{F}_2 dy + \mathbf{F}_3 dz$$

The main vector operators

$$\nabla : (\mathbb{R}^n \to \mathbb{R}) \to (\mathbb{R}^n \to \mathbb{R}^n)$$
$$\varphi \mapsto \nabla \varphi := \sum_{i=1}^n \frac{\partial \varphi}{\partial x_i} \mathbf{e}_i$$

$$\operatorname{div}(\mathbf{F}) : (\mathbb{R}^n \to \mathbb{R}^n) \to (\mathbb{R}^n \to \mathbb{R})$$
$$\mathbf{F} \mapsto \nabla \cdot \mathbf{F} := \sum_{i=1}^n \frac{\partial \mathbf{F}_i}{\partial x_i}$$

$$\mathrm{curl}(\mathbf{F}): (\mathbb{R}^3 \to \mathbb{R}^3) \to (\mathbb{R}^3 \to \mathbb{R}^3)$$
$$\mathbf{F} \mapsto \nabla \times \mathbf{F}$$

Some terminology:

Scalar Field $\varphi: X \to \mathbb{R}$ Vector Field $\mathbf{F}: X \to \mathbb{R}^n$

Gradient Field $\mathbf{F}:X\to\mathbb{R}^n\ \Big|\ \exists \varphi:X\to\mathbb{R}\ \Big|\ \nabla\varphi=F$

- The Gradient: lifts scalar fields on \mathbb{R}^n to vector fields on \mathbb{R}^n
- Divergence: drops vector fields on \mathbb{R}^n to scalar fields on \mathbb{R}^n
- Curl: takes vector fields on \mathbb{R}^3 to vector fields on \mathbb{R}^3

5 Vector Calculus

$$\mathbf{x} \cdot \mathbf{y} = \langle \mathbf{x}, \ \mathbf{y} \rangle = \sum_{i=1}^{n} x_i y_i = x_1 y_1 + x_2 y_2 + \cdots \qquad \text{inner/dot product}$$

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \ \mathbf{x} \rangle} = \sqrt{\sum_{i=1}^{n} x_i^2} = \sqrt{x_1^2 + x_2^2 + \cdots} \qquad \text{norm}$$

$$\mathbf{a} \times \mathbf{b} = \hat{\mathbf{n}} \|\mathbf{a}\| \|\mathbf{b}\| \sin \theta_{\mathbf{a}, \mathbf{b}} = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \qquad \text{cross product}$$

$$D_{\mathbf{u}}(\varphi) = \nabla \varphi \cdot \hat{\mathbf{u}}$$
 directional derivative

$$\nabla \coloneqq \sum_{i=1}^{n} \frac{\partial}{\partial x_i} \mathbf{e}_i = \left[\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \cdots, \frac{\partial}{\partial x_n} \right]$$
 del operator

$$\nabla \varphi := \sum_{i=1}^{n} \frac{\partial \varphi}{\partial x_i} \ \mathbf{e}_i = \left[\frac{\partial \varphi}{\partial x_1}, \frac{\partial \varphi}{\partial x_2}, \cdots, \frac{\partial \varphi}{\partial x_n} \right]$$
gradient

$$\Delta \varphi := \nabla \cdot \nabla \varphi := \sum_{i=1}^{n} \frac{\partial^{2} \varphi}{\partial x_{i}^{2}} = \frac{\partial^{2} \varphi}{\partial x_{1}^{2}} + \frac{\partial^{2} \varphi}{\partial x_{2}} + \dots + \frac{\partial^{2} \varphi}{\partial x_{n}^{2}}$$
 Laplacian

$$\nabla \cdot \mathbf{F} := \sum_{i=1}^{n} \frac{\partial \mathbf{F}_{i}}{\partial x_{i}} = \frac{\partial \mathbf{F}_{1}}{\partial x_{1}} + \frac{\partial \mathbf{F}_{2}}{\partial x_{2}} + \dots + \frac{\partial \mathbf{F}_{n}}{\partial x_{n}}$$
 divergence

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{e}_{1} & \mathbf{e}_{2} & \mathbf{e}_{3} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \mathbf{F}_{1} & \mathbf{F}_{2} & \mathbf{F}_{3} \end{vmatrix} = [\mathbf{F}_{3y} - \mathbf{F}_{2z}, \mathbf{F}_{1z} - \mathbf{F}_{3x}, \mathbf{F}_{2x} - \mathbf{F}_{1y}]$$
 curl
$$\iint_{\mathcal{C}} (\nabla \times \mathbf{F}) \cdot d\mathbf{S} = \iint_{\mathcal{C}} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \ dS$$
 surface integral

5.2 Big Theorems

5.2.1 Stokes' and Consequences

5.2 Big Theorems 31

Theorem 5.2.1 (Stokes' Theorem).

$$\oint_{\partial S} \mathbf{F} \cdot d\mathbf{r} = \iint_{S} (\nabla \times \mathbf{F}) \cdot d\mathbf{S}.$$

Remark 5.2.1: Note that if S is a closed surface, so $\partial S = \emptyset$, this integral vanishes.

Corollary 5.2.1 (Green's Theorem).

$$\oint_{\partial R} (L \ dx + M \ dy) = \iint_{R} \left(\frac{\partial M}{\partial x} - \frac{\partial L}{\partial y} \right) dx dy.$$

Proof(?).

Recovering Green's Theorem from Stokes' Theorem:

Let
$$\mathbf{F} = [L, M, 0]$$
, then $\nabla \times \mathbf{F} = [0, 0, \frac{\partial M}{\partial x} - \frac{\partial L}{\partial y}]$

Corollary 5.2.2 (Divergence Theorem).

$$\iint_{\partial V} \mathbf{F} \cdot d\mathbf{S} = \iiint_{V} (\nabla \cdot \mathbf{F}) \ dV.$$

Remark 5.2.2 :

- $\nabla \times (\nabla \varphi) = 0$
- $\nabla \cdot (\nabla \times \mathbf{F}) = 0$

5.2.2 Directional Derivatives

Definition 5.2.1 (Directional Derivative).

$$D_{\mathbf{v}}f(\mathbf{p}) \coloneqq \frac{\partial f}{\partial t} \left(\mathbf{p} + t\mathbf{v} \right) \Big|_{t=0}.$$

Remark 5.2.3: Note that the directional derivative uses a normalized direction vector!

Theorem 5.2.2 (Dot product expression of directional derivative).

Suppose $f: \mathbb{R}^n \to \mathbb{R}$ and $\mathbf{v} \in \mathbb{R}^n$. Then

$$D_{\mathbf{v}}f(\mathbf{p}) = \langle \nabla f(\mathbf{p}), \mathbf{v} \rangle.$$

5.2 Big Theorems 32

Proof (?).

We first use the fact that we can find L, the best linear approximation to f:

$$L(\mathbf{x}) := f(\mathbf{p}) + D_f(\mathbf{p})(\mathbf{x} - \mathbf{p})$$

$$D_{\mathbf{v}}f(\mathbf{p}) = D_{\mathbf{v}}L(\mathbf{p})$$

$$= \lim_{t \to 0} \frac{L(\mathbf{p} + t\mathbf{v}) - L(\mathbf{p})}{t}$$

$$= \lim_{t \to 0} \frac{f(\mathbf{p}) + D_f(\mathbf{p})(\mathbf{p} + t\mathbf{v} - \mathbf{p}) - (f(\mathbf{p}) + D_f(\mathbf{p})(\mathbf{p} - \mathbf{p}))}{t}$$

$$= \lim_{t \to 0} \frac{D_f(\mathbf{p})(t\mathbf{v})}{t}$$

$$= D_f(\mathbf{p})\mathbf{v}$$

$$\coloneqq \nabla f(\mathbf{p}) \cdot \mathbf{v}.$$

Need a better proof, not clear that this works.

5.3 Computing Integrals

~

5.3.1 Changing Coordinates

Multivariable Chain Rule

todo

Polar and Cylindrical Coordinates

$$x = r \cos \theta$$
$$y = r \sin \theta$$
$$dV \mapsto r - dr \ d\theta$$

Spherical Coordinates

$$x = r \cos \theta = \rho \sin \varphi \cos \theta$$
$$y = r \sin \theta = \rho \sin \varphi \sin \theta$$
$$dV \mapsto r^2 \sin \varphi \quad dr \, d\varphi \, d\theta$$

5.3 Computing Integrals

Vector Calculus

5.3.2 Line Integrals

Curves

• Parametrize the path C as $\{\mathbf{r}(t): t \in [a,b]\}$, then

$$\int_{C} f \ ds := \int_{a}^{b} (f \circ \mathbf{r})(t) \|\mathbf{r}'(t)\| \ dt$$
$$= \int_{a}^{b} f(x(t), y(t), z(t)) \sqrt{x_{t}^{2} + y_{t}^{2} + z_{t}^{2}} \ dt$$

Vector Fields

• If exact:

$$\frac{\partial}{\partial y} \mathbf{F_1} = \frac{\partial}{\partial x} \mathbf{F_2} \implies \int \mathbf{F_1} \ dx + \mathbf{F_2} \ dy = \varphi(\mathbf{p_1}) - \varphi(\mathbf{p_0})$$

The function φ can be found using the same method from ODEs.

• Parametrize the path C as $\{\mathbf{r}(t): t \in [a,b]\}$, then

$$\int_{C} \mathbf{F} \cdot d\mathbf{r} := \int_{a}^{b} (\mathbf{F} \circ \mathbf{r})(t) \cdot \mathbf{r}'(t) dt$$

$$= \int_{a}^{b} [\mathbf{F}_{1}(x(t), y(t), \cdots), \mathbf{F}_{2}(x(t), y(t), \cdots)] \cdot [x_{t}, y_{t}, \cdots] dt$$

$$= \int_{a}^{b} \mathbf{F}_{1}(x(t), y(t), \cdots) x_{t} + \mathbf{F}_{2}(x(t), y(t), \cdots) y_{t} + \cdots dt$$

• Equivalently written:

$$\int_a^b \mathbf{F}_1 \ dx + \mathbf{F}_2 \ dy + \dots := \int_C \mathbf{F} \cdot d\mathbf{r}$$

in which case $[dx, dy, \cdots] := [x_t, y_t, \cdots] = \mathbf{r}'(t)$.

• Remember to substitute dx back into the integrand!!

5.3.3 Flux

$$\iint_{S} \mathbf{F} \cdot d\mathbf{S} = \iint_{S} \mathbf{F} \cdot \hat{\mathbf{n}} \ dS.$$

5.3.4 Area

Proposition 5.3.1(Areas can be computed with Green's Theorem). Given R and f(x, y) = 0,

$$A(R) = \oint_{\partial R} x \ dy = -\oint_{\partial R} y \ dx = \frac{1}{2} \oint_{\partial R} -y \ dx + x \ dy.$$

Proof (?). Compute

$$\oint_{\partial R} x \ dy = -\oint_{\partial R} y \ dx$$
$$= \frac{1}{2} \oint_{\partial R} -y \ dx + x \ dy = \frac{1}{2} \iint_{R} 1 - (-1) \ dA = \iint_{R} 1 \ dA$$

5.3.5 Surface Integrals

• For a paramterization $\mathbf{r}(s,t):U\to S$ of a surface S and any function $f:\mathbb{R}^n\to\mathbb{R}$,

$$\iint_{S} f \ dA = \iint_{U} (f \circ \mathbf{r})(s, t) \|\mathbf{n}\| \ dA$$

• Can obtain a normal vector $\mathbf{n} = T_u \times T_v$

5.4 Other Results

Example 5.4.1 (?): $\nabla \cdot \mathbf{F} = 0 \implies \exists G: \mathbf{F} = \nabla \times G$. A counterexample

$$\mathbf{F}(x, y, z) = \frac{1}{\sqrt{x^2 + y^2 + z^2}} [x, y, z] , \quad S = S^2 \subset \mathbb{R}^3$$

$$\implies \nabla \mathbf{F} = 0 \text{ but } \iint_{S^2} \mathbf{F} \cdot d\mathbf{S} = 4\pi \neq 0$$

Where by Stokes' theorem,

$$\mathbf{F} = \nabla \times \mathbf{G} \implies \iint_{S^2} \mathbf{F} = \iint_{S^2} \nabla \times \mathbf{G}$$

$$= \oint_{\partial S^2} \mathbf{G} \ d\mathbf{r}$$
 by Stokes
$$= 0$$

5.4 Other Results 35

since $\partial S^2 = \emptyset$.

Proposition 5.4.1 (Sufficient Conditions).


Sufficient condition: if **F** is everywhere C^1 ,

$$\exists \mathbf{G}: \ \mathbf{F} = \nabla \times \mathbf{G} \iff \iint_{S} \mathbf{F} \cdot d\mathbf{S} = 0 \text{ for all closed surfaces } S.$$

6 | Linear Algebra

Remark 6.0.1: The underlying field will be assumed to be \mathbb{R} for this section.

6.1 Notation



6.2 Big Theorems



Theorem 6.2.1 (Rank-Nullity).

$$|\ker(A)| + |\operatorname{im}(A)| = |\operatorname{dom}(A)|,$$

where $\operatorname{nullspace}(A) = |\operatorname{im} A|, \operatorname{rank}(A) = |\operatorname{im}(A)|,$ and n is the number of columns in the corresponding matrix.

Generalization: the following sequence is always exact:

$$0 \to \ker(A) \stackrel{\mathrm{id}}{\hookrightarrow} \mathrm{dom}(A) \xrightarrow{A} \mathrm{im}(A) \to 0.$$

Moreover, it always splits, so dom $A = \ker A \oplus \operatorname{im} A$ and thus $|\operatorname{dom}(A)| = |\ker(A)| + |\operatorname{im}(A)|$.

Remark 6.2.1: We also have

$$\dim(\operatorname{rowspace}(A)) = \dim(\operatorname{colspace}(A)) = \operatorname{rank}(A).$$

6.3 Big List of Equivalent Properties

Let A be an $m \times n$ matrix. TFAE: - A is invertible and has a unique inverse A^{-1} - A^T is invertible - $\det(A) \neq 0$ - The linear system $A\mathbf{x} = \mathbf{b}$ has a unique solution for every $b \in \mathbb{R}^m$ - The homogeneous system $A\mathbf{x} = 0$ has only the trivial solution $\mathbf{x} = 0$ - $\operatorname{rank}(A) = n$ - i.e. A is full rank - $\operatorname{nullity}(A) := \dim \operatorname{nullspace}(A) = 0$ - $A = \prod_{i=1}^k E_i$ for some finite k, where each E_i is an elementary matrix. - A is row-equivalent to the identity matrix I_n - A has exactly n pivots - The columns of A are a basis for \mathbb{R}^n - i.e. $\operatorname{colspace}(A) = \mathbb{R}^n$ - The rows of A are a basis for \mathbb{R}^m - i.e. $\operatorname{rowspace}(A) = \mathbb{R}^n$ - Zero is not an eigenvalue of A. - A has n linearly independent eigenvectors - The rows of A are coplanar.

Similarly, by taking negations, TFAE:

- A is not invertible
- A is singular
- A^T is not invertible
- $\det A = 0$
- The linear system $A\mathbf{x} = \mathbf{b}$ has either no solution or infinitely many solutions.
- The homogeneous system $A\mathbf{x} = \mathbf{0}$ has nontrivial solutions
- $\operatorname{rank} A < n$
- dim nullspace A > 0
- At least one row of A is a linear combination of the others
- The RREF of A has a row of all zeros.

6.2 Big Theorems 37

Reformulated in terms of linear maps T, TFAE: - T^{-1} : $\mathbb{R}^m \to \mathbb{R}^n$ exists - $\operatorname{im}(T) = \mathbb{R}^n$ - $\ker(T) = 0$ - T is injective - T is surjective - T is an isomorphism - The system $A\mathbf{x} = 0$ has infinitely many solutions

6.4 Vector Spaces



6.4.1 Linear Transformations

Definition 6.4.1 (Linear Transformation).

todo

Remark 6.4.1: It is common to want to know the range and kernel of a specific linear transformation T. T can be given in many ways, but a general strategy for deducing these properties involves:

- Express an arbitrary vector in V as a linear combination of its basis vectors, and set it equal to an arbitrary vector in W.
- \bullet Use the linear properties of T to make a substitution from known transformations
- Find a restriction or relation given by the constants of the initial linear combination.

Remark 6.4.2: Useful fact: if $V \leq W$ is a subspace and $\dim(V) \geq \dim(W)$, then V = W.

Definition 6.4.2 (Kernel).

todo

Proposition 6.4.1(Two-step vector subspace test).

If $V \subseteq W$, then V is a subspace of W if the following hold:

$$\mathbf{0} \in V$$

(2)
$$\mathbf{a}, \mathbf{b} \in V \implies t\mathbf{a} + \mathbf{b} \in V.$$

6.4.2 Linear Independence

Proposition 6.4.2(?).

Any set of two vectors $\{\mathbf{v}, \mathbf{w}\}$ is linearly **dependent** $\iff \exists \lambda : \mathbf{v} = \lambda \mathbf{w}$, i.e. one is not a scalar multiple of the other.

6.4.3 Bases

Definition 6.4.3 (Basis and dimension).

A set S forms a **basis** for a vector space V iff

- 1. S is a set of linearly independent vectors, so $\sum \alpha_i \vec{s_i} = 0 \implies \alpha_i = 0$ for all i.
- 2. S spans V, so $\vec{v} \in V$ implies there exist α_i such that $\sum \alpha_i \vec{s_i} = \vec{v}$

In this case, we define the **dimension** of V to be |S|.

Show how to compute basis of kernel.

Show how to compute basis of row space (nonzero rows in ??A)?)

Show how to compute basis of column space: leading ones

6.4.4 The Inner Product

The point of this section is to show how an inner product can induce a notion of "angle", which agrees with our intuition in Euclidean spaces such as \mathbb{R}^n , but can be extended to much less intuitive things, like spaces of functions.

Definition 6.4.4 (The standard inner product).

The Euclidean inner product is defined as

$$\langle \mathbf{a}, \ \mathbf{b} \rangle = \sum_{i=1}^{n} a_i b_i = a_1 b_1 + a_2 b_2 + \dots + a_n b_n.$$

Also sometimes written as $\mathbf{a}^T \mathbf{b}$ or $\mathbf{a} \cdot \mathbf{b}$.

Proposition 6.4.3 (Inner products induce norms and angles).

Yields a norm

$$\|\mathbf{x}\| \coloneqq \sqrt{\langle \mathbf{x}, \ \mathbf{x} \rangle}$$

which has a useful alternative formulation

$$\langle \mathbf{x}, \mathbf{x} \rangle = \|\mathbf{x}\|^2.$$

This leads to a notion of angle:

$$\langle \mathbf{x}, \ \mathbf{y} \rangle = \|\mathbf{x}\| \|\mathbf{y}\| \cos \theta_{x,y} \implies \cos \theta_{x,y} := \frac{\langle \mathbf{x}, \ \mathbf{y} \rangle}{\|\mathbf{x}\| \|\mathbf{y}\|} = \langle \widehat{\mathbf{x}}, \ \widehat{\mathbf{y}} \rangle$$

where $\theta_{x,y}$ denotes the angle between the vectors **x** and **y**.

Remark 6.4.3: Since $\cos \theta = 0$ exactly when $\theta = \pm \frac{\pi}{2}$, we can can declare two vectors to be **orthogonal** exactly in this case:

$$\mathbf{x} \in \mathbf{y}^{\perp} \iff \langle \mathbf{x}, \ \mathbf{y} \rangle = 0.$$

Note that this makes the zero vector orthogonal to everything.

Definition 6.4.5 (Orthogonal Complement/Perp).

Given a subspace $S \subseteq V$, we define its **orthogonal complement**

$$S^{\perp} = \left\{ \mathbf{v} \in V \mid \forall \mathbf{s} \in S, \ \langle \mathbf{v}, \ \mathbf{s} \rangle = 0 \right\}.$$

Remark 6.4.4 : Any choice of subspace $S \subseteq V$ yields a decomposition $V = S \oplus S^{\perp}$.

Proposition 6.4.4 (Formula expanding a norm and 'Pythagorean theorem').

A useful formula is

$$\|\mathbf{x} + \mathbf{y}\|^2 = \|\mathbf{x}\|^2 + 2\langle \mathbf{x}, \mathbf{y} \rangle + \|\mathbf{y}\|^2,$$

When $\mathbf{x} \in \mathbf{y}^{\perp}$, this reduces to

$$\|\mathbf{x} + \mathbf{y}\|^2 = \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2.$$

Proposition 6.4.5 (Properties of the inner product).

1. Bilinearity:

$$\left\langle \sum_{j} \alpha_{j} \mathbf{a}_{j}, \sum_{k} \beta_{k} \mathbf{b}_{k} \right\rangle = \sum_{j} \sum_{i} \alpha_{j} \beta_{i} \langle \mathbf{a}_{j}, \mathbf{b}_{i} \rangle.$$

2. Symmetry:

$$\langle \mathbf{a},\ \mathbf{b}\rangle = \langle \mathbf{b},\ \mathbf{a}\rangle$$

3. Positivity:

$$\mathbf{a} \neq \mathbf{0} \implies \langle \mathbf{a}, \ \mathbf{a} \rangle > 0$$

4. Nondegeneracy:

$$\mathbf{a} = \mathbf{0} \iff \langle \mathbf{a}, \ \mathbf{a} \rangle = 0$$

Proof of Cauchy-Schwarz: See Goode page 346

6.4.5 Gram-Schmidt Process

Extending a basis $\{\mathbf{x}_i\}$ to an orthonormal basis $\{\mathbf{u}_i\}$

$$\mathbf{u}_{1} = N(\mathbf{x}_{1})$$

$$\mathbf{u}_{2} = N(\mathbf{x}_{2} - \langle \mathbf{x}_{2}, \mathbf{u}_{1} \rangle \mathbf{u}_{1})$$

$$\mathbf{u}_{3} = N(\mathbf{x}_{3} - \langle \mathbf{x}_{3}, \mathbf{u}_{1} \rangle \mathbf{u}_{1} - \langle \mathbf{x}_{3}, \mathbf{u}_{2} \rangle \mathbf{u}_{2})$$

$$\vdots \qquad \vdots$$

$$\mathbf{u}_{k} = N(\mathbf{x}_{k} - \sum_{i=1}^{k-1} \langle \mathbf{x}_{k}, \mathbf{u}_{i} \rangle \mathbf{u}_{i})$$

where N denotes normalizing the result.

In more detail The general setup here is that we are given an orthogonal basis $\{\mathbf{x}_i\}_{i=1}^n$ and we want to produce an **orthonormal** basis from them.

Why would we want such a thing? Recall that we often wanted to change from the standard basis \mathcal{E} to some different basis $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \cdots\}$. We could form the change of basis matrix $B = [\mathbf{b}_1, \mathbf{b}_2, \cdots]$ acts on vectors in the \mathcal{B} basis according to

$$B[\mathbf{x}]_{\mathcal{B}} = [\mathbf{x}]_{\mathcal{E}}.$$

But to change from \mathcal{E} to \mathcal{B} requires computing B^{-1} , which acts on vectors in the standard basis according to

$$B^{-1}[\mathbf{x}]_{\mathcal{E}} = [\mathbf{x}]_{\mathcal{B}}.$$

If, on the other hand, the \mathbf{b}_i are orthonormal, then $B^{-1} = B^T$, which is much easier to compute. We also obtain a rather simple formula for the coordinates of \mathbf{x} with respect to \mathcal{B} . This follows because we can write

$$\mathbf{x} = \sum_{i=1}^{n} \langle \mathbf{x}, \mathbf{b}_i \rangle \mathbf{b}_i \coloneqq \sum_{i=1}^{n} c_i \mathbf{b}_i,$$

and we find that

$$[\mathbf{x}]_{\mathcal{B}} = \mathbf{c} := [c_1, c_2, \cdots, c_n]^T..$$

This also allows us to simplify projection matrices. Supposing that A has orthonormal columns and letting S be the column space of A, recall that the projection onto S is defined by

$$P_S = Q(Q^T Q)^{-1} Q^T.$$

Since Q has orthogonal columns and satisfies $Q^TQ = I$, this simplifies to

$$P_S = QQ^T$$
..

The Algorithm Given the orthogonal basis $\{\mathbf{x}_i\}$, we form an orthonormal basis $\{\mathbf{u}_i\}$ iteratively as follows.

First define

$$N: \mathbb{R}^n \to S^{n-1}$$

 $\mathbf{x} \mapsto \widehat{\mathbf{x}} := \frac{\mathbf{x}}{\|\mathbf{x}\|}$

which projects a vector onto the unit sphere in \mathbb{R}^n by normalizing. Then,

$$\mathbf{u}_{1} = N(\mathbf{x}_{1})$$

$$\mathbf{u}_{2} = N(\mathbf{x}_{2} - \langle \mathbf{x}_{2}, \mathbf{u}_{1} \rangle \mathbf{u}_{1})$$

$$\mathbf{u}_{3} = N(\mathbf{x}_{3} - \langle \mathbf{x}_{3}, \mathbf{u}_{1} \rangle \mathbf{u}_{1} - \langle \mathbf{x}_{3}, \mathbf{u}_{2} \rangle \mathbf{u}_{2})$$

$$\vdots$$

$$\mathbf{u}_{k} = N(\mathbf{x}_{k} - \sum_{i=1}^{k-1} \langle \mathbf{x}_{k}, \mathbf{u}_{i} \rangle \mathbf{u}_{i})$$

In words, at each stage, we take one of the original vectors \mathbf{x}_i , then subtract off its projections onto all of the \mathbf{u}_i we've created up until that point. This leaves us with only the component of \mathbf{x}_i that is orthogonal to the span of the previous \mathbf{u}_i we already have, and we then normalize each \mathbf{u}_i we obtain this way.

Alternative Explanation:

Given a basis

$$S = \{\mathbf{v_1}, \mathbf{v_2}, \cdots \mathbf{v_n}\},\,$$

the Gram-Schmidt process produces a corresponding orthogonal basis

$$S' = \{\mathbf{u_1}, \mathbf{u_2}, \cdots \mathbf{u_n}\}$$

that spans the same vector space as S.

S' is found using the following pattern:

$$\begin{aligned} \mathbf{u}_1 &= \mathbf{v}_1 \\ \mathbf{u}_2 &= \mathbf{v}_2 - \mathrm{proj}_{\mathbf{u}_1} \mathbf{v}_2 \\ \mathbf{u}_3 &= \mathbf{v}_3 - \mathrm{proj}_{\mathbf{u}_1} \mathbf{v}_3 - \mathrm{proj}_{\mathbf{u}_2} \mathbf{v}_3 \end{aligned}$$

where

$$\mathrm{proj}_{\mathbf{u}}\mathbf{v} = (\mathrm{scal}_{\mathbf{u}}\mathbf{v})\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{\langle \mathbf{v}, \mathbf{u} \rangle}{\|\mathbf{u}\|} \frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{\langle \mathbf{v}, \ \mathbf{u} \rangle}{\|\mathbf{u}\|^2}\mathbf{u}$$

is a vector defined as the orthogonal projection of \mathbf{v} onto \mathbf{u} .




Figure 1: Image

The orthogonal set S' can then be transformed into an orthonormal set S'' by simply dividing the vectors $s \in S'$ by their magnitudes. The usual definition of a vector's magnitude is

$$\|\mathbf{a}\| = \sqrt{\langle \mathbf{a}, \ \mathbf{a} \rangle} \text{ and } \|\mathbf{a}\|^2 = \langle \mathbf{a}, \ \mathbf{a} \rangle$$

As a final check, all vectors in S' should be orthogonal to each other, such that

$$\langle \mathbf{v_i}, \ \mathbf{v_j} \rangle = 0 \text{ when } i \neq j$$

and all vectors in S'' should be orthonormal, such that

$$\langle \mathbf{v_i}, \ \mathbf{v_i} \rangle = \delta_{ij}$$

6.4.6 The Fundamental Subspaces Theorem

Given a matrix $A \in Mat(m, n)$, and noting that

$$A: \mathbb{R}^n \to \mathbb{R}^m,$$
$$A^T: \mathbb{R}^m \to \mathbb{R}^n$$

We have the following decompositions:


$$\mathbb{R}^n \qquad \cong \ker A \oplus \operatorname{im} A^T \qquad \cong \operatorname{nullspace}(A) \oplus \operatorname{colspace}(A^T)$$

$$\mathbb{R}^m \qquad \cong \operatorname{im} A \oplus \ker A^T \qquad \cong \operatorname{colspace}(A) \oplus \operatorname{nullspace}(A^T)$$

6.4.7 Computing change of basis matrices

todo

6.5 Matrices



Marning 6.1: The space of matrices is not an integral domain! Counterexample: if A is singular and nonzero, there is some nonzero \mathbf{v} such that $A\mathbf{v} = \mathbf{0}$. Then setting $B = [\mathbf{v}, \mathbf{v}, \cdots]$ yields AB = 0 with $A \neq 0, B \neq 0$.

Definition 6.5.1 (Rank of a matrix).

The **rank** of a matrix A representing a linear transformation T is dim colspace (A), or equivalently dim im T.

Proposition 6.5.1(?).

rank(A) is equal to the number of nonzero rows in RREF(A).

Definition 6.5.2 (Trace of a Matrix).

$$\operatorname{Trace}(A) = \sum_{i=1}^{m} A_{ii}$$

6.5 Matrices 44

Definition 6.5.3 (Elementary Row Operations).

The following are **elementary row operations** on a matrix:

- Permute rows
- Multiple a row by a scalar
- Add any row to another

Proposition 6.5.2 (Formula for matrix multiplication).

If $A = [\mathbf{a}_1, \mathbf{a}_2, \cdots] \in \operatorname{Mat}(m, n)$ and $B = [\mathbf{b}_1, \mathbf{b}_2, \cdots] \in \operatorname{Mat}(n, p)$, then

$$C := AB \implies c_{ij} = \sum_{k=1}^{n} a_{ik} b_{kj} = \langle \mathbf{a_i}, \ \mathbf{b_j} \rangle$$

where $1 \le i \le m$ and $1 \le j \le p$. In words, each entry c_{ij} is obtained by dotting row i of A against column j of B.

6.5.1 Systems of Linear Equations

Definition 6.5.4 (Consistent and inconsistent).

A system of linear equations is **consistent** when it has at least one solution. The system is **inconsistent** when it has no solutions.

Definition 6.5.5 (Homogeneous Systems).

Remark 6.5.2: Homogeneous systems are always consistent, i.e. there is always at least one solution.

Remark 6.5.3:

- Tall matrices: more equations than unknowns, overdetermined
- Wide matrices: more unknowns than equations, underdetermined

Proposition 6.5.3 (Characterizing solutions to a system of linear equations).

There are three possibilities for a system of linear equations:


- 1. No solutions (inconsistent)
- 2. One unique solution (consistent, square or tall matrices)
- 3. Infinitely many solutions (consistent, underdetermined, square or wide matrices)


6.5 Matrices 45


Unique solution

No solutions


Infinitely many solutions







$$3x + 2y = 7
3x + 2y = 4$$



$$3x + 2y = 7
6x + 4y = 14$$

These possibilities can be check by considering $r := \operatorname{rank}(A)$:

- $r < r_b$: case 1, no solutions.
- $r = r_b$: case 1 or 2, at least one solution.
 - $-r_b=n$: case 2, a unique solution.
 - $-r_b < n$: case 3, infinitely many solutions.

6.5.2 Determinants

Proposition 6.5.4(?).

 $\det(A \mod p) \mod p \equiv (\det A) \mod p$

Proposition 6.5.5 (Inverse of a 2×2 matrix).

For 2×2 matrices,

$$A^{-1} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{\det A} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

In words, swap the main diagonal entries, and flip the signs on the off-diagonal.

Proposition 6.5.6 (Properties of the determinant).

Let $A \in Mat(m, n)$, then there is a function

$$\det: \operatorname{Mat}(m, m) \to \mathbb{R}$$
$$A \mapsto \det(A)$$

satisfying the following properties:

• det is a group homomorphism onto (\mathbb{R},\cdot) :

$$\det(AB) = \det(A)\det(B)$$

- Some corollaries:

$$\det A^k = k \det A$$

$$\det(A^{-1}) = (\det A)^{-1} \det(A^t)$$
 = \det(A).

• Invariance under adding scalar multiples of any row to another:

$$\det \begin{bmatrix} \vdots \\ -\mathbf{a}_i \\ \vdots \end{bmatrix} = \det \begin{bmatrix} \vdots \\ -\mathbf{a}_i + t\mathbf{a}_j \\ \vdots \end{bmatrix}$$

• Sign change under row permutation:

$$\det \begin{bmatrix} \vdots \\ -\mathbf{a}_i & - \\ \vdots \\ -\mathbf{a}_j & - \end{bmatrix} = (-1) \det \begin{bmatrix} \vdots \\ -\mathbf{a}_j & - \\ \vdots \\ -\mathbf{a}_i & - \end{bmatrix}$$

– More generally, for a permutation $\sigma \in S_n$,

$$\det \begin{bmatrix} \vdots \\ -\mathbf{a}_{i} & - \\ \vdots \\ -\mathbf{a}_{j} & - \\ \vdots \end{bmatrix} = (-1)^{\operatorname{sgn}(\sigma)} \det \begin{bmatrix} \vdots \\ -\mathbf{a}_{\sigma(j)} & - \\ \vdots \\ -\mathbf{a}_{\sigma(i)} & - \\ \vdots \end{bmatrix}$$

• Multilinearity in rows:

$$\det \begin{bmatrix} \vdots \\ -t\mathbf{a}_{i} \\ -t\mathbf{a}_{i} \end{bmatrix} = t \det \begin{bmatrix} \vdots \\ -\mathbf{a}_{i} \\ -t\mathbf{a}_{i} \end{bmatrix}$$

$$\det \begin{bmatrix} -t\mathbf{a}_{1} & -t\mathbf{a}_{2} \\ -t\mathbf{a}_{2} & -t\mathbf{a}_{2} \end{bmatrix} = t^{m} \det \begin{bmatrix} -\mathbf{a}_{1} & -t\mathbf{a}_{2} \\ -t\mathbf{a}_{2} & -t\mathbf{a}_{3} \end{bmatrix}$$

$$\det \begin{bmatrix} -t\mathbf{a}_{1} & -t\mathbf{a}_{2} \\ -t\mathbf{a}_{2} & -t\mathbf{a}_{3} \end{bmatrix} = \prod_{i=1}^{m} t_{i} \det \begin{bmatrix} -\mathbf{a}_{1} & -t\mathbf{a}_{2} \\ -t\mathbf{a}_{2} & -t\mathbf{a}_{3} \end{bmatrix}.$$

$$\det \begin{bmatrix} -t\mathbf{a}_{1} & -t\mathbf{a}_{2} \\ -t\mathbf{a}_{3} & -t\mathbf{a}_{3} \end{bmatrix} = \prod_{i=1}^{m} t_{i} \det \begin{bmatrix} -\mathbf{a}_{1} & -t\mathbf{a}_{3} \\ -t\mathbf{a}_{3} & -t\mathbf{a}_{3} \end{bmatrix}.$$

• Linearity in each row:

$$\det \begin{bmatrix} \vdots \\ -\mathbf{a}_i + \mathbf{a}_j \\ \vdots \end{bmatrix} = \det \begin{bmatrix} \vdots \\ -\mathbf{a}_i \\ \vdots \end{bmatrix} + \det \begin{bmatrix} \vdots \\ -\mathbf{a}_j \\ \vdots \end{bmatrix}.$$

- det(A) is the volume of the parallelepiped spanned by the columns of A.
- If any row of A is all zeros, det(A) = 0.

Proposition 6.5.7 (Characterizing singular matrices).

TFAE:

- $\det(A) = 0$
- A is singular.

6.5.3 Computing Determinants

Useful shortcuts:

• If A is upper or lower triangular, $det(A) = \prod_{i} a_{ii}$.

Definition 6.5.6 (Minors).

The **minor** M_{ij} of $A \in \text{Mat}(n, n)$ is the *determinant* of the $(n-1) \times (n-1)$ matrix obtained by deleting the *i*th row and *j*th column from A.

Definition 6.5.7 (Cofactors).

The **cofactor** C_{ij} is the scalar defined by

$$C_{ij} := (-1)^{i+j} M_{ij}$$
.

Proposition 6.5.8 (Laplace/Cofactor Expansion).

For any fixed i, there is a formula

$$\det(A) = \sum_{i=1}^{n} a_{ij} C_{ij}.$$

Example 6.5.1 (?): Let

$$A = \left[\begin{array}{rrr} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{array} \right].$$

Then

$$\det A = 1 \cdot \left| \begin{array}{cc|c} 5 & 6 \\ 8 & 9 \end{array} \right| - 2 \cdot \left| \begin{array}{cc|c} 4 & 6 \\ 7 & 9 \end{array} \right| + 3 \cdot \left| \begin{array}{cc|c} 4 & 5 \\ 7 & 8 \end{array} \right| = 1 \cdot (-3) - 2 \cdot (-6) + 3 \cdot (-3) = 0.$$

Proposition 6.5.9 (Computing determinant from RREF).

 $\det(A)$ can be computed by reducing A to $\operatorname{RREF}(A)$ (which is upper triangular) and keeping track of the following effects:

- $R_i \leftarrow R_i \pm tR_i$: no effect.
- $R_i \rightleftharpoons R_j$: multiply by (-1).
- $R_i \leftarrow tR_i$: multiply by t.

6.5.4 Inverting a Matrix

Proposition 6.5.10 (Cramer's Rule).

Given a linear system $A\mathbf{x} = \mathbf{b}$, writing $\mathbf{x} = [x_1, \dots, x_n]$, there is a formula

$$x_i = \frac{\det(B_i)}{\det(A)}$$

where B_i is A with the ith column deleted and replaced by **b**.

Proposition 6.5.11 (Gauss-Jordan Method for inverting a matrix).

Under the equivalence relation of elementary row operations, there is an equivalence of augmented matrices:

$$\left[A \mid I\right] \sim \left[I \mid A^{-1}\right]$$

where I is the $n \times n$ identity matrix.

Proposition 6.5.12 (Cofactor formula for inverse).

$$A^{-1} = \frac{1}{\det(A)} [C_{ij}]^t.$$

where C_{ij} is the cofactor(Definition 6.5.7) at position i, j, a

^aNote that the matrix appearing here is sometimes called the *adjugate*.

Example 6.5.2 (Inverting a 2×2 matrix):

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \quad \text{where } ad - bc \neq 0$$

What's the pattern?

- 1. Always divide by determinant
- 2. Swap the diagonals
- 3. Hadamard product with checkerboard

Example 6.5.3 (Inverting a 3×3 matrix):

$$\begin{bmatrix} + & - \\ - & + \end{bmatrix}$$

$$A^{-1} \coloneqq \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}^{-1} = \frac{1}{\det A} \begin{bmatrix} ei-fh & -(bi-ch) & bf-ce \\ -(di-fg) & ai-cg & -(af-cd) \\ dh-eg & -(ah-bg) & ae-bd \end{bmatrix}.$$

The pattern:

- 1. Divide by determinant
- 2. Each entry is determinant of submatrix of A with corresponding col/row deleted
- 3. Hadamard product with checkerboard

4. Transpose at the end!!

6.5.5 Bases for Spaces of a Matrix

Let $A \in \text{Mat}(m, n)$ represent a map $T : \mathbb{R}^n \to \mathbb{R}^m$.

Add examples.

Definition 6.5.8 (Pivot).

todo

Proposition 6.5.13.

 $\dim \operatorname{rowspace}(A) = \dim \operatorname{colspace}(A).$

The row space

$$\operatorname{im}(T)^{\vee} = \operatorname{rowspace}(A) \subset \mathbb{R}^n.$$

Reduce to RREF, and take nonzero rows of RREF(A).

The column space

$$im(T) = colspace(A) \subseteq \mathbb{R}^m$$

Reduce to RREF, and take columns with pivots from original A.

Remark 6.5.4: Not enough pivots implies columns don't span the entire target domain

The nullspace

$$\ker(T) = \operatorname{nullspace}(A) \subseteq \mathbb{R}^n$$

Reduce to RREF, zero rows are free variables, convert back to equations and pull free variables out as scalar multipliers.

Eigenspaces For each $\lambda \in \operatorname{Spec}(A)$, compute a basis for $\ker(A - \lambda I)$.

6.5.6 Eigenvalues and Eigenvectors

Definition 6.5.9 (Eigenvalues, eigenvectors, eigenspaces).

A vector **v** is said to be an **eigenvector** of A with **eigenvalue** $\lambda \in \operatorname{Spec}(A)$ iff

$$A\mathbf{v} = \lambda \mathbf{v}$$

6.5 Matrices 51

For a fixed λ , the corresponding **eigenspace** E_{λ} is the span of all such vectors.

$Remark\ 6.5.5$:

- Similar matrices have identical eigenvalues and multiplicities.
- Eigenvectors corresponding to distinct eigenvalues are always linearly independent
- A has n distinct eigenvalues \implies A has n linearly independent eigenvectors.
- A matrix A is diagonalizable \iff A has n linearly independent eigenvectors.

Proposition $6.5.14 (How\ to\ find\ eigenvectors).$

For $\lambda \in \operatorname{Spec}(A)$,

$$\mathbf{v} \in E_{\lambda} \iff \mathbf{v} \in \ker(A - I\lambda).$$

 $Remark\ 6.5.6$: Some miscellaneous useful facts:

- $\lambda \in \operatorname{Spec}(A) \implies \lambda^2 \in \operatorname{Spec}(A^2)$ with the same eigenvector.
- $\prod \lambda_i = \det A$
- $\sum \lambda_i = \operatorname{Tr} A$

Finding generalized eigenvectors

todo

Diagonalizability

Remark 6.5.7: An $n \times n$ matrix P is diagonalizable iff its eigenspace is all of \mathbb{R}^n (i.e. there are n linearly independent eigenvectors, so they span the space.)

Remark 6.5.8:A is diagonalizable if there is a basis of eigenvectors for the range of P.

6.5 Matrices 52

6.5.7 Useful Counterexamples

$$A := \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \implies A^n = \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}, \qquad \operatorname{Spec}(A) = [1, 1]$$

$$A := \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \implies A^2 = I_2, \qquad \operatorname{Spec}(A) = [1, -1]$$

6.6 Example Problems

Problem. (?)

Determine a basis for

$$S = \left\{ a_0 + a_1 x + a_2 x^2 \mid a_0, a_1, a_2 \in \mathbb{R} \land a_0 - a_1 - 2a_2 = 0 \right\}.$$

Solution:

Let $a_2 = t$, $a_1 = s$, $a_0 = s + 2t$, then

$$S = \left\{ (s+2t) + (sx + tx^2) \mid s, t \in \mathbb{R} \right\}$$

$$= \left\{ (s+sx) + (2t + tx^2) \mid s, t \in \mathbb{R} \right\}$$

$$= \left\{ s(1+x) + t(2+x^2) \mid s, t \in \mathbb{R} \right\}$$

$$= \operatorname{span} \left\{ (1+x), (2+x^2) \right\}$$

and a basis for S is

$$\{(1+x), (2+x^2)\}$$

Problem. (?)

 \mathbf{T}/\mathbf{F} : If V is an n-dimensional vector space, then every set S with fewer than n vectors can be extended to a basis for V.

Solution:

False. Only *linearly independent* sets with fewer than n vectors can be extended to form a basis for V.

6.6 Example Problems 53

Problem. (?)

T/F: The set of all 3 x 3 upper triangular matrices forms a three-dimensional subspace of

Solution:

False. This set forms a 6-dimensional subspace. A basis for this space would require six elements.

Problem. (?)Given A =

$$\begin{bmatrix} 1 & 3' \\ -2 & -6 \end{bmatrix}$$

what is the dimension of the null space of A?

Solution:

The augmented matrix for the system $A\mathbf{x} = \mathbf{0}$ is

$$\begin{bmatrix} [cc|c]1 & 3 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

which has one free variable.

Writing one variable in terms of another results in $x_1 + 3x_2 = 0 \Rightarrow x_1 = 3x_2$.

Let
$$x_2 = r$$
 where $r \in R$, then $S = \{x \in \mathbb{R}^2 : \mathbf{x} = r(3, 1), r \in \mathbb{R}\} = \text{span}\{(3, 1)\}.$

Let $x_2 = r$ where $r \in R$, then $S = \{x \in \mathbb{R}^2 : \mathbf{x} = r(3,1), r \in \mathbb{R}\} = \text{span}\{(3,1)\}.$ So, the set $B = \{(3,1)\}$ is a basis for the null space of A and the dimension of the null space is 1.

Problem. (?)

Let S be the subspace of \mathbb{R}^3 that consists of all solutions to the equation x-3y+z=0. Determine a basis for S, and find $\dim[S]$.

Solution:

The first goal is to find a way to express the set of 3-tuples that satisfy this equation. Let y = r and z = s, then x = r - s. Then vectors **v** that satisfy the equation are all of the form

$$\mathbf{v} = (3r - s, r, s) = (3r, r, 0) + (-s, 0, s) = r(3, 1, 0) + s(-1, 0, 1).$$

(Note - the goal here is to separate the dependent variables into different vectors so they can be written as a linear combination of something.)

The set S that satisfies this equation is then

$$S = \left\{ \mathbf{v} \in \mathbb{R}^3 : \mathbf{v} = r(3, 1, 0) + s(-1, 0, 1) \land r, s \in \mathbb{R} \right\}$$
$$= \operatorname{span} \left\{ (3, 1, 0), (-1, 0, 1) \right\}$$

All that remains is to check that the vectors in this span are linearly independent. This can be done by showing that **if**

$$a(3,1,0) + b(-1,0,1) = (0,0,0)$$

then a = b = 0.

Since the two vectors are linearly independent and span the solution set S, they form a basis for S of dimension 2.

Problem. (?)

Determine a basis for the subspace of $M_2(\mathbb{R})$ spanned by

$$\begin{bmatrix} -1 & 3 \\ -1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} -1 & 4 \\ 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 6 \\ -5 & 1 \end{bmatrix}$$

Solution:

Note that because the set contains the zero matrix, it is linearly dependent. So only consider the other three, as they span the same subspace as the original set.

First, determine if $\{A_1, A_2, A_3\}$ is linearly independent. Start with the equation

$$c_1 A_1 + c_2 A_2 + c_3 A_3 = 0_2$$

which gives

$$c_1 - c_2 + 5c_3 = 0$$
$$3c_1 + 4c_2 - 6c_3 = 0$$
$$-c_1 + c_2 - 5c_3 = 0$$
$$2c_1 + c_2 + c_3 = 0$$

which has the solution (-2r, 3r, r). So the set is linearly dependent by the relation

$$-2A_1 + 3A_2 + A_3 = 0$$
 or $A_3 = 2A_1 - 3A_2$

So $\{A_1, A_2\}$ spans the same subspace as the original set. It is also linearly independent, and therefore forms a basis for the original subspace.

Problem. (?)

Let $A, B, C \in M_2(\mathbb{R})$. Define $\langle A, B \rangle = a_{11}b_{11} + 2a_{12}b_{12} + 3a_{21}b_{21}$. Does this define an inner product on $M_2(\mathbb{R})$?

Problem. (?)

Instead, let $\langle A, B \rangle = a_{11} + b_{22}$. Does this define an inner product on $M_2(\mathbb{R})$?

Problem. (?)

Let $p = a_0 + a_1 x + a_2 x^2$ and $q = b_0 + b_1 x + b_2 x^2$. Define $\langle p, q \rangle = \sum_{i=0}^{2} (i+1)a_i b_i$. Does this define an inner product on P_2 ?

7 Linear Algebra: Advanced Topics

7.1 Changing Basis

Proposition 7.1.1 (Changing to the standard basis).

The transition matrix from a given basis $\mathcal{B} = \{\mathbf{b}_i\}_{i=1}^n$ to the standard basis is given by

$$A \coloneqq \begin{bmatrix} | & | & | \\ \mathbf{b}_1 & \mathbf{b}_2 & \cdots & \mathbf{b}_n \\ | & | & | \end{bmatrix},$$

and the transition matrix from the standard basis to \mathcal{B} is A^{-1} .

7.2 Orthogonal Matrices

Given a notion of orthogonality for vectors, we can extend this to matrices. A square matrix is said to be orthogonal iff $QQ^T = Q^TQ = I$. For rectangular matrices, we have the following characterizations:

$$QQ^T = I \implies$$
 The rows of Q are orthogonal,

$$Q^TQ = I \implies$$
 The columns of Q are orthogonal.

To remember which condition is which, just recall that matrix multiplication AB takes the inner product between the **rows** of A and the **columns** of B. So if, for example, we want to inspect whether or not the columns of Q are orthogonal, we should let B = Q in the above formulation—then we just note that the rows of Q^T are indeed the columns of Q, so Q^TQ computes the inner products between all pairs of the columns of Q and stores them in a matrix.

7.3 Projections

Remark 7.3.1: A projection P induces a decomposition

$$dom(P) = \ker(P) \oplus \ker(P)^{\perp}.$$

Check! Domain or range..?

Distance from a point **p** to a line $\mathbf{a} + t\mathbf{b}$: let $\mathbf{w} = \mathbf{p} - \mathbf{a}$, then: $\|\mathbf{w} - P(\mathbf{w}, \mathbf{v})\|$

Proposition 7.3.1 (Projection onto range).

$$\operatorname{Proj}_{\operatorname{range}(A)}(\mathbf{x}) = A(A^t A)^{-1} A^t \mathbf{x}.$$

57

7.2 Orthogonal Matrices

Mnemonic:

$$P \approx \frac{A^t A}{A A^t}.$$

With an inner product in hand and a notion of orthogonality, we can define a notion of **orthogonal projection** of one vector onto another, and more generally of a vector onto a subspace spanned by multiple vectors.

7.3.1 Projection Onto a Vector

Say we have two vectors \mathbf{x} and \mathbf{y} , and we want to define "the component of \mathbf{x} that lies along \mathbf{y} ", which we'll call \mathbf{p} . We can work out what the formula should be using a simple model:

We notice that whatever p is, it will in the direction of \mathbf{y} , and thus $\mathbf{p} = \lambda \hat{\mathbf{y}}$ for some scalar λ , where in fact $\lambda = \|\mathbf{p}\|$ since $\|\hat{\mathbf{y}}\| = 1$. We will find that $\lambda = \langle \mathbf{x}, \hat{\mathbf{y}} \rangle$, and so

$$\mathbf{p} = \langle \mathbf{x}, \ \widehat{\mathbf{y}} \rangle \widehat{\mathbf{y}} = \frac{\langle \mathbf{x}, \ \mathbf{y} \rangle}{\langle \mathbf{y}, \ \mathbf{y} \rangle} \mathbf{y}.$$

Notice that we can then form a "residual" vector $\mathbf{r} = \mathbf{x} - \mathbf{p}$, which should satisfy $\mathbf{r}^{\perp}\mathbf{p}$. If we were to let λ vary as a function of a parameter t (making \mathbf{r} a function of t as well) we would find that this particular choice minimizes $\|\mathbf{r}(t)\|$.

7.3.2 Projection Onto a Subspace

In general, supposing one has a subspace $S = \text{span}\{\mathbf{y}_1, \mathbf{y}_2, \cdots, \mathbf{y}_n\}$ and (importantly!) the \mathbf{y}_i are orthogonal, then the projection of \mathbf{p} of x onto S is given by the sum of the projections onto each basis vector, yielding

$$\mathbf{p} = \sum_{i=1}^{n} \frac{\langle \mathbf{x}, \ \mathbf{y}_i \rangle}{\langle \mathbf{y}_i, \ \mathbf{y}_i \rangle} \mathbf{y}_i = \sum_{i=1}^{n} \langle \mathbf{x}, \ \mathbf{y}_i \rangle \widehat{\mathbf{y}}_i.$$

Note: this is part of why having an orthogonal basis is desirable!

Letting $A = [\mathbf{y}_1, \mathbf{y}_2, \cdots]$, then the following matrix projects vectors onto S, expressing them in terms of the basis \mathbf{y}_i^1 :

$$\tilde{P}_A = (AA^T)^{-1}A^T,$$

7.3 Projections 58

¹For a derivation of this formula, see the section on least-squares approximations.

while this matrix performs the projection and expresses it in terms of the standard basis:

$$P_A = A(AA^T)^{-1}A^T.$$

Equation of a plane: given a point \mathbf{p}_0 on a plane and a normal vector \mathbf{n} , any vector \mathbf{x} on the plane satisfies

$$\langle \mathbf{x} - \mathbf{p}_0, \ \mathbf{n} \rangle = 0$$

To find the distance between a point \mathbf{a} and a plane, we need only project \mathbf{a} onto the subspace spanned by the normal \mathbf{n} :

$$d = \langle \mathbf{a}, \mathbf{n} \rangle.$$

One important property of projections is that for any vector \mathbf{v} and for any subspace S, we have $\mathbf{v} - P_S(\mathbf{v}) \in S^{\perp}$. Moreover, if $\mathbf{v} \in S^{\perp}$, then $P_s(\mathbf{v})$ must be zero. This follows by noting that in equation ??, every inner product appearing in the sum vanishes, by definition of $\mathbf{v} \in S^{\perp}$, and so the projection is zero.

7.3.3 Least Squares

Proposition 7.3.2 (Normal Equations).

 \mathbf{x} is a least squares solution to $A\mathbf{x} = \mathbf{b}$ iff

$$A^t A \mathbf{x} = A^t \mathbf{b}$$

Derivation of normal equations.

The general setup here is that we would like to solve $A\mathbf{x} = \mathbf{b}$ for \mathbf{x} , where \mathbf{b} is not in fact in the range of A. We thus settle for a unique "best" solution $\tilde{\mathbf{x}}$ such that the error $||A\tilde{\mathbf{x}} - \mathbf{b}||$ is minimized.

Geometrically, the solution is given by projecting **b** onto the column space of A. To see why this is the case, define the residual vector $\mathbf{r} = A\tilde{\mathbf{x}} - \mathbf{b}$. We then seek to minimize $\|\mathbf{r}\|$, which happens exactly when \mathbf{r}^{\perp} im A. But this happens exactly when $\mathbf{r} \in (\text{im } A)^{\perp}$, which by the fundamental subspaces theorem, is equivalent to $\mathbf{r} \in \ker A^T$.

From this, we get the equation

$$A^{T}\mathbf{r} = \mathbf{0}$$

$$\implies A^{T}(A\tilde{\mathbf{x}} - \mathbf{b}) = \mathbf{0}$$

$$\implies A^{T}A\tilde{\mathbf{x}} = A^{T}\mathbf{b}.$$

7.3 Projections 59

where the last line is described as the **normal equations**.

If A is an $m \times n$ matrix and is of full rank, so it has n linearly independent columns, then one can show that $A^T A$ is nonsingular, and we thus arrive at the least-squares solution

$$\tilde{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b} \blacksquare$$

These equations can also be derived explicitly using Calculus applied to matrices, vectors, and inner products. This requires the use of the following formulas:

$$\frac{\partial}{\partial \mathbf{x}} \langle \mathbf{x}, \ \mathbf{a} \rangle = \mathbf{a}$$
$$\frac{\partial}{\partial \mathbf{x}} \langle \mathbf{x}, \ \mathbf{A} \mathbf{x} \rangle = (A + A^T) \mathbf{x}$$

as well as the adjoint formula

$$\langle A\mathbf{x}, \ \mathbf{x} \rangle = \langle \mathbf{x}, \ A^T\mathbf{x} \rangle..$$

From these, by letting A = I we can derive

$$\frac{\partial}{\partial \mathbf{x}} \|\mathbf{x}\|^2 = \frac{\partial}{\partial \mathbf{x}} \langle \mathbf{x}, \ \mathbf{x} \rangle = 2\mathbf{x}$$

The derivation proceeds by solving the equation

$$\frac{\partial}{\partial \mathbf{x}} \left\| \mathbf{b} - A\mathbf{x} \right\|^2 = \mathbf{0}..$$

7.4 Normal Forms

Remark 7.4.1: Every square matrix is similar to a matrix in Jordan canonical form.

7.5 Decompositions

7.5.1 The QR Decomposition

Gram-Schmidt is often computed to find an orthonormal basis for, say, the range of some matrix A. With a small modification to this algorithm, we can write A = QR where R is upper triangular and Q has orthogonal columns.

7.4 Normal Forms 60

Why is this useful? One reason is that this also allows for a particularly simple expression of least-squares solutions. If A = QR, then R will be invertible, and a bit of algebraic manipulation will show that

$$\tilde{\mathbf{x}} = R^{-1}Q^T\mathbf{b}..$$

How does it work? You simply perform Gram-Schmidt to obtain $\{\mathbf{u}_i\}$, then

$$Q = [\mathbf{u}_1, \mathbf{u}_2, \cdots].$$

The matrix R can then be written as

$$r_{ij} = \begin{cases} \langle \mathbf{u}_i, \ \mathbf{x}_j \rangle, & i \leq j, \\ 0, & \text{else.} \end{cases}$$

Explicitly, this yields the matrix


$$R = \begin{bmatrix} \langle \mathbf{u}_1, \ \mathbf{x}_1 \rangle & \langle \mathbf{u}_1, \ \mathbf{x}_2 \rangle & \langle \mathbf{u}_1, \ \mathbf{x}_3 \rangle & \cdots \\ 0 & \langle \mathbf{u}_2, \ \mathbf{x}_2 \rangle & \langle \mathbf{u}_2, \ \mathbf{x}_3 \rangle & \cdots \\ 0 & 0 & \langle \mathbf{u}_3, \ \mathbf{x}_3 \rangle & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Explain shortcut for diagonal

8 | Appendix: Lists of things to know

 ${\bf Textbook:\ Leon,\ } Linear\ Algebra\ with\ Applications$

8.1 Topics



- 3.5: Change of Basis
- 4.1: Linear Transformations
- 4.2: Matrix Representations
- 4.3: Similarity
 - Exam 1

- 5.1: Scalar Product in \mathbb{R}^n
- 5.2: Orthogonal Subspaces
- 5.3: Least Squares
- 5.4: Inner Product Spaces
- 5.5: Orthonormal Sets
- 5.6: Gram-Schmidt
- 6.1: Eigenvalues and Eigenvectors
 - Exam 2
- 6.2: Systems of Linear Differential Equations
- 6.3: Diagonalization
- 6.6: Quadratic Forms
- 6.7: Positive Definite Matrices
- 6.5: Singular Value Decomposition
- 7.7: The Moore-Penrose Pseudo-Inverse
 - Final Exam

8.2 Definitions

- System of equations
- Homogeneous system
- Consistent/inconsistent system
- Matrix
- Matrix (i.e. $A\mathbf{x} = \mathbf{b}$)
- Inverse matrix
- Singular matrix
- Determinant
- Trace
- Rank
- Elementary row operation
- Row equivalence
- Pivot
- Row Echelon Form
- Reduced Row Echelon Form
- Gaussian elimination
- Block matrix
- Vector space
- Vector subspace
- Linear transformation
- Span
- Linear independence
- Basis
- Change of basis
- Dimension
- Row space

8.2 Definitions 62

- Column space
- Image
- Null space
- Kernel
- Direct sum
- Projection
- Orthogonal subspaces
- Orthogonal complement
- Normal equations
- Least-squares solution
- Orthonormal
- Eigenvalue
- Eigenvector
- Characteristic polynomial
- Similarity
- Diagonalizable
- Inner product
- Bilinearity
- Multilinearity
- Defective
- Singular value decomposition
- QR factorization
- Gram-Schmidt process
- Spectral theorem
- Symmetric matrix
- Orthogonal matrix
- Positive-definite
- Quadratic form

8.3 Lower-division review



- Systems of linear equations
 - Consistent vs. Inconsistent
 - Possibilities for solutions
 - Geometric interpretation
- Matrix Inverses
 - Detecting if a matrix is singular
 - Computing the inverse
 - \diamondsuit Formula for 2x2 case
 - ♦ Augment with the identity
 - ♦ Cramer's Rule
- Vector Spaces

8.3 Lower-division review 63

- Definition in terms of closures
- Span
- Linear Independence
- Subspace and the subspace test
- Basis
- Common Computations
 - Reduction to RREF
 - Eigenvalues and eigenvectors
 - Basis for the column space
 - Basis for the nullspace
 - Basis for the eigenspace
 - Construct matrix from a given linear map
 - Construct change of basis matrix
 - Construct matrix projection onto subspace
 - Convert a basis to an orthonormal basis

8.4 Things to compute



- Construct a matrix representing a linear map
 - With respect to the standard basis in both domain and range
 - With respect to a nonstandard basis in the range
 - With respect to a nonstandard basis in the domain
 - With respect to nonstandard bases in both the domain and range
- Construct a change of basis matrix
- Check that a map is a linear transformation
- Compute the following spaces of a matrix and their orthogonal complements:
 - Row space
 - Column space
 - Null space
- Compute the shortest distance between a point and a plane
- Compute the least squares solution to linear system
- Prove that something is a vector space
- Prove that a map is an inner product
- Compute determinants
- Compute the RREF of a matrix
- Compute characteristic polynomials, eigenvalues, and eigenvectors
- Diagonalize a matrix
- Solve a system of ODEs resulting arising from tank mixing
- Compute the singular value decomposition of a matrix
- Compute the rank and nullity of a matrix
- Convert a set of vectors to a basis
- Convert a basis to an orthonormal basis

8.4 Things to compute 64

- Determine if a matrix is diagonalizable
- Compute the matrix for a projection onto a subspace
- Find the QR factorization of a matrix

Things to prove



- Prove facts about block matrices
- Prove facts about injective linear maps
- Prove facts about similar matrices
- Prove facts about orthogonal spaces and orthogonal complements
- Prove facts about inner products
- Prove facts about orthonormal sets
- Prove facts about eigenvalues/eigenvectors
- Understand when a matrix can be diagonalized
- Prove facts about diagonalizable matrices
- Prove facts about the orthogonal decomposition theorem

Ordinary Differential Equations

9.1 Techniques Overview

p(y)y' = q(x)separable y' + p(x)y = q(x)integrating factor y' = f(x, y), f(tx, ty) = f(x, y)y = xV(x) COV reduces to separable Bernoulli, divide by y^n and COV $u = y^{1-n}$ $y' + p(x)y = q(x)y^n$ $M_y = N_x : \varphi(x, y) = c(\varphi_x = M, \varphi_y = N)$ M(x,y)dx + N(x,y)dy = 0 $x^k e^{rx}$ for each root P(D)y = f(x, y)

Where e^{zx} yields $e^{ax} \cos bx$, $e^{ax} \sin bx$

8.5 Things to prove 65

9.2 Types of Equations



$$p(y)\frac{dy}{dx} - q(x) = 0 \implies \int p(y)dy = \int q(x)dx + C$$

$$\frac{dy}{dx} = f(x)g(y) \implies \int \frac{1}{g(y)} dy = \int f(x) dx + C$$

- Population growth:

$$\frac{dP}{dt} = kP \implies P = P_0 e^{kt}$$

- Logistic growth:

 \diamondsuit General form: $\frac{dP}{dt} = (B(t) - D(t))P(t)$

 \diamondsuit Assume birth rate is constant $B(t) = B_0$ and death rate is proportional to instantaneous population $D(t) = D_0 P(t)$. Then let $r = B_0, C = B_0/D_0$ be the carrying capacity:

$$\frac{dP}{dt} = r\left(1 - \frac{P}{C}\right)P \implies P(t) = \frac{P_0}{\frac{P_0}{C} + e^{-rt}(1 - \frac{P_0}{C})}$$

• First order linear:

$$\frac{dy}{dx} + p(x)y = q(x) \implies I(x) = e^{\int p(x)dx}, \qquad y(x) = \frac{1}{I(x)} \left(\int q(x)I(x)dx + C \right)$$

• Exact:

$$-M(x,y)dx + N(x,y)dy = 0 \text{ is exact } \iff \exists \varphi : \frac{\partial \varphi}{\partial x} = M(x,y), \ \frac{\partial \varphi}{\partial y} = N(x,y)$$
$$\iff \frac{\partial M}{\partial y} = \frac{\partial N}{x}$$

- General solution:

$$\varphi(x,y) = \int_{-\infty}^{\infty} M(s,y)ds + \int_{-\infty}^{y} N(x,t)dt - \int_{-\infty}^{y} \frac{\partial}{\partial t} \left(\int_{-\infty}^{\infty} M(s,t)ds \right) dt$$

(where $\int_{-\infty}^{x} f(t)dt$ means take the antiderivative of f and consider it a function of x)

• Cauchy Euler: #todo

• Bernoulli: todo

9.3 Linear Homogeneous

General form:

$$y^{(n)} + c_{n-1}y^{(n-1)} + \dots + c_2y'' + cy' + cy = 0$$
$$p(D)y = \prod (D - r_i)^{m_i} y = 0$$

where p is a polynomial in the differential operator D with roots r_i :

• Real roots: contribute m_i solutions of the form

$$e^{rx}, xe^{rx}, \cdots, x^{m_i-1}e^{rx}$$

• Complex conjugate roots: for r = a + bi, contribute $2m_i$ solutions of the form

$$e^{(a\pm bi)x}, xe^{(a\pm bi)x}, \dots, x^{m_i-1}e^{(a\pm bi)x}$$

= $e^{ax}\cos(bx), e^{ax}\sin(bx), xe^{ax}\cos(bx), xe^{ax}\sin(bx), \dots,$

Example: by cases, second order equation of the form

$$ay'' + by' + cy = 0$$

- Two distinct roots: $c_1e^{r_1x} + c_2e^{r_2x}$ - One real root: $c_1e^{rx} + c_2xe^{rx}$ - Complex conjugates $\alpha \pm i\beta$: $e^{\alpha x}(c_1\cos\beta x + c_2\sin\beta x)$

9.4 Linear Inhomogeneous

General form:

$$y^{(n)} + c_{n-1}y^{(n-1)} + \dots + c_2y'' + cy' + cy = F(x)$$
$$p(D)y = \prod (D - r_i)^{m_i} y = 0$$

Then solutions are of the form $y_c + y_p$, where y_c is the solution to the associated homogeneous system and y_p is a particular solution.

Methods of obtaining particular solutions

9.4.1 Undetermined Coefficients

- Find an operator p(D) the annihilates F(x) (so q(D)F = 0)
- Find solution of q(D)p(D) = 0, subtract of known solutions from homogeneous part to obtain the form of the trial solution $A_0 f(x)$, where A_0 is the undetermined coefficient
- Substitute trial solution into original equation to determine A_0

Useful Annihilators:

$$\begin{split} F(x) &= p(x): & D^{\deg(p)+1} \\ F(x) &= p(x)e^{ax}: & (D-a)^{\deg(p)+1} \\ F(x) &= \cos(ax) + \sin(ax): & D^2 + a^2 \\ F(x) &= e^{ax}(a_0\cos(bx) + b_0\sin(bx)): & (D-z)(D-\overline{z}) = D^2 - 2aD + a^2 + b^2 \\ F(x) &= p(x)e^{ax}\cos(bx) + p(x)e^{ax}\cos(bx): & ((D-z)(D-\overline{z}))^{\max(\deg(p),\deg(q))+1} \end{split}$$

9.4.2 Variation of Parameters

todo

9.4.3 Reduction of Order

todo

9.5 Systems of Differential Equations

General form:

$$\frac{\partial \mathbf{x}(t)}{\partial t} = A\mathbf{x}(t) + \mathbf{b}(t) \iff \mathbf{x}'(t) = A\mathbf{x}(t) + \mathbf{b}(t)$$

General solution to homogeneous equation:

$$c_1\mathbf{x_1}(t) + c_2\mathbf{x_2}(t) + \dots + c_n\mathbf{x_n}(t) = \mathbf{X}(t)\mathbf{c}$$

If A is a matrix of constants: $\mathbf{x}(t) = e^{\lambda_i t} \mathbf{v}_i$ is a solution for each eigenvalue/eigenvector pair $(\lambda_i, \mathbf{v}_i)$ - If A is defective, you'll need generalized eigenvectors.

Inhomogeneous Equation: particular solutions given by

$$\mathbf{x}_p(t) = \mathbf{X}(t) \int_0^t \mathbf{X}^{-1}(s) \mathbf{b}(s) \ ds$$

9.6 Laplace Transforms

Definitions:

$$H_{a}(t) := \begin{cases} 0, & 0 \le t < a \\ 1, & t \ge a \end{cases}$$

$$\delta(t) : \int_{\mathbb{R}} \delta(t - a) f(t) \ dt = f(a), \quad \int_{\mathbb{R}} \delta(t - a) \ dt = 1$$

$$(f * g)(t) = \int_{0}^{t} f(t - s) g(s) \ ds$$

$$L[f(t)] = L[f] = \int_{0}^{\infty} e^{-st} f(t) dt = F(s).$$

Useful property: for $a \leq b$, $H_a(t) - H_b(t) = \mathbb{1}[[a, b]]$.

$$t^{n}, n \in \mathbb{N} \iff \frac{n!}{s^{n+1}}, s > 0$$

$$t^{-\frac{1}{2}} \iff \sqrt{\pi}s^{-\frac{1}{2}} s > 0$$

$$e^{at} \iff \frac{1}{s-a}, s > a$$

$$\cos(bt) \iff \frac{s}{s^{2}+b^{2}}, s > 0$$

$$\sin(bt) \iff \frac{b}{s^{2}+b^{2}}, s > 0$$

$$\cosh(bt) \iff \frac{s}{s^{2}-b^{2}}, ?$$

$$\sinh(bt) \iff \frac{b}{s^{2}-b^{2}}, ?$$

$$\delta(t-a) \iff e^{-as}$$

$$H_{a}(t) \iff s^{-1}e^{-as}$$

$$H_{a}(t) \iff f(s-a)$$

$$f'(t) \iff sL(f) - f(0)$$

$$f''(t) \iff s^{n}L(f) - \sum_{i=0}^{n-1} s^{n-1-i}f^{(i)}(0)$$

$$f(t)g(t) \iff F(s) * G(s)$$

• For f periodic with period T, $L(f) = \frac{1}{1 + e^{-sT}} \int_0^T e^{-st} f(t) dt$

$$p(y)y'=q(x)$$
 separable
$$y'+p(x)y=q(x)$$
 integrating factor
$$y'=f(x,y), f(tx,ty)=f(x,y)$$

$$y=xV(x) \text{ COV reduces to separable}$$

$$y'+p(x)y=q(x)y^n \qquad \text{Bernoulli, divide by } y^n \text{ and COV } u=y^{1-n}$$

$$M(x,y)dx+N(x,y)dy=0 \qquad M_y=N_x: \varphi(x,y)=c(\varphi_x=M,\varphi_y=N)$$

$$P(D)y=f(x,y) \qquad x^ke^{rx} \text{ for each root}$$

9.6 Laplace Transforms

Theorem 9.6.1 (First Shifting Theorem).

$$L[e^{at}f(t)] = \int_0^\infty e^{(a-s)}f(t)dt = F(s-a),.$$

Remark 9.6.1: The general technique for solving differential equations with Laplace Transforms: - Take the Laplace Transform of all terms on both sides. - Solve for L[y] in terms of s. - Attempt an inverse Laplace Transformations - This may involve partial fraction decomposition, completing the square, and splitting numerators to match terms with known inverse transformations.

9.7 Systems of Differential Equations

Definition 9.7.1 (Wronksian).

For a collection of n functions $f_i: \mathbb{R}^n \to \mathbb{R}$, define the $n \times 1$ column vector

$$W(f_i)(\mathbf{p}) \coloneqq \begin{bmatrix} f_i(\mathbf{p}) \\ f'_i(\mathbf{p}) \\ f''_i(\mathbf{p}) \\ \vdots \\ f^{(n-1)}(\mathbf{p}) \end{bmatrix}.$$

The Wronskian of this collection is defined as

$$W(f_1, \cdots, f_n)(\mathbf{p}) := \det \begin{bmatrix} | & | & | \\ W(f_1)(\mathbf{p}) & W(f_2)(\mathbf{p}) & \cdots & W(f_n)(\mathbf{p}) \end{bmatrix}.$$

Proposition 9.7.1 (Wronskian detects linear dependence of functions).

A set of functions $\{f_i\}$ is linearly independent on $I \iff \exists x_0 \in I : W(x_0) \neq 0$.

 \triangle Warning 9.1: $W \equiv 0$ on I does not imply that $\{f_i\}$ is linearly dependent! Counterexample: $\{x, x + x^2, 2x - x^2\}$ where $W \equiv 0$ but $x + x^2 = 3(x) + (2x - x^2)$ is a linear combination of the other two functions.

Sufficient condition: each f_i is the solution to a linear homogeneous ODE L(y) = 0.

9.7.1 Linear Equations of Order n

The standard form of such equations is

$$y^{(n)} + a_1 y^{(n-1)} + a_2 y^{(n-2)} + \dots + a_n y'' + a_{n-1} y' + y = F(x).$$

All solutions will be the sum of the solution to the associated homogeneous equation and a single particular solution.

In the homogeneous case, examine the discriminant of the characteristic polynomial. Three cases arise:

- 1. $D > 0 \Rightarrow 2$ Real solutions, $c_1 e^{r_1 x} + c_2 e^{r_2 x}$
- 2. $D=0 \Rightarrow 1$ Real, 1 Complex, $(c_1+c_2x)e^{r_1x}$
- 3. $D < 0 \Rightarrow 2$ Complex, $e^{ax}(c_1 \cos bx + c_2 \sin bx)$

That is, every real root contributes a term of ce^{rx} , while a multiplicity of m multiplies the solution by a polynomial in x of degree m-1.

Every pair of complex roots contributes a term $ce^{r}(a\cos\omega x + b\sin\omega x)$, where r is the real part of the roots and ω is the complex part.

In the nonhomogeneous case, assume a solution in the most general form of F(x), and substitute it into the equation to solve for constant terms. For example,

1.
$$F(x) = P(x^n) \Rightarrow y_p = a + bx + cx^2 + \dots + (n+1)x^n$$

- 2. $F(x) = e^x \Rightarrow y_p = Ae^x$
- 3. $F(x) = A\cos(\omega x) \Rightarrow y_p = a\cos(\omega x) + b\sin(\omega x)$

9.8 Annihilators

Use to reduce a nonhomogeneous equation to a homogeneous one as a polynomial in the operator D

1.
$$(D-a) \Rightarrow e^{ax}$$

2.
$$(D-a)^{k+1} \Rightarrow x^k e^{ax}, x^{k-1} e^{ax}, \dots, e^{ax}$$

3.
$$D^{k+1} \Rightarrow x^k, x^{k-1}, \cdots, C$$

4.
$$D^2 - 2aD + a^2 + b^2 \Rightarrow e^{ax}\cos(bx), e^{ax}\sin(bx)$$

5.
$$(D^2 - 2aD + a^2 + b^2)^{k+1} \Rightarrow x^k e^{ax} \cos(bx), x^{k-1} e^{ax} \cos(bx), x^k e^{ax} \sin(bx), x^{k-1} e^{ax} \sin(bx), \cdots$$

9.8 Annihilators 71

10 Algebra

9.9 Complex Solutions

 \sim

F(x) of the form $e^{ax}sin(kx)$ can be rewritten as $e^{(a+ki)x}$

10 | Algebra

This section is very sketchy!

10.1 To Sort




- Burnside's Lemma
- Cauchy's Theorem
 - If $|G| = n = \prod p_i^{k_i}$, then for each i there exists a subgroup H of order p_i .
- The Sylow Theorems
 - If $|G| = n = \prod p_i^{k_i}$, for each ii and each $1 \le k_j \le k_i$ then there exists a subgroup H of order $p_i^{k_j}$.
- Galois Theory
- More terms: http://mathroughguides.wikidot.com/glossary:abstract-algebra
- Order p: One, Z_p
- Order p^2 : Two abelian groups, Z_{p^2}, Z_p^2
- Order p^3 :
 - 3 abelian $Z_{p^3}, Z_p \times Z_{p^2}.Z_p^3$,
 - -2 others $Z_p \rtimes Z_{p^2}$.
 - \Diamond The other is the quaternion group for p=2 and a group of exponent p for p>2.
- Order pq:

$$-p \mid q-1$$
: Two groups, Z_{pq} and $Z_q \rtimes Z_p$
- Else cyclic, Z_{pq}

- Every element in a permutation group is a product of disjoint cycles, and the order is the lcm of the order of the cycles.
- The product ideal IJ is not just elements of the form ij, it is all sums of elements of this form! The product alone isn't enough.
- The intersection of any number of ideals is also an ideal

10.2 Big List of Notation



10.3 Group Theory

Notation: H < G a subgroup, N < G a normal subgroup, concatenation is a generic group operation.

- \mathbb{Z}_n the unique cyclic group of order n
- **Q** the quaternion group
- $G^n = G \times G \times \cdots G$
- Z(G) the center of G
- o(G) the order of a group

10.2 Big List of Notation 73

- S_n the symmetric group
- A_n the alternating group
- D_n the dihedral group of order 2n
- Group Axioms
 - Closure: $a, b \in G \implies ab \in G$
 - Identity: $\exists e \in G \mid a \in G \implies ae = ea = a$
 - Associativity: $a, b, c \in G \implies (ab)c = a(bc)$
 - Inverses: $a \in G \implies \exists b \in G \mid ab = ba = e$
- Definitions:
 - Order
 - \Diamond Of a group: o(G) = |G|, the cardinality of G
 - \Diamond Of an element: $o(g) = \min \{ n \in \mathbb{N} : g^n = e \}$
 - Index
 - Center: the elements that commute with everything
 - Centralizer: all elements that commute with a given element/subgroup.
 - Group Action: a function $f: X \times G \to G$ satisfying

$$\diamondsuit \ x \in X, g_1, g_2 \in G \implies g_1.(g_2.x) = (g_1g_2).x$$

$$\Diamond x \in X \implies e.x = x$$

- Orbits partition any set
- Transitive Action
- Conjugacy Class: $C \subset G$ is a conjugacy class \iff

$$\diamondsuit \ x \in C, g \in G \implies gxg^{-1} \in C$$

$$\Diamond x, y \in C \implies \exists g \in G : gxg^{-1} = y$$

- \Diamond i.e. subsets that are closed under G acting on itself by conjugation and on which the action is transitive
- ♦ i.e. orbits under the conjugation action
- \Diamond The order of any conjugacy class divides the order of G
- p-group: Any group of order p^n .
- Simple Group: no nontrivial normal subgroups
- Normal Series: $0 ext{ ≤ } H_0 ext{ ≤ } H_1 \cdots ext{ ≤ } G$
- Composition Series: The successive quotients of the normal series
- Solvable: G is solvable \iff G has an abelian composition series.
- One step subgroup test:

$$a, b \in H \implies ab^{-1} \in H$$

• Useful isomorphism invariants:

10.3 Group Theory 74

- Order profile of elements: n_1 elements of order p_1 , n_2 elements of order p_2 , etc \diamondsuit Useful to look at elements of order 2!
- Order profile of subgroups
- $-Z(A) \stackrel{\circ}{\cong} Z(B)$
- Number of generators (generators are sent to generators)
- Number and size of conjugacy classes
- Number of Sylow-p subgroups.
- Commutativity
- "Being cyclic"
- Automorphism Groups
- Solvability
- Nilpotency
- Useful homomorphism invariants

$$-\varphi(e) = e$$

$$-|g| = m < \infty \implies |\varphi(g)| = m$$


$$- \text{ Inverses, i.e. } \varphi(a)^{-1} = \varphi(a^{-1})$$

$$- H < G \implies \varphi(H) < G'$$

$$\diamondsuit H' < G' \implies \varphi^{-1}(H') < G$$

$$-|G| < \infty \implies \varphi(G) \text{ divides } |G|, |G'|$$

10.4 Big Theorems



$$G \cong \mathbb{Z}_{p_1^{k_1}} \oplus \mathbb{Z}_{p_2^{k_2}} \oplus \cdots \oplus \mathbb{Z}_{p_n^{k_n}},$$

where (p_i, k_i) are the set of elementary divisors of G.

• Isomorphism Theorems

$$\varphi: G \to G' \implies \frac{G}{\ker \varphi} \cong \varphi(G)$$

$$H \trianglelefteq G, \ K < G \implies \frac{K}{H \cap K} \cong \frac{HK}{H}$$

$$H, K \trianglelefteq G, \ K < H \implies \frac{G/K}{H/K} \cong \frac{G}{H}$$

- Lagrange's Theorem: $H < G \implies o(H) \mid o(G)$
 - Converse is false: $o(A_4) = 12$ but has no order 6 subgroup.

- The GZ Theorem: G/Z(G) cyclic implies that $G \in \mathbf{Ab}$.
- Orbit Stabilizer Theorem: $G/x_0 \cong Gx$
- The Class Equation
 - Let $G \curvearrowright X$ and $\mathcal{O}_i \subseteq X$ be the nontrivial orbits, then

$$|X| = |X_0| + \sum_{[x_i] \in X/G} |Gx|.$$

- The right hand side is the number of fixed points, plus a sum over all of the orbits of size greater than 1, where any representative within the orbit is chosen and we look at the index of its stabilizer in G.
- Let $G \curvearrowright G$ and for each nontrivial conjugacy class C_G choose a representative $[x_i]$ $C_G = C_G(x_i)$ to obtain

$$|G| = |Z(G)| + \sum_{[x_i] = C_G(x_i)} [G : [x_i]].$$

- Useful facts:
 - $-H < G \in \mathbf{Ab} \implies H \triangleleft G$
 - \Diamond Converse doesn't hold, even if all subgroups are normal. Counterexample: $\mathbf Q$
 - $-G/Z(G) \cong \operatorname{Inn}(G)$
 - $-H, K < G \text{ with } H \cong K \not \Longrightarrow G/H \cong G/K$
 - \diamondsuit Counterexample: $G = \mathbb{Z}_4 \times \mathbb{Z}_2, H = <(0,1)>, K = <(2,0)>$. Then $G/H \cong \mathbb{Z}_4 \not\cong \mathbb{Z}_2^2 \cong G/K$
 - $-G \in \mathbf{Ab} \implies$ for each p dividing o(G), there is an element of order p
 - Any surjective homomorphism $\varphi: A \to B$ where o(A) = o(B) is an isomorphism
 - If G is abelian, for each $d \mid |G|$ there is exactly one subgroup of order d.
- Sylow Subgroups:
 - Todo
- Big List of Interesting Groups
 - $-\mathbb{Z}_4,\mathbb{Z}_2^2$

 - $-Q=\langle a,b|a^4=1,a^2=b^2,ab=ba^3\rangle$ the quaternion group $-S^3$, the smallest nonabelian group

10.4 Big Theorems 76

• Chinese Remainder Theorem:

$$\mathbb{Z}_{pq} \cong \mathbb{Z}_p \oplus \mathbb{Z}_q \iff (p,q) = 1$$

- Fundamental Theorem of Finitely Generated Abelian Groups:
- $-G = \mathbb{Z}^n \oplus \bigoplus \mathbb{Z}_{q_i}$
- Finding all of the unique groups of a given order: #todo

10.4.1 Cyclic Groups

- Generated by ?
- For each d dividing o(G), there exists a subgroup H of order d.
 - If $G = \langle a \rangle$, then take $H = \langle a^{\frac{n}{d}} \rangle$

10.4.2 The Symmetric Group

- Generated by:
 - Transpositions
 - #todo
- Cycle types: characterized by the number of elements in the cycle.
 - Two elements are in the same conjugacy class \iff they have the same cycle type.
- Inversions: given $\tau = (p_1 \cdots p_n)$, a pair p_i, p_j is inverted iff i < j but $p_j < p_i$
- Can count inversions $N(\tau)$
 - Equal to minimum number of transpositions to obtain non-decreasing permutation
- Sign of a permutation: $\sigma(\tau) = (-1)^{N(\tau)}$
- Parity of permutations $\cong (\mathbb{Z}, +)$
 - even \circ even = even
 - $\text{ odd } \circ \text{ odd } = \text{even}$
 - even \circ odd = odd

10.5 Ring Theory



Ring Axioms

- Examples:
- Non-Examples:

10.5 Ring Theory 77

- Definition of an Ideal
- Definitions of types of rings:
 - Field
 - Unique Factorization Domain (UFD)
 - Principal Ideal Domain (PID)
 - Euclidean Domain:
 - Integral Domain
 - Division Ring

field
$$\implies$$
 Euclidean Domain \implies PID \implies UFD \implies integral domain.

- Counterexamples to inclusions are strict:
 - An ED that is not a field:
 - A PID that is not an ED: $\mathbb{Q}[\sqrt{19}]$
 - A UFD that is not a PID:
 - An integral domain that is not a UFD:
- Integral Domains
- Unique Factorization Domains
- Prime Elements
- Prime Ideals
- Field Extensions
- The Chinese Remainder Theorem for Rings
- Polynomial Rings
 - Irreducible Polynomials
 - \Diamond Over \mathbb{Z}_2 :


$$x$$
, $x + 1$, $x^2 + x + 1$, $x^3 + x + 1$, $x^3 + x^2 + 1$.

- ♦ Eisenstein's Criterion
- Gauss' Lemma

When is $\mathbb{Q}(\sqrt{d})$ a field?

$oldsymbol{1}oldsymbol{1}$ Number Theory

11.1 Notation and Basic Definitions



 \mathbb{Z}_n

 \mathbb{Z}_n^{\times}

the greatest common divisor the ring of integers $\mod n$ the group of units $\mod n$.

78

Number Theory

Definition 11.1.1 (Multiplicative Functions).

A function $f: \mathbb{Z} \to \mathbb{Z}$ is said to be **multiplicative** iff

$$(a,b) = 1 \implies f(ab) = f(a)f(b).$$

11.2 Big Theorems

 \sim

Link to theorems

11.3 Primes

<u>_</u>

Theorem 11.3.1 (The fundamental theorem of arithmetic).

Every $n \in \mathbb{Z}$ can be written uniquely as

$$n = \prod_{i=1}^{m} p_i^{k_i}$$

where the p_i are the m distinct prime divisors of n.

Remark 11.3.1: Note that the number of distinct prime factors is m, while the total number of factors is $\prod_{i=1}^{m} (k_i + 1)$.

11.4 Divisibility



Definition 11.4.1 (Divisibility).

$$a \mid b \iff b \equiv 0 \mod a \iff \exists k \text{ such that } ak = b$$

11.4.1 gcd, lcm

Remark 11.4.1: gcd(a, b) can be computed by taking prime factorizations of a and b, intersecting the primes occurring, and taking the lowest exponent that appears. Dually, lcm(a, b) can be computed by taking the *union* and the *highest* exponent.

11.2 Big Theorems 79

Check

Proposition 11.4.1 (Relationship between gcd and lcm).

$$xy = \gcd(x, y) \operatorname{lcm}(x, y)$$

Proposition 11.4.2(?).

If $d \mid x$ and $d \mid y$, then

$$\gcd(x,y) = d \cdot \gcd\left(\frac{x}{d}, \frac{y}{d}\right)$$
$$\operatorname{lcm}(x,y) = d \cdot \operatorname{lcm}\left(\frac{x}{d}, \frac{y}{d}\right)$$

Check

Proposition 11.4.3 (Useful properties of gcd).

$$\gcd(x, y, z) = \gcd(\gcd(x, y), z)$$

$$gcd(x, y) = gcd(x \mod y, y)$$

$$\gcd(x,y) = \gcd(x-y,y).$$

11.4.2 The Euclidean Algorithm

gcd(a, b) can be computed via the Euclidean algorithm, taking the final bottom-right coefficient.

Example of Euclidean algorithm,

11.5 Modular Arithmetic



Generally concerned with the multiplicative group (\mathbb{Z}_n, \times) .

Theorem 11.5.1 (The Chinese Remainder Theorem).

The system

$$x \equiv a_1 \mod m_1$$

$$x \equiv a_2 \mod m_2$$

$$\vdots$$

$$x \equiv a_r \mod m_r$$

has a unique solution $x \mod \prod m_i \iff (m_i, m_j) = 1$ for each pair i, j, given by

$$x = \sum_{j=1}^{r} a_j \frac{\prod_i m_i}{m_j} \left[\frac{\prod_i m_i}{m_j} \right]^{-1} \mod m_j.$$

Theorem 11.5.2 (Euler's Theorem).

$$a^{\varphi(p)} \equiv 1 \mod n$$
.

Theorem 11.5.3 (Fermat's Little Theorem).

$$x^p \equiv x \mod p$$

$$x^{p-1} \equiv 1 \mod p \quad \text{if } p \nmid a$$

11.5.1 Diophantine Equations

Proposition 11.5.1 (Solutions to linear Diophantine equations).

Consider ax + by = c. This has solutions iff $c = 0 \mod(a, b) \iff \gcd(a, b)$ divides c.

How to obtain solutions

11.5.2 Computations

Proposition 11.5.2(?).

If $x \equiv 0 \mod n$, then $x \equiv 0 \mod p^k$ for all p^k appearing in the prime factorization of n.

Remark 11.5.1: If there are factors of the modulus in the equation, peel them off with addition, using the fact that $nk \equiv 0 \mod n$.

$$x \equiv nk + r \mod n$$
$$\equiv r \mod n$$

11.5 Modular Arithmetic 81

So take x = 463, n = 4, then use $463 = 4 \cdot 115 + 4$ to write

$$463 \equiv y \mod 4$$

$$\implies 4 \cdot 115 + 3 \equiv y \mod 4$$

$$\implies 3 \equiv y \mod 4.$$

Proposition 11.5.3(Repeated square/fast exponentiation). For any n,

$$x^k \mod n \equiv (x^{k/d} \mod n)^d \mod n.$$

Example 11.5.1 (?):

$$2^{25} \equiv (2^5 \mod 5)^5 \mod 5$$
$$\equiv 2^5 \mod 5$$
$$\equiv 2 \mod 5$$

Remark 11.5.2: Make things easier with negatives! For example, mod 5,

$$4^{25} \equiv (-1)^{25} \mod 5$$
$$\equiv (-1) \mod 5$$
$$\equiv 4 \mod 5$$

11.5.3 Invertibility

Proposition 11.5.4 (Reduction of modulus).

$$xa = xb \mod n \implies a = b \mod \frac{n}{(x,n)}.$$

Proposition 11.5.5 (Characterization of invertibility).

$$x \in \mathbb{Z}_n^{\times} \iff (x, n) = 1,$$

and thus

$$\mathbb{Z}_n^{\times} = \{1 \le x \le n : (x, n) = 1\}$$

and
$$|\mathbb{Z}_n^{\times}| = \varphi(n)$$
.

Example 11.5.2 (Using invertibility): One can reduce equations by dividing through by a unit. Pick any x such that $x \mid a$ and $x \mid b$ with (x, n) = 1, then

$$a = b \mod n \implies \frac{a}{x} = \frac{b}{x} \mod n.$$

11.6 The Totient Function

Definition 11.6.1 (Euler's Totient Function).

$$\varphi(n) = |\{1 \le x \le n : (x, n) = 1\}|$$

Example 11.6.1 (?):

$$\varphi(1) = |\{1\}| = 1$$

$$\varphi(2) = |\{1\}| = 1$$

$$\varphi(3) = |\{1, 2\}| = 2$$

$$\varphi(4) = |\{1, 3\}| = 2$$

$$\varphi(5) = |\{1, 2, 3, 4\}| = 4$$

Proposition 11.6.1 (Formulas involving the totient).

$$\varphi(p) = p - 1$$

$$\varphi(p^k) = p^{k-1}(p - 1)$$

$$\varphi(n) = n \prod_{i=1}^{?} \left(1 - \frac{1}{p_i}\right)$$

$$n = \sum_{d \mid n} \varphi(d)$$

Proof (?).

All numbers less than p are coprime to p; there are p^k numbers less than p^k and the only numbers not coprime to p^k are multiples of p, i.e. $\left\{p, p^2, \cdots p^{k-1}\right\}$ of which there are k-1, yielding p^k-p^{k-1}

Along with the fact that φ is multiplicative, so $(p,q)=1 \implies \varphi(pq)=\varphi(p)\varphi(q)$, compute this for any n by taking the prime factorization.

With these properties, one can compute:

$$\varphi(n) = \varphi\left(\prod_{i} p_{i}^{k_{i}}\right)$$

$$= \prod_{i} p_{i}^{k_{i}-1}(p_{i}-1)$$

$$= n\left(\frac{\prod_{i} (p_{i}-1)}{\prod_{i} p_{i}}\right)$$

$$= n\prod_{i} \left(1 - \frac{1}{p_{i}}\right)$$

\todo[inline]{Check and explain}

11.7 Quadratic Residues

 $\begin{tabular}{ll} \textbf{Definition 11.7.1} & (Quadratic Residue). \\ \end{tabular}$

x is a quadratic residue mod n iff there exists an a such that $a^2 = x \mod n$.

Proposition 11.7.1(?).

In \mathbb{Z}_p , exactly half of the elements (even powers of generator) are quadratic residues.

Proposition 11.7.2(?).

-1 is a quadratic residue in $\mathbb{Z}_p \iff p = 1 \mod 4$.

Definition 11.7.2 (The Jacobi Symbols).

todo

Definition 11.7.3 (The Legendre Symbol).

todo

11.7 Quadratic Residues

12 Sequences

11.8 Primality Tests



Proposition 11.8.1 (Fermat Primality Test).

If n is prime, then

$$a^{n-1} = 1 \mod n$$

Proposition 11.8.2 (Miller-Rabin Primality Test).

n is prime iff

$$x^2 = 1 \mod n \implies x = \pm 1$$

11.9 Sequences in Metric Spaces



Theorem 11.9.1 (Bolzano-Weierstrass).

Every bounded sequence has a convergent subsequence.

Theorem 11.9.2 (Heine-Borel).

In \mathbb{R}^n , X is compact \iff X is closed and bounded.

Remark 11.9.1: Necessity of \mathbb{R}^n : $X = (\mathbb{Z}, d(x, y) = 1)$ is closed, complete, bounded, but not compact since $\{1, 2, \dots\}$ has no convergent subsequence

Proposition 11.9.1 (Converse of Heine-Borel).


Converse holds iff bounded is replaced with totally bounded

12 | Sequences

Notation: $\{a_n\}_{n\in\mathbb{N}}$ is a **sequence**, $\sum_{i\in\mathbb{N}}a_i$ is a **series**.

11.8 Primality Tests 85

12.1 Known Examples



$$c, c^2, c^3, \dots = \{c^n\}_{n=1}^{\infty} \to 0$$

$$\forall |c| < 1$$

$$\frac{1}{c},\frac{1}{c^2},\frac{1}{c^3},\ldots = \left\{\frac{1}{c^n}\right\}_{n=1}^{\infty} \to 0$$

$$\forall |c| > 1$$

$$1, \frac{1}{2^c}, \frac{1}{3^c}, \dots = \left\{\frac{1}{n^c}\right\}_{n=1}^{\infty} \to 0$$

$$\forall c > 0$$

12.2 Convergence

Definition 12.2.1 (Convergence of a Sequence).

A sequence $\{x_i\}$ converges to L iff

$$\forall \varepsilon > 0, \exists N > 0 \text{ such that } n \geq N \implies |x_n - L| < \varepsilon.$$

Theorem 12.2.1 (Squeeze Theorem).

$$b_n \leq a_n \leq c_n$$
 and $b_n, c_n \to L \implies a_n \to L$

Theorem 12.2.2 (Monotone Convergence Theorem for Sequences).

If $\{a_j\}$ monotone and bounded, then $a_j \to L = \limsup a_i < \infty$.

Theorem 12.2.3 (Cauchy Criteria).

 $|a_m - a_n| \to 0 \in \mathbb{R} \implies \{a_i\} \text{ converges.}$

12.2.1 Checklist

- Is the sequence bounded?
 - $-\{a_i\}$ not bounded \implies not convergent
 - If bounded, is it monotone?
 - $\Diamond \{a_i\}$ bounded and monotone \implies convergent

- Use algebraic properties of limits
- Epsilon-delta definition
- Algebraic properties and manipulation:
 - Limits commute with \pm, \times , Div and $\lim C = C$ for constants.
 - \diamondsuit E.g. Divide all terms by *n* before taking limit
 - \diamondsuit Clear denominators

13 | Sums ("Series")

Definition 13.0.1 (Series).

A series is an function of the form

$$f(x) = \sum_{j=1}^{\infty} c_j x^j.$$



13.1 Known Examples



13.1.1 Conditionally Convergent

$$\sum_{k=1}^{\infty} k^{p} < \infty \qquad \iff p \le 1$$

$$\sum_{k=1}^{\infty} \frac{1}{k^{p}} < \infty \qquad \iff p > 1$$

$$\sum_{k=1}^{\infty} \frac{1}{k} = \infty$$

13.1.2 Convergent

$$\sum_{n=1}^{\infty} \frac{1}{n^2} < \infty$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} < \infty$$

$$\sum_{n=1}^{\infty} \frac{1}{n^{\frac{3}{2}}} < \infty$$

$$\sum_{n=1}^{\infty} \frac{1}{n!} = e$$

$$\sum_{n=1}^{\infty} \frac{1}{c^n} = \frac{c}{c-1}$$

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{c^n} = \frac{c}{c+1}$$

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n} = \ln 2$$

13.1.3 Divergent

$$\sum_{n=1}^{\infty} \frac{1}{n} = \infty$$

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \infty$$

13.2 Convergence

Useful reference: http://math.hawaii.edu/~ralph/Classes/242/SeriesConvTests.pdf

Definition 13.2.1 (Absolutely Convergent).

todo

Remark 13.2.1: $a_n \to 0$ does not imply $\sum a_n < \infty$. Counterexample: the harmonic series.

Proposition 13.2.1(?).

Absolute convergence \implies convergence

Proposition 13.2.2 (The Cauchy Criterion).

$$\limsup a_i \to 0 \implies \sum a_i \text{ converges}$$

13.2.1 The Big Tests

Theorem 13.2.1 (Comparison Test).

- $a_n < b_n \sum b_n < \infty \implies \sum a_n < \infty$ $b_n < a_n \sum b_n = \infty \implies \sum a_n = \infty$

Theorem 13.2.2 (Ratio Test).

$$R = \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right|$$

- R < 1: absolutely convergent
- R > 1: divergent
- R = 1: inconclusive

Theorem 13.2.3 (Root Test).

$$R = \limsup_{n \to \infty} \sqrt[n]{|a_n|}$$

- R < 1: convergent - R > 1: divergent - R = 1: inconclusive

Theorem 13.2.4 (Integral Test).

$$f(n) = a_n \implies \sum a_n < \infty \iff \int_1^\infty f(x)dx < \infty$$

Theorem $13.2.5(Limit\ Test)$.

$$\lim_{n \to \infty} \frac{a_n}{b_n} = L < \infty \implies \sum a_n < \infty \iff \sum b_n < \infty$$

Theorem 13.2.6 (Alternating Series Test).

$$a_n \downarrow 0 \implies \sum (-1)^n a_n < \infty$$

Theorem 13.2.7 (Weierstrass M-Test).

$$\sum_{n=1}^{\infty} \|f_n\|_{\infty} < \infty \implies \exists f \text{ such that } \left\|\sum_{n=1}^{\infty} f_n - f\right\|_{\infty} \to 0$$

In other words, the series converges uniformly.

Slogan: Convergence of the sup norms implies uniform convergence'

Remark 13.2.2: The M in the name comes from defining $\sup \{f_k(x)\} := M_n$ and requiring $\sum |M_n| < M_n$

13.2.2 Checklist

- Do the terms tend to zero?
 - $-a_i \not\to 0 \implies \sum a_i = \infty.$
 - ♦ Can check with L'Hopital's rule
- There are exactly 6 tests at our disposal:
 - Comparison, root, ratio, integral, limit, alternating
- Is the series alternating?
 - If so, does $a_n \downarrow 0$?
 - \Diamond If so, **convergent**
- Is this series bounded above by a known convergent series?
 - p series with p > 1, i.e. : $\sum a_n \le \sum \frac{1}{n^p} < \infty$ Geometric series with |x| < 1, i.e. $\sum a_n \le \sum x^n$
- Is this series bounded below by a known divergent series?
 - p series with $p \leq 1$, i.e. $\infty = \sum_{n} \frac{1}{n^p} \leq \sum_{n} a_i$
- Are the ratios strictly less than or greater than 1?
 - $< 1 \implies$ convergent
 - $->1 \implies$ convergent
- Does the integral analogue converge?

- Integral converges ⇐⇒ sum converges
- Try the root test
 - $< 1 \implies$ convergent
 - $->1 \implies$ convergent
- Try the limit test
 - Attempt to divide each term to obtain a known convergent/divergent series

Some Pattern Recognition:

- (stuff)!: Ratio test (only test that will work with factorials!!)
- $(stuff)^n$: Root test or ratio test
- Replace a_n with an f(x) that's easy to integrate integral test
- p(x) or $\sqrt{p(x)}$: comparison or limit test

13.3 Radius of Convergence



Proposition 13.3.1 (Finding the radius of convergence).

Use the fact that

$$\lim_{k \to \infty} \left| \frac{a_{k+1} x^{k+1}}{a_k x^k} \right| = |x| \lim_{k \to \infty} \left| \frac{a_{k+1}}{a_k} \right| < 1 \implies \sum a_k x^k < \infty,$$


so take $L\coloneqq\lim_{k\to\infty}\frac{a_{k+1}}{a_k}$ and then obtain the radius as

$$R = \frac{1}{L} = \lim_{k \to \infty} \frac{a_k}{a_{k+1}}$$

Remark 13.3.1:

- Note $L=0 \implies$ absolutely convergent everywhere
- $L = \infty \implies$ convergent only at x = 0.
- Also need to check endpoints R, -R manually.

14 Real Analysis





Definition 14.1.1 (Continuously Differentiable).

A function is **continuously differentiable** iff f is differentiable and f' is continuous. Conventions:

 $\bullet \ \ Integrable \ {\it means} \ Riemann \ integrable.$

f	a functional $\mathbb{R}^n \to \mathbb{R}$
${f f}$	a function $\mathbb{R}^n \to \mathbb{R}^m$
A, E, U, V	open sets
A'	the limit points of A
\overline{A}	the closure of A
$A^{\circ} \coloneqq A \setminus A'$	the interior of A
K	a compact set
\mathcal{R}_A	the space of Riemann integral functions on A
$C^{j}(A)$	the space of j times continuously differentiable functions $f:\mathbb{R}^n\to\mathbb{R}$
$\{f_n\}$	a sequence of functions
$\{x_n\}$	a sequence of real numbers
$f_n \to f$	pointwise convergence
$f_n ightrightarrows f$	uniform convergence
$x_n \nearrow x$	$x_i \leq x_j$ and x_j converges to x
$x_n \searrow x$	$x_i \ge x_j$ and x_j converges to x
$\sum_{k\in\mathbb{N}}f_k$	a series
D(f)	the set of discontinuities of f .

14.2 Big Ideas



Summary for GRE:

- Limits,
- Continuity,
- Boundedness,
- Compactness,

14.1 Notation 92

- Definitions of topological spaces,
- Lipschitz continuity
- Sequences and series of functions.
- Know the interactions between the following major operations:
 - Continuity (pointwise limits)
 - Differentiability
 - Integrability
 - Limits of sequences
 - Limits of series/sums
- The derivative of a continuous function need not be continuous
- A continuous function need not be differentiable
- A uniform limit of differentiable functions need not be differentiable
- A limit of integrable functions need not be integrable
- An integrable function need not be continuous
- An integrable function need not be differentiable

Theorem 14.2.1 (Generalized Mean Value Theore).

$$f, g$$
 differentiable on $[a, b] \implies \exists c \in [a, b] : [f(b) - f(a)] g'(c) = [g(b) - g(a)] f'(c)$

Corollary 14.2.1 (Mean Value Theorem).

todo

14.3 Important Examples

14.4 Limits

todo

14.5 Commuting Limits



- Suppose $f_n \to f$ (pointwise, not necessarily uniformly)
- Let $F(x) = \int f(t)$ be an antiderivative of f
- Let $f'(x) = \frac{\partial f}{\partial x}(x)$ be the derivative of f.

Then consider the following possible ways to commute various limiting operations:

Does taking the derivative of the integral of a function always return the original function?

$$\left[\frac{\partial}{\partial x}, \int dx\right]: \qquad \qquad \frac{\partial}{\partial x} \int f(x,t) dt =_{?} \int \frac{\partial}{\partial x} f(x,t) dt$$

Answer: Sort of (but possibly not).

Counterexample:

$$f(x) = \begin{cases} 1 & x > 0 \\ -1 & x \le 0 \end{cases} \implies \int f \approx |x|,$$

which is not differentiable. (This is remedied by the so-called "weak derivative")

Sufficient Condition: If f is continuous, then both are always equal to f(x) by the FTC.

Is the derivative of a continuous function always continuous?

$$\left[\frac{\partial}{\partial x}, \lim_{x_i \to x}\right]: \qquad \lim_{x_i \to x} f'(x_n) =_{?} f'(\lim_{x_i \to x} x)$$

Answer: No.

Counterexample:

$$f(x) = \begin{cases} x^2 \sin(1/x) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases} \implies f'(x) = \begin{cases} 2x \sin\left(\frac{1}{x}\right) - \cos\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

which is discontinuous at zero.

Sufficient Condition: There doesn't seem to be a general one (which is perhaps why we study C^k functions).

14.5 Commuting Limits 94

Is the limit of a sequence of differentiable functions differentiable and the derivative of the limit?

$$\left[\frac{\partial}{\partial x}, \lim_{f_n \to f}\right]:$$

$$\lim_{f_n \to f} \frac{\partial}{\partial x} f_n(x) =_{?} \frac{\partial}{\partial x} \lim_{f_n \to f} f_n(x)$$

Answer: Super no – even the uniform limit of differentiable functions need not be differentiable!

Counterexample:
$$f_n(x) = \frac{\sin(nx)}{\sqrt{n}} \Rightarrow f = 0 \text{ but } f'_n \not\to f' = 0$$

Sufficient Condition: $f_n \Rightarrow f$ and $f_n \in C^1$.

Is the limit of a sequence of integrable functions integrable and the integral of the limit?

$$\left[\int dx, \lim_{f_n \to f}\right](f): \qquad \lim_{f_n \to f} \int f_n(x) dx = \int \lim_{f_n \to f} f_n(x) dx$$

Answer: No.

Counterexample: Order $\mathbb{Q} \cap [0,1]$ as $\{q_i\}_{i \in \mathbb{N}}$, then take

$$f_n(x) = \sum_{i=1}^n \mathbb{1}[q_n] \to \mathbb{1}[\mathbb{Q} \cap [0,1]]$$

where each f_n integrates to zero (only finitely many discontinuities) but f is not Riemann-integrable.

Sufficient Condition: - $f_n \Rightarrow f$, or - f integrable and $\exists M : \forall n, |f_n| < M$ (f_n uniformly bounded)

Is the integral of a continuous function also continuous?

$$\left[\int dx, \lim_{x_i \to x}\right] : \qquad \lim_{x_i \to x} F(x_i) = \operatorname{P}\left(\lim_{x_i \to x} x_i\right)$$

14.5 Commuting Limits 95

Answer: Yes.

Proof: |f(x)| < M on I, so given c pick a sequence $x \to c$. Then

$$|f(x)| < M \implies \left| \int_{c}^{x} f(t)dt \right| < \int_{c}^{x} Mdt \implies |F(x) - F(c)| < M(b-a) \to 0$$

Is the limit of a sequence of continuous functions also continuous?

$$\left[\lim_{x_i \to x}, \lim_{f_n \to f}\right] : \qquad \qquad \lim_{f_n \to f} \lim_{x_i \to x} f(x_i) =_{?} \lim_{x_i \to x} \lim_{f_n \to f} f_n(x_i)$$

Answer: No.

Counterexample: $f_n(x) = x^n \to \delta(1)$

Sufficient Condition: $f_n \rightrightarrows f$

Does a sum of differentiable functions necessarily converge to a differentiable function?

$$\left[\frac{\partial}{\partial x}, \sum_{f_n}\right]: \qquad \frac{\partial}{\partial x} \sum_{k=1}^{\infty} f_k = \sum_{k=1}^{\infty} \frac{\partial}{\partial x} f_k$$

Answer: No.

Counterexample:
$$f_n(x) = \frac{\sin(nx)}{\sqrt{n}} \Rightarrow 0 := f$$
, but $f'_n = \sqrt{n}\cos(nx) \neq 0 = f'$ (at, say, $x = 0$)

Sufficient Condition: When $f_n \in C^1$, $\exists x_0 : f_n(x_0) \to f(x_0)$ and $\sum ||f'_n||_{\infty} < \infty$ (continuously differentiable, converges at a point, and the derivatives absolutely converge)

14.6 Continuity

Definition 14.6.1 (Limit definition of continuity).

$$f$$
 continuous $\iff \lim_{x \to p} f(x) = f(p)$

Definition 14.6.2 (ε - δ definition of continuity).

$$f:(X,d_X) \to (Y,d_Y)$$
 continuous $\iff \forall \varepsilon, \ \exists \delta \ \middle| \ d_X(x,y) < \delta \implies d_Y(f(x),f(y)) < \varepsilon$

Example 14.6.1 (A nonobviously discontinuous function):

$$f(x) = \sin\left(\frac{1}{x}\right) \implies 0 \in D(f)$$

Proof (?).

 $\binom{\text{todo}}{}$

Example 14.6.2 (The Dirichlet function): The Dirichlet function is nowhere continuous:

$$f(x) = \mathbb{1} \left[\mathbb{Q} \right]$$

Proposition 14.6.1(Thomae's function: the set of points of continuity and of discontinuity can both be infinite).

The following function continuous at infinitely many points and discontinuous at infinitely many points:

$$f(x) = \begin{cases} 0 & x \in \mathbb{R} \setminus \mathbb{Q} \\ \frac{1}{q} & x = \frac{p}{q} \in \mathbb{Q} \end{cases}$$

Then f is discontinuous on \mathbb{Q} and continuous on $\mathbb{R} \setminus \mathbb{Q}$.

Proof (?).

f is continuous on \mathbb{Q} :

• Fix ε , let $x_0 \in \mathbb{R} - \mathbb{Q}$, choose $n : \frac{1}{n} < \varepsilon$ using Archimedean property.

14.6 Continuity 97

- Define
$$S = \left\{ x \in \mathbb{Q} : x \in (0,1), x = \frac{m}{n'}, n' < n \right\}$$

- Then $|S| \le 1 + 2 + \dots + (n-1)$, so choose $\delta = \min_{s \in S} |s - x_0|$

$$x \in N_{\delta}(x_0) \implies f(x) < \frac{1}{n} < \varepsilon.$$

f is discontinuous on $\mathbb{R} \setminus \mathbb{Q}$:

• Let
$$x_0 = \frac{p}{q} \in \mathbb{Q}$$
 and $\{x_n\} = \left\{x - \frac{1}{n\sqrt{2}}\right\}$. Then
$$x_n \uparrow x_0 \text{ but } f(x_n) = 0 \to 0 \neq \frac{1}{q} = f(x_0)$$

Remark 14.6.1 : There are no functions that are continuous on $\mathbb Q$ but discontinuous on $\mathbb R-\mathbb Q$

Definition 14.6.3 (Uniform Continuity).

todo

Definition 14.6.4 (Absolute Continuity).

Theorem 14.6.1 (Extreme Value Theorem).

A continuous function on a compact space attains its extrema.

14.6.1 Lipschitz Continuity

14.7 Differentiability

$$f'(p) := \frac{\partial f}{\partial x}(p) = \lim_{x \to p} \frac{f(x) - f(p)}{x - p}$$

- For multivariable functions: existence and continuity of $\frac{\partial \mathbf{f}}{\partial x_i} \forall i \implies \mathbf{f}$ differentiable
 - Necessity of continuity: example of a continuous functions with all partial and directional

14.7 Differentiability 98

derivatives that is not differentiable:

$$f(x,y) = \begin{cases} \frac{y^3}{x^2 + y^2} & (x,y) \neq (0,0) \\ 0 & \text{else} \end{cases}.$$

14.7.1 Properties, strongest to weakest

$$C^{\infty} \subseteq C^k \subseteq \text{ differentiable } \subseteq C^0 \subset \mathcal{R}_K$$
.

- Example showing $f \in C^0 \implies f$ is differentiable and f not differentiable $\implies f \notin C^0$.
 - Take f(x) = |x| at x = 0.
- Example showing that f differentiable $\implies f \in C^1$:
 - Take

$$f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & x \neq 0 \\ 0 & x = 0 \end{cases} \implies f'(x) = \begin{cases} -\cos\left(\frac{1}{x}\right) + 2x\sin\left(\frac{1}{x}\right) & x \neq 0 \\ 0 & x = 0 \end{cases}$$


but $\lim_{x\to 0} f'(x)$ does not exist and thus f' is not continuous at zero.

Proof that f differentiable $\implies f \in C^0$:

$$f(x) - f(p) = \frac{f(x) - f(p)}{x - p} (x - p) \stackrel{\text{hypothesis}}{=} f'(p) (x - p) \stackrel{x \to p}{\rightrightarrows} 0$$

14.8 Giant Table of Relations

Bold are assumed hypothesis, regular text is the strongest conclusion you can reach, strikeout denotes implications that aren't necessarily true.



Explanation of items in table:

- K-integrable: compactly integrable.
- f integrable $\implies F$ differentiable $\implies F \in C_0$
 - By definition and FTC, and differentiability \implies continuity
- f differentiable and K compact $\implies f$ integrable on K.
 - In general, f differentiable $\implies f$ integrable. Necessity of compactness:

$$f(x) = e^x \in C^{\infty}(\mathbb{R}) \text{ but } \int_{\mathbb{R}} e^x dx \to \infty.$$

- f integrable $\implies f$ differentiable
 - An integrable function that is not differentiable: f(x) = |x| on \mathbb{R}
- f differentiable $\implies f$ continuous a.e.

14.9 Integrability

- Sufficient criteria for Riemann integrability:
 - f continuous
 - -f bounded and continuous almost everywhere, or
 - f uniformly continuous
- f integrable \iff bounded and continuous a.e.

Theorem 14.9.1 (FTC for the Riemann Integral).

If F is a differentiable function on the interval [a, b], and F' is bounded and continuous a.e., then $F' \in L_R([a, b])$ and

$$\forall x \in [a, b] : \int_{a}^{x} F'(t) \ dt = F(x) - F(a)$$

Suppose f bounded and continuous a.e. on [a, b], and define

$$F(x) := \int_{a}^{x} f(t) dt$$

Then F is absolutely continuous on [a, b], and for $p \in [a, b]$,

$$f \in C^0(p) \implies F$$
 differentiable at $p, F'(p) = f(p)$, and $F' \stackrel{\text{a.e.}}{=} f$.

Proposition 14.9.1.

The Dirichlet function is Lebesgue integrable but not Riemann integrable:

$$f(x) = \mathbb{1} [x \in \mathbb{Q}]$$

14.9 Integrability

Proof(?).

todo

14.10 List of Free Conclusions:

- f integrable on $U \Longrightarrow$:
 - f is bounded
 - -f is continuous a.e. (finitely many discontinuities)
 - $-\int f$ is continuous
 - $-\int f$ is differentiable
- f continuous on U:
 - $-\ f$ is integrable on compact subsets of U
 - f is bounded
 - f is integrable
- f differentiable at a point p:
 - f is continuous
- f is differentiable in U
 - -f is continuous a.e.
- Defining the Riemann integral: #todo

14.11 Convergence

14.11.1 Sequences and Series of Functions

Definition 14.11.1 (Convergence of an infinite series). Define

$$s_n(x) \coloneqq \sum_{k=1}^n f_k(x)$$

and

$$\sum_{k=1}^{\infty} f_k(x) := \lim_{n \to \infty} s_n(x),$$

which can converge pointwise, absolutely, uniformly, or not all.

Proposition 14.11.1(?).

If $\limsup |f_k(x)| \neq 0$ then f_k is not convergent.

Proposition 14.11.2(?).

If f is injective, then f' is nonzero in some neighborhood of ???

14.11.2 Pointwise convergence

$$f_n \to f = \lim_{n \to \infty} f_n.$$

Summary:

$$\lim_{f_n \to f} \lim_{x_i \to x} f_n(x_i) \neq \lim_{x_i \to x} \lim_{f_n \to f} f_n(x_i).$$

$$\lim_{f_n \to f} \int_I f_n \neq \int_I \lim_{f_n \to f} f_n.$$

Proposition 14.11.3(?).

Pointwise convergence is strictly weaker than uniform convergence.

 $f_n(x) = x^n$ on [0, 1] converges pointwise but not uniformly.

- Towards a contradiction let $\varepsilon = \frac{1}{2}$.
- Let $n = N\left(\frac{1}{2}\right)$ and $x = \left(\frac{3}{4}\right)^{\frac{1}{n}}$. Then f(x) = 0 but

$$|f_n(x) - f(x)| = x^n = \frac{3}{4} > \frac{1}{2}$$

Proposition 14.11.4(A pointwise limit of continuous functions is not necessarily continuous.).

 f_n continuous $\implies f := \lim_n f_n$ is continuous.

14.11 Convergence 102 Proof (?).

Take

$$f_n(x) = x^n, \quad f_n(x) \to 1 \text{ [[] } x = 1\text{]}.$$

Proposition 14.11.5 (The limit of derivatives need not equal the derivative of the limit).

$$f_n$$
 differentiable $\implies f'_n$ converges f'_n converges $\implies \lim f'_n = f'$.

Proof (?).

Take

$$f_n(x) = \frac{1}{n}\sin(n^2x) \to 0,$$
 but $f'_n = n\cos(n^2x)$ does not converge.

Proposition 14.11.6(?).

$$f_n \in \mathcal{R}_I \implies \lim_{f_n \to f} \int_I f_n \neq \int_I \lim_{f_n \to f} f_n.$$

Proof(?).

May fail to converge to same value, take

$$f_n(x) = \frac{2n^2x}{(1+n^2x^2)^2} \to 0$$
 but $\int_0^1 f_n = 1 - \frac{1}{n^2+1} \to 1 \neq 0$.

14.11.3 Uniform Convergence

Notation:

$$f_n \rightrightarrows f = \lim_{n \to \infty} f_n \text{ and } \sum_{n=1}^{\infty} f_n \rightrightarrows S.$$

_

14.11 Convergence 103

Summary:

$$\lim_{x_i \to x} \lim_{f_n \to f} f_n(x_i) = \lim_{f_n \to f} \lim_{x_i \to x} f_n(x_i) = \lim_{f_n \to f} f_n(\lim_{x_i \to x} x_i).$$

$$\lim_{f_n \to f} \int_I f_n = \int_I \lim_{f_n \to f} f_n.$$

$$\sum_{n=1}^{\infty} \int_{I} f_n = \int_{I} \sum_{n=1}^{\infty} f_n.$$

"The uniform limit of a(n) x function is x", for $x \in \{\text{continuous}, \text{bounded}\}$

• Equivalent to convergence in the uniform metric on the metric space of bounded functions on X:

$$f_n \rightrightarrows f \iff \sup_{x \in X} |f_n(x) - f(x)| \to 0.$$

- $(B(X,Y), \|\|_{\infty})$ is a metric space and $f_n \rightrightarrows f \iff \|f_n f\|_{\infty} \to 0$ (where B(X,Y) are bounded functions from X to Y and $\|f\|_{\infty} = \sup_{x \in I} \{f(x)\}$
- $f_n \rightrightarrows f \implies f_n \to f$ pointwise
- f_n continuous $\implies f$ continuous
 - − i.e. "the uniform limit of continuous functions is continuous"
- $f_n \in C^1$, $\exists x_0 : f_n(x_0) \to f(x_0)$, and $f'_n \rightrightarrows g \implies f$ differentiable and f' = g (i.e. $f'_n \to f'$)
 - Necessity of \mathbb{C}^1 look at failures of f_n' to be continuous:

$$\diamondsuit$$
 Take $f_n(x) = \sqrt{\frac{1}{n^2} + x^2} \Longrightarrow |x|$, not differentiable

$$\diamondsuit$$
 Take $f_n(x) = n^{-\frac{1}{2}} \sin(nx) \Rightarrow 0$ but $f'_n \not\to f' = 0$ and $f' \neq g$

- f_n integrable $\implies f$ integrable and $\int f_n \to \int f$
- f_n bounded $\Longrightarrow f$ bounded
- $f_n \rightrightarrows f_n \implies f'_n$ converges
 - Says nothing about it general

- $f'_n \rightrightarrows f' \implies f_n \rightrightarrows f$
 - Unless f converges at one or more points.

Proposition 14.11.7 (All subsequences of a convergent sequence share a limit). $\{x_i\} \to p \implies$ every subsequence also converges to p.

Definition 14.11.2 (Cauchy Sequence).

todo

Proposition 14.11.8(?).

Every convergent sequence in X is a Cauchy sequence.

Remark 14.11.1: The converse need not hold in general, but if X is complete, every Cauchy sequence converges. An example of a Cauchy sequence that doesn't converge: take $X = \mathbb{Q}$ and set $x_i = \pi$ truncated to i decimal places.

 $Remark\ 14.11.2:$ If any subsequence of a Cauchy sequence converges, the entire sequence converges.

Definition 14.11.3 (Metric).

$$d(x,y) \geq 0$$
 Positive $d(x,y) = 0 \iff x = y$ Nondegenerate $d(x,y) = d(y,x)$ Symmetric $d(x,y) \leq d(x,p) + d(p,y)$ $\forall p$ Triangle Inequality.

Definition 14.11.4 (Complete).

todo

 $\textbf{Definition 14.11.5} \ (\text{Bounded}).$

todo

14.12 Topology



14.11 Convergence 105

Definition 14.12.1 (Axioms for a Topology).

Open Set Characterization: Arbitrary unions and finite intersections of open sets are open. **Closed Set Characterization**: Arbitrary intersections and finite unions of closed sets are closed.

Remark 14.12.1: The best source of examples and counterexamples is the open/closed unit interval in \mathbb{R} . Always test against these first!

Remark 14.12.2: If f is a continuous function. the preimage of every open set is open and the preimage of every closed set is closed.

Proposition 14.12.1(?).

In \mathbb{R} , singleton sets and finite discrete sets are closed.

Proof (?).

A singleton set can be written

$$\{p_0\} = (-\infty, p) \cup (p, \infty).$$

A finite discrete set $\{p_0\}$, which wlog (by relabeling) can be assumed to satisfy $p_0 < p_1 < \cdots$, can be written

$$\{p_0, p_1, \dots, p_n\} = (-\infty, p_0) \cup (p_0, p_1) \cup \dots \cup (p_n, \infty).$$

Proposition 14.12.2(?).

This yields a good way to produce counterexamples to continuity.

In \mathbb{R} , singletons are closed. This means any finite subset is closed, as a finite union of singleton sets!

Proposition 14.12.3(?).

If X is a compact metric space, then X is complete and bounded.

Proposition 14.12.4(?).

If X complete and $X \subset Y$, then X closed in Y.

Remark 14.12.3: The converse generally does not hold, and completeness is a necessary condition. Counterexample: $\mathbb{Q} \subset \mathbb{Q}$ is closed but $\mathbb{Q} \subset \mathbb{R}$ is not.

14.12 Topology 106

Proposition 14.12.5(?).

If X is compact, then $Y \subset X \implies Y$ is compact $\iff Y$ closed.

Definition 14.12.2 (Sequential Compactness).


A topological space X is **sequentially compact** iff every sequence $\{x_n\}$ has a subsequence converging to a point in X.

Proposition 14.12.6 (Compactness and sequential compactness).

If X is a metric space, X is compact iff X is sequentially compact.

Remark 14.12.4: Note that in general, neither form of compactness implies the other.

14.13 Counterexamples



There are functions differentiable only at a single point. Example:

$$f(x) = \begin{cases} x^2 & x \in QQ \\ -x^2 & x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}.$$

This is discontinuous everywhere except for x = 0, and you can compute

$$\lim_{h \to 0} \frac{f(x+h) - f(x)}{h} \Big|_{x=0} = \lim_{h \to 0} \begin{cases} h & x \in \mathbb{Q} \\ -h & x \in \mathbb{R} \setminus \mathbb{Q} \end{cases} = 0.$$

Proposition 14.13.2(?).

The product of two non-differentiable functions can be differentiable: take f(x) = g(x) = |x| which are not differentiable at x = 0, then $fg(x) = |x|^2$ is differentiable at x = 0.

Proposition 14.13.3(?).

A continuous function that is zero on a dense set $A \subset X$ is identically zero.

Proof(?).

Since A is dense, for any $x \in X \setminus A$ take a sequence $\{x_n\}$ in A converging to x. Then $0 = f(x_n) \to f(x)$ implies f(x) = 0.

14.13 Counterexamples 107

15 | Point-Set Topology

~

15.1 Definitions

Bring in Rudin's list

• Epsilon-neighborhood

$$- N_r(p) = \left\{ q \mid d_X(p,q) < r \right\}$$

- Limit Point
 - p is a limit point of E iff $\forall N_r(p), \exists q \neq p \mid q \in N_r(p)$
 - Equivalently, $\forall N_r(p), N_r(p) \cap E \neq \emptyset$
 - Let L(E) be the set of limit points of E.
 - Example: $E = (0,1) \implies 0 \in L(E)$
- Isolated Point
 - -p is an isolated point of E iff p is not a limit point of E
 - Equivalently, $\exists N_r(p) \mid N_r(p) \cap E = \emptyset$
 - Equivalently, E L(E)
- Perfect
 - -E is perfect iff E is closed and $E \subseteq L(E)$
 - Equivalently, L(E) = E
- Interior
 - p is an interior point of E iff $\exists N_r(p) \mid N_r(p) \subsetneq E$
 - Denote the interior of E by E°
- Exterior
- Closed sets
 - -E is closed iff p a limit point of $E \implies p \in E$
 - Equivalently if $L(E) \subseteq E$
 - Closed under finite unions, arbitrary intersections

- Open sets
 - -E is open iff $p \in E \implies p \in E^{\circ}$ Equivalently, if $E \subseteq E^{\circ}$

 - Closed under arbitrary unions, finite intersections
- Boundary
- Closure
- Dense
 - -E is dense in X iff $X \subseteq E \cup L(E)$
- Connected
 - Space of connected sets closed under union, product, closures
 - Convex \implies connected
- Disconnected
- Path Connected

$$\begin{array}{ll} - \ \forall x,y \in X \exists f: I \to X \ \Big| \ f(0) = x, f(1) = y \\ - \ \text{Path connected} \implies \text{connected} \end{array}$$

- Simply Connected
- Totally Disconnected
- Hausdorff
- Compact
 - Every covering has a finite subcovering.
 - -X compact and $U \subset X : (U \text{ closed} \implies U \text{ compact})$
 - $\Diamond U \text{ compact } \Longrightarrow U \text{ closed } \text{iff } X \text{ is Hausdorff}$
 - Closed under products

Example 15.1.1 (?): The space
$$\left\{\frac{1}{n}\right\}_{n\in\mathbb{N}}$$
.

List of properties preserved by continuous maps:

• Connectedness


• Compactness

Checking if a map is homeomorphism:

• f continuous, X compact and Hausdorff $\implies f$ is a homeomorphism.

16 Probability





Definition 16.1.1 (Random Variable).

For (Σ, E, μ) a probability space with sample space Σ and probability measure μ , a random variable is a function $X : \Sigma \to \mathbb{R}$

Definition 16.1.2 (Probability Density Function (PDF)).

For any $U \subset \mathbb{R}$, given by the relation

$$P(X \in U) = \int_{U} f(x) \ dx$$

$$\implies P(a \le X \le b) = \int_{a}^{b} f(x) \ dx$$

 $\textbf{Definition 16.1.3} \ (\textbf{Cumulative Distribution Function} \ (\textbf{CDF})).$

The antiderivative of the PDF

$$F(x) = P(X \le x) = \int_{-\infty}^{x} f(x) \ dx.$$

Probability 110

Yields
$$\frac{\partial F}{\partial x} = f(x)$$

Definition 16.1.4 (Mean/Expected Value).

$$E[X] := \langle \mathrm{id}, f \rangle = \int_{\mathbb{R}} x f(x) dx.$$

Also denoted μ_X .

Proposition 16.1.1 (Linearity of Expectation).

$$E\left[\sum_{i\in\mathbb{N}}a_iX_i\right] = \sum_{i\in\mathbb{N}}a_iE[X_i].$$

Does not matter whether or not the X_i are independent.

Definition 16.1.5 (Variance).

$$Var(X) = E[(X - E[X])^{2}]$$

$$= \int (x - E[X])^{2} f(x) dx$$

$$= E[X^{2}] - E[X]^{2}$$

$$\coloneqq \sigma^{2}(X)$$

where σ is the standard deviation. Can also defined as $\langle (\mathrm{id} - \langle \mathrm{id}, f \rangle)^2, f \rangle$ Take the portion of the id function in the orthogonal complement of f, squared, and project it back onto f?

Proposition 16.1.2 (Properties of Variance).

$$\operatorname{Var}(aX + b) = a^{2}\operatorname{Var}(X)$$

$$\operatorname{Var}\left(\sum_{\mathbb{N}} X_{i}\right) = \sum_{i} \operatorname{Var}(X_{i}) + 2\sum_{i < j} \operatorname{Cov}(X_{i}, X_{j}).$$

Definition 16.1.6 (Covariance).

16.1 Definitions

$$Cov(X,Y) = E[(X - \mu_X)(Y - \mu_Y)]$$
$$= E[XY] - E[X]E[Y]$$

Proposition 16.1.3 (Properties of Covariance).

$$Cov(X, X) = Var(X)$$

$$Cov(aX, Y) = aCov(X, Y)$$

$$Cov(\sum_{\mathbb{N}} X_i, \sum_{\mathbb{N}} Y_j) = \sum_{i} \sum_{j} Cov(X_i, Y_j)$$

Proposition 16.1.4 (Stirling's Approximation).

$$k! \sim k^{\frac{k+1}{2}} e^{-k} \sqrt{2\pi}.$$

Proposition 16.1.5 (Markov Inequality).

$$P(X \ge a) \le \frac{1}{a} E[X]$$

One-sided Markov:

$$P(X \in N_{\varepsilon}(\mu)) = 2\frac{\sigma^2}{\sigma^2 + a^2}.$$

Proposition 16.1.6 (Chebyshev's Inequality).

$$P(|X - \mu| \ge a) \le \left(\frac{\sigma}{k}\right)^2$$

Proof (?).

Apply Markov to the variable $(X - \mu)^2$ and $a = k^2$

Theorem $16.1.1(Central\ Limit\ Theorem).$

For X_i i.i.d.,

$$\lim_{n} \frac{\sum_{i=1}^{n} X_i - n\mu}{\sigma\sqrt{n}} \sim N(0, 1).$$

16.1 Definitions 112

Theorem 16.1.2 (Strong Large of Large Numbers).

$$P(\frac{1}{n}\sum X_i \to \mu) = 1.$$

Proposition 16.1.7 (Chernoff Bounds).

For all t > 0,

$$P(X \in N_{\varepsilon}(a)^c) \le 2e^{-at}M_X(t)$$

Proposition 16.1.8 (Jensen's Inequality).

$$E[f(X)] \ge f(E[X])$$

Definition 16.1.7 (Entropy).

$$H(X) = -\sum p_i \ln p_i$$

16.2 Theory and Background

 $\textbf{Definition 16.2.1} \ (\text{Axioms for a Probability Space}).$

Given a sample space Σ with events S, 1. $\mu(\Sigma) = 1$ 1. Yields $S \in \Sigma \implies 0 \le P(S) \le 1$ 2. For mutually exclusive events, $P(\cup_{\mathbb{N}} S_i) = \sum_{\mathbb{N}} P(S_i)$ 1. Yields $P(\emptyset) = 0$

Proposition 16.2.1 (Properties that follow from axioms).

- $P(S^c) = 1 P(S)$
- $E \subseteq F \implies P(E) \le P(F)$
- Proof: $E \subseteq F \implies F = E \cup (E^c \cap F)$, which are disjoint, so $P(E) \leq P(E) + P(E^c \cap F) = P(F)$.
- $P(E \cup F) = P(E) + P(F) P(E \cap F)$

Definition 16.2.2 (Conditional Probability).

$$P(F)P(E \mid F) = P(E \cap F) = P(E)P(F \mid E)$$

Generalization:

$$P(\cap_{\mathbb{N}} E_i) = P(E_1)P(E_2 \mid E_1)P(E_3 \mid E_1 \cap E_2) \cdots$$

Theorem 16.2.1 (Bayes' Rule).

$$P(E) = P(F)P(E \mid F) + P(F^{c})P(E \mid F^{c})$$
$$P(E) = \sum_{i} P(A_{i})P(E \mid A_{i})$$

Generalization: for $\coprod_{i=1}^{n} A_i = \Sigma$ and $A = A_i$ for some i,

$$P(A \mid B) = \frac{P(A)P(B \mid A)}{\sum_{j=1}^{n} P(B \mid A_j)}.$$

The LHS: the posterior probability, while $P(A_i)$ are the priors.

Definition 16.2.3 (Odds).

$$P(A)/P(A^c)$$

Conditional odds:

$$\frac{P(A \mid E)}{P(A^c \mid E)} = \frac{P(A)}{P(A^c)} \frac{P(E \mid A)}{P(E \mid A^c)}$$

Definition 16.2.4 (Independence).

$$P(A \cap B) = P(A)P(B)$$

Proposition 16.2.2 (Change of Variables for PDFs).

If g is differentiable and monotonic and Y = g(X), then

$$f_Y(y) = \begin{cases} (f_X \circ g^{-1})(y) \middle| \frac{\partial}{\partial y} g^{-1}(y) \middle| & y \in \text{im}(g) \\ 0 & \text{else} \end{cases}$$

Proposition 16.2.3 (PDF for a sum of independent random variables).

$$f_{X+Y} = (F_X * f_y)$$

16.3 Distributions

Let X be a random variable, and f be its probability density function satisfying f(k) = P(X = k)

16.3.1 Uniform

• Consider an event with n mutually exclusive outcomes of equal probability, and let $X \in \{1, 2, ..., n\}$ denote which outcome occurs. Then,

$$f(k) = \frac{1}{n}$$
$$\mu = \frac{n}{2}$$
$$\sigma^2 = a$$

- Examples:
 - Dice rolls where n = 6.
 - Fair coin toss where n=2.
- Continuous: $\mu = (1/2)(b+a), \sigma^2 = (1/12)(b-a)^2$

16.3.2 Bernoulli

• Consider a trial with either a positive or negative outcome, and let $X \in \{0, 1\}$ where 1 denotes a success with probability p. Then,

$$f(k) = \begin{cases} 1 - p, & k = 0 \\ p, & k = 1 \end{cases}$$
$$\mu = p$$
$$\sigma^2 = p(1 - p)$$

- Examples: - A weighted coin with P(Heads) = p

16.3.3 Binomial

• Consider a sequence of independent Bernoulli trials, let $X \in \{1, ..., n\}$ denote the number of successes occurring in n trials. Then,

$$f(k) = \binom{n}{k} p^k (1-p)^{n-k}$$
$$\mu = np$$
$$\sigma^2 = np(1-p)$$

- Examples:
 - A sequence of coin flips and the numbers of total heads occurring.

16.3.4 Poisson

• Given a parameter $\lambda > 0$ that denotes the rate per unit time of an event occurring and X the number of times the event occurs in one unit of time,

$$f(k) = \frac{\lambda^k}{k!} e^{-\lambda}$$
$$\mu = \lambda$$
$$\sigma^2 = \lambda^2$$

• Approximates binomial when n >> 1 and p << 1 by using $\lambda = np$

16.3.5 Negative Binomial

• $B^-(r,p)$: similar to binomial, where X is the number of trials it takes to accumulate r successes

$$f(k) = {k-1 \choose r-1} p^r (1-p)^{k-r}$$
$$\mu = \frac{r}{p}$$
$$\sigma^2 = \frac{r(1-p)}{p^2}$$

16.3.6 Geometric

• Consider a sequence of independent Bernoulli trials, let $X \in \{1, ..., n\}$ where X = k denotes the first success happening on the k-th trial. Then,

$$f(k) = (1-p)^{k-1}p$$

$$\mu = \frac{1}{p}$$

$$\sigma^2 = \frac{1-p}{p^2}$$

- Note this is equal to $B^-(1,p)$
- Examples:
 - A sequence of coin flips and the number of flips before the first heads appears.

16.3.7 Hypergeometric

• H(n, m, s) An urn filled with n balls, where m are white and n - m are black; pick a sample of size s and let X denote the number of white balls:

$$f(k) = {m \choose k} {n-m \choose s-k} {n \choose s}^{-1}$$

$$\mu = \frac{ms}{n}$$

$$\sigma^2 = \frac{ms}{n} (1 - \frac{m}{n}) \left(1 - \frac{s-1}{n-1}\right)$$

16.3.8 Normal

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

$$\begin{array}{c|cc} z & \Phi(z) \\ \hline 0 & 0.5 \\ 1 & 0.69 \\ 1.5 & 0.84 \\ 2 & 0.93 \\ 2.5 & 0.97 \\ > 3 & 0.99 \\ \hline \end{array}$$

16.3 Distributions

16.4 Table of Distributions

Table: let q = 1 - p.

Distribution f(x)	μ	$\sigma^2 M(t)$
$B(n,p)\binom{n}{x}p^xq^{n-x}$	np	$npq(pe^t + q)^n$
$P(\lambda) \frac{\lambda^x}{x!} e^{-\lambda}$	λ	$\lambda e^{\lambda(e^t-1)}$
$G(p)q^{x-1}p$	$\frac{1}{p}$	$\frac{q}{p^2} \frac{pe^t}{1 - qe^t}$
$B^{-}(r,p)\binom{n-1}{r-1}p^{r}q^{n-r}$	$rac{r}{p}$	$\frac{rq}{p^2} \left(\frac{pe^t}{1 - qe^t} \right)^r$
$U(a,b)1 [a \le x \le b] \frac{1}{b-a}$	$\frac{1}{2}(a+b)$	$\frac{1}{12}(b-a)^2 \frac{e^{tb} - e^{ta}}{t(b-a)}$
$Exp(\lambda)\mathbb{1}\left[0 \le x\right]\lambda e^{-\lambda x}$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}\frac{\lambda}{\lambda-t}$
$\Gamma(s,\lambda)\mathbb{1}\left[0 \le x\right] \frac{\lambda e^{-\lambda x}(\lambda x)^{s-1}}{\Gamma(s)}$	$rac{s}{\lambda}$	$rac{s}{\lambda^2}\left(rac{\lambda}{\lambda-t} ight)^s$
$N(\mu, \sigma^2) \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	μ	$\sigma^2 e^{\mu t + \frac{1}{2}\sigma^2 t^2}$

• Why you need the Stieltjes Integral: let $X \sim B(n, \frac{1}{2}), Y \sim U(0, 1)$, and

$$Z = \begin{cases} X, & X = 1 \\ Y, & else \end{cases}$$

then $|Z|=|\mathbb{R}|$ so Z is not discrete, but $P(X=1)=\frac{1}{2}\neq 0$ so Z is not continuous. Definition:

$$\int_{a}^{b} g(x) \ dF(x) = \lim \sum_{i=1}^{n} g(x_i) (F(x_i) - F(x_{i-1})).$$

16.5 Common Problems



- Birthday Paradox
- Coupon Collectors
 - Given $X = \{1, \dots n\}$, what is the expected number of draws until all n outcomes are seen?

16.6 Notes, Shortcuts, Misc



- When computing expected values, variation, etc, just insert a parameter k and compute the moments $E[X^k]$. Then with a solution in terms of k, let k=1,2 etc.
- Neat property of pdfs: $P(X \in N_{\varepsilon}(a)) \approx \varepsilon f(a)$

Definition 16.6.1 (The Gamma Function).

$$\Gamma(x+1) = \int_{\mathbb{R}^{>0}} e^{-t} t^x dt.$$

Integrate by parts to obtain functional relation $\Gamma(x+1) = x\Gamma(x)$

Proposition 16.6.1 (Boole's Inequality).

$$P(\cup_{\mathbb{N}} A_i) \le \sum_{\mathbb{N}} P(A_i)$$

• For any function $f: X \to \mathbb{R}$, there is a lower bound: $\max_{x \in X} f(x) \ge E[f(x)]$

Definition 16.6.2 (Moment Generating Functions).


Define

$$M(t) = E[e^{Xt}]$$

- Then $M^{(n)}(0)$ is the *n*-th moment, i.e. $M'(0) = E[X], M''(0) = E[X^2]$, etc.

- $M_{X+Y}(t) = M_X(t)M_Y(t)$ (if independent) $M_{aX+b}(t) = e^{bt}M_X(at)$ $f_X = \mathcal{F}^{-1}(M_X(it))$, denoting the inverse Fourier transform,

17 Combinatorics



$$S_n = \{1, 2, \dots n\}$$
 the symmetric group
$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$
 binomial coefficient
$$n^{\underline{k}} = n(n-1)\cdots(n-k+1) = k!\binom{n}{k}$$
 falling factorial
$$n^{\overline{k}} = n(n+1)\cdots(n+k-1) = k!\binom{n+n-1}{n}$$
 rising factorial
$$\binom{n}{m_1, m_2, \cdots m_k} = \frac{n!}{\prod_{i=1}^k m_i!}$$
 multinomial coefficient

Note that the rising and falling factorials always have exactly k terms.

Multinomial: A set of n items divided into k distinct, disjoint subsets of sizes $m_1 \cdots m_k$. Multinomial theorem:

$$(x_1 + x_2 + \dots + x_k)^n = \sum_{\substack{m_1, m_2, \dots, m_k \\ \sum m_i = n}} \binom{n}{m_1, m_2, \dots, m_k} x_1^{m_1} x_2^{m_2} \dots x_k^{m_k}$$

which contains $\binom{n+r-1}{r-1}$ terms.

Inclusion-Exclusion:

$$|\cup_{i=1}^{n} A_{i}| = \sum_{i} |A|_{i} - \sum_{i_{1} < i_{2}}^{\binom{n}{2^{2}}} |A_{i_{1}} \cap A_{i_{2}}| + \sum_{i_{1} < i_{2} < i_{3}}^{\binom{n}{2^{3}}} |A_{i_{1}} \cap A_{i_{2}} \cap A_{i_{3}}| + \dots + (-1)^{n+1} |\cap_{i=1}^{n} A_{i}|$$

$$= \sum_{k=1}^{n} \sum_{i_{1} < \dots < i_{k}} (-1)^{k+1} |\cap_{j=1}^{k} A_{i_{j}}|$$

17.2 The Important Numbers

- Binomial Coefficients
 - The Binomial Theorem:

$$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

- Choosing: $\binom{n}{k}$
- Choosing with repetition allowed: $\binom{n+k-1}{k}$

17.1 Notation 120

- Signed Stirling Numbers of the First Kind: s(n,k)
 - Count the number of permutations of n elements with k disjoint cycles, i.e. the number of elements elements in S_n that are the product of k disjoint cycles (including trivial cycles that fix a point).
 - Recurrence relation:

$$s(n,k) = s(n-1, k-1) + ks(n-1, k).$$

- Relation to falling factorial: $x^{\underline{n}} = \sum_{k=1}^{n} s(n,k)x^{k}$
- Stirling Numbers of the Second Kind: $\binom{n}{k}$
 - Counts the number of ways to partition a set N into k non-empty subsets S_i (i.e. such that $S_i \cap S_j = \emptyset$, $\prod_{i=1}^k S_i = N$)
 - Recurrence relation:

$${n+1 \brace k} = k {n \brace k} + {n \brace k-1}$$

$${0 \brace 0} = 1, \quad {n \brace 0} = {0 \brack n} = 0$$

- Explicit formula:
$${n \brace k} = \frac{1}{k!} \sum_{i=0}^{k} (-1)^{k-i} {k \choose i} i^n$$
-
$$B_n = \sum_{i=0}^{n} {n \brace i}$$

- Betti Numbers
- Bell Numbers
- Compositions
 - A composition of n is a way of writing n as a sum of strictly positive integers, ie. $k_1 + k_2 + \cdots + k_i = n$ where each $0 < k_i \le n$, where order matters (and distinct orders count as distinct compositions).
 - Weak compositions: identical, but some terms are allowed to be zero.
 - Number of compositions of n into k parts: $\binom{n-1}{k-1}$
 - Number of weak compositions of n into k parts: $\binom{n+k-1}{n}$
 - Total number of compositions of n (into any number of parts): 2^{n-1}
- Partitions
 - A partition of n is a composition of n quotiented by permutations of the ordering of terms.
 - \diamondsuit Example: 2 compositions of 5 involving 1 and 4, given by 4+1 and 1+4, whereas there is only one such partition of 5 given by 4+1.
 - Visualize with Young diagrams

17.2.1 Common Problems

• Stars and Bars

– No two bars adjacent: $\binom{n-1}{k-1}$ – Allowing adjacent bars: $\binom{n+k-1}{k-1}$

Coupon Collectors Problem

17.2.2 The Twelvefold Way

Consider a function $f: N \to K$ where |N| = n, |K| = k.

A number of valid interpretations: - f labels elements of N by elements of K - For each element of N, f chooses an element of K - f partitions N into classes that are mapped to the same element of K - Throw each of N balls into some of K boxes

Dictionary: - No restrictions: - Assign n labels, repetition allowed - Form a multiset of K of size n - Injectivity - Assign n labels without repetition - Select n distinct elements from K - Number of n-combinations of k elements - No more than one ball per box - Surjectivity: - Use every label at least once - Every element of K is selected at least once - "Indistinguishable" - Quotient by the action of S_n or S_k - n-permutations = injective functions - n-combinations = injective functions / S_n - n-multisets = all functions / S^n - Partitions of N into k subsets = surjective functions / S_k -Compositions of n into k parts = surjective functions / S_n

Permutations Restrictions	$N \xrightarrow{f} K$	$N \hookrightarrow K$	$N \twoheadrightarrow K$
f	k^n	$k^{\underline{n}}$	$k! \begin{Bmatrix} n \\ k \end{Bmatrix}$
$f\circ\sigma_N$	$\binom{n+k-1}{n}$	$\binom{k}{n}$	$\binom{n-1}{n-k}$
$\sigma_X \circ f$	$\sum_{i=0}^{k} {n \brace i}$	$1 [n \le k]$	$\begin{Bmatrix} n \\ k \end{Bmatrix}$
$\sigma_X \circ f \circ \sigma_N$	$p_k(n+k)$	$1 [n \le k]$	$p_k(n)$

In words (todo: explain)

Perm. / Rest.	_	Injective	Surjective
	A sequence of any n elements of K	Sequences of n distinct elements of K	Compositions of N with exactly k subsets
Permutations of N	Multisets of K with n elements	An n -element subset of K	Compositions of n with k terms

Perm. / Rest.	_	Injective	Surjective
Permutations of X	Partitions of N into $\leq k$ subsets	?	Partitions of N into exactly k nonempty subsets
Both	Partitions of n into $\leq k$ parts	?	Partitions of n into exactly k parts

Proofs/Explanations: todo

• Counting non-isomorphic things: Pick a systematic way. Can descend my maximum vertex degree, or ascend by adding nodes/leaves.

18 Complex Analysis

• $\lim_{z\to z_0} f(z) = x_0 + iy_0$ iff the component functions limit to x_0 and y_0 respectively. Moreover, both ways are equal!

Notation: z = a + ib, f(z) = u(x, y) + iv(x, y)





Complex Analysis 123

$$|z| = \sqrt{a^2 + b^2}$$

$$|z|^2 = z\overline{z} = a^2 + b^2$$

$$\frac{z\overline{z}}{|z|^2} = \frac{(a+ib)(a-ib)}{a^2 + b^2} = 1$$

$$\frac{1}{z} = \frac{\overline{z}}{|z|^2} = \frac{a-ib}{a^2 + b^2}$$

$$e^{zx} = e^{(a+ib)x} = e^{ax}(\cos(bx) + i\sin(bx))$$

$$x^z := e^{z\ln x}$$

$$\text{Log}(z) = \ln|z| + i\operatorname{Arg}(z)$$

$$\cos z = \frac{1}{2}(e^{iz} + e^{-iz})$$

$$\sin z = \frac{1}{2i}(e^{iz} - e^{-iz})$$

$$(x-z)(x-\overline{z}) = x^2 - 2\mathcal{R}(z)x + (a^2 + b^2)$$

$$\frac{\partial}{\partial z} = \frac{1}{2}\left(\frac{\partial}{\partial x} - i\frac{\partial}{\partial y}\right)$$

$$\frac{\partial}{\partial \overline{z}} = \frac{1}{2}\left(\frac{\partial}{\partial x} + i\frac{\partial}{\partial y}\right)$$

18.2 Complex Arithmetic and Calculus

• *n*-th roots:

$$e^{\frac{ki}{2\pi n}}, \qquad k = 1, 2, \dots n - 1$$

18.2.1 Complex Differentiability

$$z' = \lim_{h \to 0} \frac{f(z+h) - f(z)}{h}$$

- A complex function that is not differentiable at a point: $f(z)=z/\bar{z}$ at z=0

• Cauchy-Riemann Equations

$$u_x = v_y u_y = -v_x$$

• Alternatively:

$$\begin{split} &-\frac{\partial[}{\partial f}\,]\bar{z}=0\\ &-\langle\nabla u,\;\nabla v\rangle=0\\ &-\Delta u=\Delta v=0\;\text{(both components are harmonic)} \end{split}$$

18.3 Complex Integrals

The main theorem:

$$\oint_C f(z) \ dz = 2\pi i \sum_k \operatorname{Res}(f, z_k)$$

Computing residues:

$$\operatorname{Res}(f, c) = \frac{1}{(n-1)!} \lim_{z \to c} \frac{d^{n-1}}{dz^{n-1}} ((z-c)^n f(z))$$
$$f(z) = \frac{g(z)}{h(z)} \implies \operatorname{Res}(f, c) = \frac{g(c)}{h'(c)}$$

Definitions

- Analytic: differentiable everywhere
- Entire
- Holomorphic
- Meromorphic

Complex Analytic \implies smooth and all derivatives are analytic

Not true in real case, take the everywhere differentiable but not C^1 function

$$f(x) = \begin{cases} -\frac{1}{2}x^2 & x < 0\\ \frac{1}{2}x^2 & x \ge 0 \end{cases}$$

18.4 Definitions

In these notes, C generally denotes some closed contour, \mathbb{H} is the upper half-plane, C_R is a semicircle of radius R in \mathbb{H} , f will denote a complex function.

1. Analytic

f is analytic at z_0 if it can be expanded as a convergent power series in some neighborhood of z_0 .

2. Holomorphic

A function f is holomorphic at a point z_0 if $f'(z_0)$ exists in a neighborhood of z_0 .

(Note - this is more than just being differentiable at a single point!)

Big Theorem: f is a holomorphic complex function iff f is analytic.

3. Meromorphic

Holomorphic, except for possibly a finite number of singularities.

4. Conformal

f is conformal at z_0 if f is analytic at z_0 and $f'(z_0) \neq 0$.

5. Harmonic

A function u(x, y) is harmonic if it satisfies Laplace's equation,

$$\Delta u = u_{xx} + u_{yy} = 0$$

Some other notions to look up:

- Conformal maps
- Analytic
- Theorem: Analytic \implies conformal
- The Sterographic projection. Is it conformal?
- Branch points and branch cuts
- Loxodromic transformations
- Horocycles
- Analytic Continuation
- The complex logarithm
- Mobius transformations
- Curvature
- Angular Excess

18.5 Preliminary Notions

18.4 Definitions

18.5.1 What is the Complex Derivative?

In small neighborhoods, the derivative of a function at a point rotates it by an angle $\Delta\theta$ and scales it by a real number λ according to

$$\Delta\theta = \arg f'(z_0), \ \lambda = |f'(z_0)|$$

18.5.2 nth roots of a complex number

The *n*th roots of z_0 are given by writing $z_0 = re^{i\theta}$, and are

$$\zeta = \left\{ \sqrt[n]{r} \exp \left[i \left(\frac{\theta}{n} + \frac{2k\pi}{n} \right) \right] \mid k = 0, 1, 2, \dots, n - 1 \right\}$$

or equivalently

$$\zeta = \left\{ \sqrt[n]{r} \omega_n^k \mid k = 0, 1, 2, \dots, n - 1 \right\} \text{ where } \omega_n = e^{\frac{2\pi i}{n}}$$

This can be derived by looking at $\left(re^{i\theta+2k\pi}\right)^{\frac{1}{n}}$.

It is also useful to immediately recognize that $z^2 + a = (z - i\sqrt{a})(z + i\sqrt{a})$.

18.5.3 The Cauchy-Riemann Equations

If f(x+iy)=u(x,y)+iv(x,y) or $f(re^{i\theta})=u(r,\theta)+iv(r,\theta)$, then f is complex differentiable if u,v satisfy

$$u_x = v_y$$
 $u_y = -v_x$
 $ru_r = v_\theta$ $u_\theta = -rv_r$

In this case,

$$f'(x+iy) = u_x(x,y) + iv_x(x,y)$$

or in polar coordinates,

$$f'(re^{i\theta}) = e^{i\theta}(u_r(r,\theta) + iv_r(r,\theta))$$

18.5 Preliminary Notions

18.6 Integration



18.6.1 The Residue Theorem

If f is meromorphic inside of a closed contour C, then

$$\oint_C f(z)dz = 2\pi i \sum_{z_k} \operatorname{Res}_{z=z_k} f(z)$$

where $\underset{z=z_k}{\operatorname{Res}} f(z)$ is the coefficient of z^{-1} in the Laurent expansion of f.

If f is analytic everywhere in the interior of C, then $\oint_C f(z)dz = 0$.

If f is meromorphic inside of a contour C and analytic everywhere else, one can equivalently calculate the residue at infinity

$$\oint_C f(z)dz = 2\pi i \sum_{z_k} \underset{z=0}{\text{Res }} z^{-2} f(z^{-1})$$

18.6.2 Computing Residues

18.6.3 Simple Poles

If z_0 is a pole of order m, define $g(z) := (z - z_0)^m f(z)$.

If g(z) is analytic and $g(z_0) \neq 0$, then

$$\operatorname{Res}_{z=z_0} f(z) = \frac{\varphi^{(m-1)}(z_0)}{(m-1)!}$$

In the case where m=1, this reduces to

$$\operatorname{Res}_{z=z_0} f(z) = \varphi(z_0)$$

To compute residues this way, attempt to write f in the form

$$f(z) = \frac{\varphi(z)}{(z - z_0)^m}$$

where φ only needs to be analytic at z_0 .

18.6 Integration

18.6.4 Rational Functions

If
$$f(z) = \frac{p(z)}{q(z)}$$
 where

- 1. $p(z_0) \neq 0$
- 2. $q(z_0) = 0$
- 3. $q'(z_0) \neq 0$

then the residue can be computed as

$$\operatorname{Res}_{z=z_0} \frac{p(z)}{q(z)} = \frac{p(z_0)}{q'(z_0)}$$

18.6.5 Computing Integrals

When computing real integrals, the following contours can be useful:

One often needs bounds, which can come from the following lemmas

The Arc Length Bound If $|f(z)| \leq M$ everywhere on C, then

$$\left| \oint_C f(z)dz \right| \le ML_C$$

where L_C is the length of C.

Jordan's Lemma: If f is analytic outside of a semicircle C_R and $|f(z)| \leq M_R$ on C_R where $M_R \to 0$, then

$$\int_{C_R} f(z) e^{iaz} dz \to 0$$

.

Can also be used for integrals of the form $\int f(z) \cos az dz$ or $\int f(z) \sin az dz$, just take real/imaginary parts of e^{iaz} respectively.

18.7 Conformal Maps



1. Linear Fractional Transformations:

$$f(z) = \frac{az+b}{cz+d} \qquad f^{-1}(z) = \frac{-dz+b}{cz-a}$$

2. $[z_1, z_2, z_3] \mapsto [w_1, w_2, w_3]$

Every linear fractional transformation is determined by its action on three points. Given 3 pairs points $z_i \mapsto w_i$, construct one using the implicit equation

$$\frac{(w-w_1)(w_2-w_3)}{(w-w_3)(w_2-w_1)} = \frac{(z-z_1)(z_2-z_3)}{(z-z_3)(z_2-z_1)}$$

3. z^k : Wedge $\mapsto \mathbb{H}$

Just multiplies the angle by k. If a wedge makes angle θ , use $z^{\frac{\pi}{\theta}}$.

It is useful to know that $z \mapsto z^2$ is equivalent to $(x,y) \mapsto (x^2 - y^2, 2xy)$.

4. $e^z: \mathbb{C} \to \mathbb{C}$

the countainer of the tho regions are marened




FIGURE 126

 $w = \exp z$.

Figure 2: image




Figure 3: image




Figure 4: image

5. $\log : \mathbb{H} \to \mathbb{R} + i[0, \pi]$

Just the inverse of what the exponential map does.

Rays	\mapsto	Horizontal Lines
Wedges	\mapsto	Horizontal Strips




Figure 5: $z \mapsto \log z$

6.
$$\sin: [0, \pi/2] + i\mathbb{R} \mapsto \mathbb{H}_{\mathcal{R}(z)>0}$$

Maps the infinite strip to the first quadrant.




Figure 6: $z \leftarrow \sin w$

7.
$$z \mapsto \frac{i-z}{i+z} : \mathbb{H} \mapsto D^{\circ}$$
.

 $\begin{array}{ccc} \mathbb{R}_{>0} & \mapsto & \text{Upper half of } D^{\circ} \\ \mathbb{R}_{<0} & \mapsto & \text{Bottom half of } D^{\circ} \end{array}$

Has inverse $w \mapsto i \frac{1-w}{1+w}$

8.
$$z \mapsto z + z^{-1} : \partial D \mapsto \mathbb{R}$$




Figure 7: $z \mapsto z + z^{-1}$

Maps the boundary of the circle to the real axis, and the plane to \mathbb{H} .

18.7.1 Applications

It is mostly important to know that composing a harmonic function on one domain with an analytic function produces a new harmonic function on the new domain.

Similarly, composing the solution to a boundary value problem on a domain with a conformal map produces a new solution to a new boundary problem in the new domain, where the new boundary is given by the conformal image of the old one.

The general technique is use solutions to the boundary value problem on a simple domain D, and compose one or several conformal maps to map a given problem into D, then pull back the solution.

Heat Flow: Steady Temperatures Generally interested in finding a harmonic function T(x, y) which represents the steady-state temperature at any point. Usually given as a Dirichlet problem on a domain D of the form




Figure 8: image

$$\Delta T = 0$$
$$T(\partial D) = f(\partial D)$$

where f is a given function that prescribes values on ∂D , the boundary of D.

Embed this in an analytic function with its harmonic conjugate to yield solutions of the form F(x+iy) = T(x,y) + iS(x,y).

The **isotherms** are given by T(x, y) = c.

The lines of flow are given by S(x,y) = c.




Figure 9: image

Any easy solution on the domain $\mathbb{R} \times i[0,\pi]$ in the u,v plane, where

$$T(x,0) = 0$$
$$T(x,\pi) = 1$$

is given by $T(u, v) = \frac{1}{\pi}v$.

It is harmonic, as the imaginary part of the analytic $F(u+iv) = \frac{1}{\pi}(u+iv)$, since every analytic function has harmonic component functions.

Similar methods work with different domains, just pick a smooth interpolation between the boundary conditions.




Figure 10: image

Fluid Flow Write $F(z) = \varphi(x,y) + i\psi(x,y)$. Then F is the complex potential of the flow, $\overline{F'}$ is the velocity, and setting $\psi(x,y) = c$ yields the streamlines.

A solution in \mathbb{H} is F(z) = Az some some velocity A. Apply conformal mapping appropriately.




Figure 11: image

18.7.2 Theorems

General Theorems

1. Liouville's Theorem:

If f is entire and bounded on \mathbb{C} , then f is constant.

- 2. If f is continuous in a region D, f is bounded in D.
- 3. If f is differentiable at z_0 , f is continuous at z_0 .

Note - the converse need not hold!

4. If f = u + iv, where u, v satisfy the Cauchy-Riemann equations **and** have continuous partials, then f is differentiable.

Note - continuous partials are not enough, consider $f(z) = |z|^2$.

5. Rouché's Theorem

If p(z) = f(z) + g(z) and |g(z)| < |f(z)| everywhere on C, then f and p have the same number of zeros with C.

6. The Argument Principle

If f is analytic on a closed contour C and meromorphic within C, then

$$W := \frac{1}{2\pi} \Delta_C \arg f(z) = Z - P$$

Proof: Evaluate the integral $\oint_C \frac{f'(z)}{f(z)} dz$ first by parameterizing, changing to polar, and using the FTC, and second by using residues directly from the Laurent series.

7. The Main Story: The following are equivalent

- f is continuous
- f' exists
- f is analytic
- f is conformal
- \bullet f satisfies the Cauchy-Riemann equations

Theorems About Analytic Functions

1. If f is analytic on D, then $\oint_C f(z)dz = 0$ for any closed contour $C \subset D$.

Note: this does not require f to be f' to be continuous on C.

2. Maximum Modulus Principle

If f is analytic in a region D and not constant, then |f(z)| attains its maximum on ∂D .

- 3. If f is analytic, then $f^{(n)}$ is analytic for every n. If f = u(x,y) + iv(x,y), then all partials of u,v are continuous.
- 4. If f is analytic at z_0 and $f'(z_0) \neq 0$, then f is conformal at z_0 .
- 5. If f = u + iv is analytic, then u, v are harmonic conjugates.
- 6. If f is holomorphic, f is C_{∞} (smooth).

7. If f is analytic, f is holomorphic.

Proof: Since f has a power series expansion at z_0 , its derivative is given by the term-by-term differentiation of this series.

18.7.3 Some Useful Formulae

$$f_{x_0}(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{1}{2!}f''(x_0)(x - x_0)^2 + \dots$$

$$\frac{1}{1-z} = \sum_{k} z^k$$

$$e^z = \sum_k \frac{1}{k!} z^k$$

$$\left(\sum_{i} a_{i} z^{i}\right) \left(\sum_{j} b_{j} z^{j}\right) = \sum_{n} \left(\sum_{i+j=n} a_{i} b_{j}\right) z^{n}$$

$$\cos z = \frac{1}{2}(e^{iz} + e^{-iz}) \qquad = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \dots$$

$$\cosh z = \frac{1}{2}(e^z + e^{-z}) \qquad = \cos iz = 1 + \frac{z^2}{2!} + \frac{z^4}{4!} + \dots$$

$$\sin z = \frac{1}{2i}(e^{iz} - e^{-iz}) \qquad = z - \frac{z^3}{3!} + \frac{z^4}{4!} - \dots$$

$$\sinh z = \frac{1}{2}(e^z - e^{-z}) \qquad = -i\sin iz = z + \frac{z^3}{3!} + \frac{z^4}{4!} + \dots$$

Mnemonic: just remember that cosine is an even function, and that the even terms of e^z are kept. Similarly, sine is an odd function, so keep the odd terms of e^z .

Harmonic Conjugate

$$v(x,y) = \int_{(0,0)}^{(x,y)} -u_t(s,t)ds + u_s(s,t)dt$$

The Gamma Function

$$\Gamma(z) = \int_0^\infty x^{z-1} e^{-x} dx$$

137

Useful to know: $\Gamma(\frac{1}{2}) = \sqrt{\pi}$.

18.8 Questions



1. True or False: If f is analytic and bounded in \mathbb{H} , then f is constant on \mathbb{H} .

False: Take $f(z) = e^{-z}$, where $|f(z)| \le 1$ in \mathbb{H} .

2. Compute
$$\int_{-\infty}^{\infty} \frac{\sin x}{x(x^2 + a^2)} dx$$

Two semicircles needed to avoid singularity at zero. Limit equals the residue at zero, solution is $\pi(\frac{1}{a^2} - \frac{e^{-a}}{a^2})$.

3. Compute
$$\int_0^{2\pi} \frac{1}{2 + \cos \theta} d\theta$$

Cosine sub, solution is $\frac{2\pi}{\sqrt{3}}$

4. Find the first three terms of the Laurent expansion of $\frac{e^z+1}{e^z-1}$.

Equals
$$2z^{-1} + 0 + 6^{-1}z + \dots$$

5. Compute
$$\int_{S_1} \frac{1}{z^2 + z - 1} dz$$

Equals
$$i\frac{2\pi}{5}$$

6. True or false: If f is analytic on the unit disk $E = \{z : |z| < 1\}$, then there exists an $a \in E$ such that $|f(a)| \ge |f(0)|$.

True, by the maximum modulus principal. Suppose otherwise. Then f(0) is a maximum of f inside S_1 . But by the MMP, f must attain its maximum on ∂S_1 .

7. Prove that if f(z) and $f(\bar{z})$ are both analytic on a domain D, then f is constant on D

Analytic \implies Cauchy-Riemann equations are satisfied. Also have the identity $f' = u_x + iv_x$, and $f' = 0 \implies f$ is constant.

$oldsymbol{19}|$ Common Mistakes

$$-x^{-2} \neq \int x^{-1} = \int \frac{1}{x} = \ln x$$

$$\frac{1}{x} \neq \int \ln x = x \ln x - x$$

$$\int x^{-k} = \frac{1}{-k+1} x^{-k+1} \neq \frac{1}{-(k+1)} x^{-(k+1)}$$
e.g.
$$\int x^{-2} = -x^{-1} \neq -\frac{1}{3} x^{-3} \lim_{n \to \infty} \frac{n}{n+1} = 1 \neq 0$$

$$\frac{\partial}{\partial x} a^x = \frac{\partial}{\partial x} e^{x \ln a} = e^{x \ln a} \ln a = a^x \ln a.$$

Exponentials: when in doubt, write $a^b = e^{b \ln a}$

$$\frac{\partial}{\partial x}x^{f(x)} = ?$$

$$\sum x^k = \frac{1}{1-x} \neq \frac{1}{1+x} = \sum (-1)^k x^k$$

20 Appendix 1

20.1 Neat Tricks

• Commuting differentials and integrals:

$$\frac{d}{dx}\int_{a(x)}^{b(x)}f(x,t)dt=f(x,b(x))\frac{d}{dx}b(x)-f(x,a(x))\frac{d}{dx}a(x)+\int_{a(x)}^{b(x)}\frac{\partial}{\partial x}f(x,t)dt$$

- Need $f, df dx f, \frac{df}{dx}$ to be continuous in both variables. Also need $a(x), b(x) \in C1a(x), b(x) \in C1$.
- If a, b are constant, boundary terms vanish.
- Recover the fundamental theorem with a(x) = a, b(x) = ba(x) = a, b(x) = b, and f(x,t) = f(t)f(x,t) = f(t).

Common Mistakes 139

20

20.2 Big Derivative / Integral Table



20.1 Neat Tricks 140

20 Appendix 1

$\frac{\partial f}{\partial x} \Leftarrow$	f	$\Rightarrow \int f dx$	
$\frac{1}{2\sqrt{x}}$	\sqrt{x}	$rac{2}{3}x^{rac{3}{2}}$	
nx^{n-1}	$x^n, n \neq -1$	$\frac{1}{n+1}x^{n+1}$	
$-nx^{-(n+1)}$	$\frac{1}{x^n}, n \neq 1$	$-\frac{1}{n-1}x^{-(n-1)}$	
$\frac{1}{x}$	ln(x)	$x \ln(x) - x$	
$a^x \ln(a)$	a^x	$\frac{a^x}{\ln a}$	
$\cos(x)$	$\sin(x)$	$-\cos(x)$	
$-\csc(x)\cot(x)$	$\csc(x)$	$\ln \csc(x) - \cot(x) $	
$-\sin(x)$	$\cos(x)$	$\sin(x)$	
$\sec(x)\tan(x)$	sec(x)	$\ln \sec(x) + \tan(x) $	
$\sec^2(x)$	$\tan(x)$	$\ln \left \frac{1}{\cos x} \right $	
$-\csc^2(x)$	$\cot(x)$	$\ln \sin x $	
$\frac{\frac{1}{1+x^2}}{\frac{1}{\sqrt{1-x^2}}}$	$\tan^{-1}(x)$	$x\tan^{-1}x - \frac{1}{2}\ln(1+x^2)$	
$\frac{1}{\sqrt{1-x^2}}$	$\sin^{-1}(x)$	$x\sin^{-1}x + \sqrt{1-x^2}$	
$-\frac{1}{\sqrt{1-x^2}}$	$\cos^{-1}(x)$	$x\cos^{-1}x - \sqrt{1-x^2}$	
$\frac{1}{\sqrt{x^2 + a}}$	$\ln\left x+\sqrt{x^2+a}\right $		
$2\sin x\cos x$	$\sin^2(x)$	$\frac{1}{2}(x - \sin x \cos x)$	
$-2\sin x\cos x$	$\cos^2(x)$	$\frac{1}{2}(x+\sin x\cos x)$	
$2\csc^2(x)\cot(x)$	$\csc^2(x)$	$-\cot(x)$	
$2\sec^2(x)\tan(x)$	$\sec^2(x)$	$\tan(x)$	
?	?	?	
?	?	?	
?	?	?	
?	?	?	
?	?	?	
?	?	?	
?	?	?	
$(ax+1)e^{ax}$	xe^{ax}	$\frac{1}{a^2}(ax-1)e^{ax}$	
?	$e^{ax}\sin(bx)$	$\frac{1}{a^2 + b^2} e^{ax} (a\sin bx - b\cos bx)$	
20.2 Big Derivative / Integral Table	nm (-)	i de la companya de	141
?	$e^{ax}\cos(bx)$	$\frac{1}{a^2 + b^2}e^{ax}(a\sin bx + b\cos bx)$	
?	?	?	

20.3 Useful Series and Sequences

~

Notation: \uparrow , \downarrow : monotonically converges from below/above.

• Taylor Series:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$$

• Cauchy Product:

$$\left(\sum_{k=0}^{\infty}a_kx^k\right)\left(\sum_{k=0}^{\infty}b_ix^n\right)=\sum_{k=0}^{\infty}\left(\sum_{i=0}^ka_nb_n\right)x^k$$

• Differentiation:

$$\frac{\partial}{\partial x} \sum_{k=i}^{\infty} a_k x^k = \sum_{k=i+1}^{\infty} k \, a_k x^{k-1}$$

• Common Series

$$\begin{split} \sum_{k=0}^{N} x^k & = \frac{1-x^{N+1}}{1-x} \\ \sum_{k=1}^{\infty} x^k & = \frac{1}{1-x} \quad \text{for } |x| < 1 \\ \sum_{k=1}^{\infty} kx^{k-1} & = \frac{1}{(1-x)^2} \quad \text{for } |x| < 1 \\ \sum_{k=2}^{\infty} k(k-1)x^{k-2} & = \frac{2}{(1-x)^3} \quad \text{for } |x| < 1 \\ \sum_{k=3}^{\infty} k(k-1)(k-2)x^{k-3} & = \frac{6}{(1-x)^4} \quad \text{for } |x| < 1 \\ \sum_{k=1}^{\infty} \binom{n}{k} x^k y^{n-k} & = (x+y)^n \\ \sum_{k=1}^{\infty} \frac{x^k}{k!} & = -\log(1-x) \\ \sum_{k=0}^{\infty} \frac{x^k}{(2n+1)!} x^{2k+1} & = x - \frac{x^3}{3!} + \frac{x^5}{5!} & = \sin(x) \\ \sum_{k=0}^{\infty} \frac{(-1)^k}{(2n)!} x^{2k} & = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} & = \cos(x) \\ \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{2n+1} & = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \cdots & = \sinh(x) \\ \sum_{k=0}^{\infty} \frac{1}{(2k+1)!} x^{2k} & = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \cdots & = \cosh(x) \\ \sum_{k=0}^{\infty} \frac{1}{(2k)!} x^{2k} & = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \cdots & = \cosh(x) \\ \sum_{k=0}^{\infty} \frac{1}{(2k)!} x^{2k} & = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \cdots & = \cosh(x) \\ \sum_{k=0}^{\infty} \frac{1}{(2k)!} \frac{1}{k} & = \infty \\ \sum_{k=1}^{\infty} (-1)^k \frac{1}{k} & = \infty \\ \sum_{k=1}^{\infty} (-1)^k \frac{1}{k} & = \infty \\ \sum_{k=1}^{\infty} \frac{1}{k} & = \infty \\ \sum_{k=1}^{\infty} (-1)^k \frac{1}{k} & = \ln(2) \\ \sum_{k=1}^{\infty} \frac{1}{k} & = \frac{\pi^2}{6} \end{split}$$

20.4 Partial Fraction Decomposition

~

Given $R(x) = \frac{p(x)}{q(x)}$, factor q(x) into $\prod q_i(x)$.

• Linear factors of the form $q_i(x) = (ax + b)^n$ contribute

$$r_i(x) = \sum_{k=1}^n \frac{A_k}{(ax+b)^k} = \frac{A_1}{ax+b} + \frac{A_2}{(ax+b)^2} + \cdots$$

• Irreducible quadratics of the form $q_i(x) = (ax^2 + bx + c)^n$ contribute

$$r_i(x) = \sum_{k=1}^n \frac{A_k x + B_k}{(ax^2 + bx + c)^k} = \frac{A_1 x + B_1}{ax^2 + bx + c} + \frac{A_2 x + B_2}{(ax^2 + bx + c)^2} + \cdots$$

- Note: $ax^2 + bx + c$ is irreducible $\iff b^2 < 4ac$
- Write $R(x) = \frac{p(x)}{\prod q_i(x)} = \sum r_i(x)$, then solve for the unknown coefficients A_k, B_k .
 - IMPORTANT SHORTCUT: don't try to solve the resulting linear system: for each $q_i(x)$, multiply through by that factor and evaluate at its root to zero out many terms!
 - For linear terms $q_i(x) = (ax + b)^n$, define $P(x) = (ax + b)^n R(x)$; then

$$A_k = \frac{1}{(n-k)!} P^{(n-k)}(a), \quad k = 1, 2, \dots n$$

$$\implies A_n = P(a), \ A_{n-1} = P'(a), \ \dots, \ A_1 = \frac{1}{(n-1)!} P^{(n-1)}(A)$$

- Note: #todo check, might need to evaluate at -b/a instead, extend to quadratics.

20.5 Properties of Norms



$$\begin{aligned} &\|t\mathbf{x}\| = |t|\|\mathbf{x}\| \\ &|\langle \mathbf{x}, \ \mathbf{y} \rangle| \le \|\mathbf{x}\|\|\mathbf{y}\| \\ &\|\mathbf{x} + \mathbf{y}\| \le \|\mathbf{x}\| + \|\mathbf{y}\| \\ &\|\mathbf{x} - \mathbf{z}\| \le \|\mathbf{x} - \mathbf{y}\| + \|\mathbf{y} - \mathbf{z}\| \end{aligned}$$

20 Appendix 1

20.6 Logic Identities

- $P \implies Q \iff Q \text{ or } \neg P$
- $\bullet \ P \implies Q \iff \neg Q \implies \neg P$
- $P \text{ or } (Q \text{ and } S) \iff (P \text{ or } Q) \text{ and } (P \text{ or } S)$
- P and $(Q \text{ or } S) \iff (P \text{ and } Q) \text{ or } (P \text{ and } S)$
- $\neg (P \text{ and } Q) \iff \neg P \text{ or } \neg Q$
- $\neg (P \text{ or } Q) \iff \neg P \text{ and } \neg Q$

20.7 Set Identities

 $A \cup (A^c \cap B)$ $A \cup B$ A $(B \cap A) \cup (B^c \cap A)$ $(\cup_{\mathbb{N}}A_i)^c$ $\cap_{\mathbb{N}} A_i^c$ $(\cap_{\mathbb{N}} A_i)^c$ $\cup_{\mathbb{N}} A_i^c$ A - B $A \cap B^c$ $(A-B)^c$ $A^c \cup B$ $(A \cup B) - C$ $(A-C)\cup(B-C)$ $(A \cap B) - C$ $(A-C)\cap (B-C)$ $A - (B \cup C)$ $(A-B)\cap (A-C)$ $A - (B \cap C)$ $(A-B)\cup(A-C)$ A - (B - C) $(A-B)\cup(A\cap C)$ $(A-B)\cap C$ $(A \cap C) - B$ $A \cap (C - B)$ $(A-B)\cup C$ $(A \cup C) - (B - C)$ $A \cup (B \cap C)$ $(A \cup B) \cap (A \cup C)$ $A \cap (B \cup C)$ $(A \cap B) \cup (A \cap C)$ $A \subseteq C$ and $B \subseteq C$ $A \cup B \subseteq C$ $C \subseteq A$ and $C \subseteq B$ $C \subseteq A \cup B$ $\prod A_k, \ \cup_{k=1}^{\infty} A_k \quad \text{countable}$ A_k countable

20.8 Preimage Identities

Summary

• Injectivity: left cancellation

20.6 Logic Identities 145

- Surjectivity: right cancellation
- Everything commutes with unions
- Preimage commutes with everything
- Image generally only results in an inequality

Preimage Equations

•
$$A \subseteq B \Longrightarrow f(A) \subseteq f(B) \text{ or } f^{-1}(A) \subseteq f^{-1}(B)$$

• $f^{-1}(\bigcup_{i \in I} A_i) = \bigcup_{i \in I} f^{-1}(A_i)$

•
$$f^{-1}(\bigcup_{i \in I} A_i) = \bigcup_{i \in I} f^{-1}(A_i)$$

- Also holds for
$$f(\bigcup_{i \in I} A_i) = \bigcup_{i \in I} f(A_i)$$

•
$$f^{-1}(\cap_{i \in I} A_i) = \cap_{i \in I} f^{-1}(A_i)$$

- Also holds for
$$f(\cap_{i\in I} A_i) = \cap_{i\in I} f(A_i)$$

•
$$f^{-1}(A) - f^{-1}(B) = f^{-1}(A - B)$$

- BUT
$$f(A) - f(B) \subseteq f(A - B)$$

• For
$$X \subset A, Y \subset B$$
:

$$- (f|_X)^{-1} = X \cap f^{-1}(Y) - (f \circ f^{-1})(Y) = Y \cap f(A)$$

- Summary: preimage commutes with:
 - Union
 - Intersection
 - Complements
 - Difference
 - Symmetric Difference

Image Equations

•
$$A \subset B \implies f(A) \subset f(B)$$

- $f(\cup A_i) = \cup f(A_i)$
- $f(\cap A_i) \subset \cap f(A_i)$
- $f(A-B)\supset f(A)-f(B)$
- $f(A^c) = im(f) f(A)$

Equations Involving Both

•
$$A \subseteq f^{-1}(f(A))$$

- Equal
$$\iff f$$
 is injective

•
$$f(f^{-1}(A)) \subseteq A$$

- Equal
$$\iff$$
 f is surjective

20 Appendix 1





\overline{n}	Sequence
3	1, 2, 1
4	1, 3, 3, 1
5	1, 4, 6, 4, 1
6	1, 5, 10, 10, 5, 1
7	1, 6, 15, 20, 15, 16, 1
8	1, 7, 21, 35, 35, 21, 7, 1

Obtain new entries by adding in L pattern rotated by π (e.g. $7=1+6,\ 12=6+15,\ \text{etc}$). Note that $\binom{n}{i}$ is given by the entry in the n-th row, i-th column.

20.10 Table of Small Factorials



\overline{n}	n!
2	2
3	6
4	24
5	120
6	720
7	5040
8	40320
9	362880
10	3628800

 $\pi \approx 3.1415926535 \ e \approx 2.71828 \ \sqrt{2} \approx 1.4142135$

20

Appendix 1

20.11 Primes Under 100:



2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 83, 89, 97, 101

20.12 Checking Divisibility by Small Numbers



Note that $n \mod 10^k$ yields the last k digits. Let d_i denote the i-th digit of n.

The recursive prime procedure (RPP): for each prime p, there exists a k such recursive application of this procedure to n yields the same remainder mod p as n itself.

- Write $n_0 = 10x + y$ where y = 0...9
- Let $n_1 = x + ky$, repeat until $n_i < 10$.

p	$p \mid n \iff$	Mnemonic
2	$n \equiv 2, 4, 6, 8 \mod 10$	Last digit is even
3	$\sum d_i \equiv 0 \mod 3$	3 divides the sum of digits (apply recursively)
4	$n \equiv 4k \mod 10^2$	Last two digits are divisible by 4
5	$n \equiv 0, 5 \mod 10$	Last digit is 0 or 5
6	$n \equiv 0 \mod 2$ and $n \equiv 0 \mod 3$	Reduce to 2, 3 case
7	RPP, $k = -2$	$-20 \equiv 1 \mod 7 \Longrightarrow 10x + y \equiv 10(x - 2y) \mod 7$
8	$n \equiv 8k \mod 10^3$	Manually divide the last 3 digits by 8 (or peel off factors of 2)

20.11 Primes Under 100: 148

20 Appendix 1

\overline{p}	$p \mid n \iff$	Mnemonic
9	$\sum d_i \equiv 0 \mod 9$	9 divides the sum of digits (apply recursively)
10	$n \equiv 0 \mod 10$	Last digit is 0
11	$\sum (-1)^i d_i \equiv 0 \mod 11 \text{ or }$	11 divides alternating sum
13	$\overline{RPP}, k = 4$	$40 \equiv 1 \mod 13 \implies$
		$10x + y \equiv 10(x + 4y) \mod 13$
17	RPP, $k = -5$	$-50 \equiv 1 \mod 17 \implies$
		$10x + y \equiv 10(x - 5y) \mod 19$
19	RPP, $k=2$	$20 \equiv 1 \mod 19 \implies$
		$10x + y \equiv 10(x + 2y) \mod 19$

20.13 Hyperbolic Functions

$$\cosh(x) = \frac{1}{2}(e^x + e^{-x})$$

$$\sinh(x) = \frac{1}{2}(e^x - e^{-x})$$

$$\cos(iz) = \cosh z$$

$$\cosh(iz) = \cos z$$

$$\sin(iz) = \sinh z$$

$$\sinh(iz) = \sin z$$

$$\sinh^{-1} x = ? = \ln(x + \sqrt{x^2 + 1})$$

$$\cosh^{-1} x = ? = \ln(x + \sqrt{x^2 - 1})$$

$$\tanh^{-1} x = \frac{1}{2}\ln(\frac{1+x}{1-x})$$

20.14 Integral Tables

$\frac{\partial f}{\partial x} \Leftarrow$	f	$\Rightarrow \int f dx$
$\frac{1}{2\sqrt{x}}$	\sqrt{x}	$\frac{2}{3}x^{\frac{3}{2}}$
nx^{n-1}	$x^n, n \neq -1$	$\frac{1}{n+1}x^{n+1}$
$\frac{1}{x}$	$\ln(x)$	$x \ln(x) - x$
$a^x \ln(a)$	a^x	$\frac{a^x}{\ln a}$
$\cos(x)$	$\sin(x)$	$-\cos(x)$
$-\sin(x)$	$\cos(x)$	$\sin(x)$
$2\sec^2(x)\tan(x)$	$\sec^2(x)$	$\tan(x)$
$2\csc^2(x)\cot(x)$	$\csc^2(x)$	$-\cot(x)$
$\sec^2(x)$	$\tan(x)$	$\ln \sec(x) $
$\sec(x)\tan(x)$	sec(x)	$ \ln \sec(x) + \tan(x) $
$-\csc(x)\cot(x)$	$\csc(x)$	$\ln \csc(x) - \cot(x) $
$\frac{1}{1+x^2}$	$\tan^{-1}(x)$	$x \tan^{-1} x - \frac{1}{2} \ln(1 + x^2)$
$\frac{1}{1+x^2}$ $\frac{1}{\sqrt{1-x^2}}$	$\sin^{-1}(x)$	$x\sin^{-1}x + \sqrt{1-x^2}$
$-\frac{1}{\sqrt{1-x^2}}$	$\cos^{-1}(x)$	$x\cos^{-1}x - \sqrt{1-x^2}$
$\frac{1}{\sqrt{x^2 + a}}$	$\ln\left x + \sqrt{x^2 + a}\right $	
$-\csc^2(x)$	$\cot(x)$?
?	$\cos^2(x)$?
?	$\sin^2(x)$?
?	xe^{ax}	$\frac{1}{a^2}(ax-1)e^{ax}$
?	$e^{ax}\sin(bx)$	$\frac{1}{a^2 + b^2}e^{ax}(a\sin bx - b\cos bx)$
?	$e^{ax}\cos(bx)$	$\frac{1}{a^2 + b^2}e^{ax}(a\sin bx + b\cos bx)$
?	?	?

20.14 Integral Tables 150

21 Definitions

$$e^x = \lim_{n \to \infty} \left(1 + \frac{1}{n}\right)^n = \lim_{n \to \infty} \left(\frac{n+1}{n}\right)^n.$$

21.1 Set Theory

• Injectivity

$$f: X \to Y$$
 injective $\iff \forall x_1, x_2 \in X, \quad f(x_1) = f(x_2) \implies x_1 = x_2$
 $\iff \forall x_1, x_2 \in X, \quad x_1 \neq x_2 \implies f(x_1) \neq f(x_2).$

• Surjectivity

$$f: X \to Y \text{ surjective } \iff \forall y \in Y, \ \exists x \in X: f(x) = y.$$

• Preimage

$$f: X \to Y, U \subseteq Y \implies f^{-1}(U) = \{x \in X : f(x) \in U\}.$$

21.2 Calculus

• Limit

$$\lim_{x \to p} f(x) = L \iff \forall \varepsilon, \ \exists \delta :$$
$$d(x, p) < \delta \implies d(f(x), L) < \varepsilon$$

- Continuity
 - Epsilon-delta definition:

$$f: X \to Y$$
 continuous at $p \iff \forall \varepsilon, \ \exists \delta:$
 $d_X(x,p) < \delta \implies d_Y(f(x),f(p)) < \varepsilon$

- Limit/Sequential definition:

$$f: X \to Y$$
 continuous at $p \iff \forall \{x_i\}_{i \in \mathbb{N}} \subseteq X: \{x_i\} \to p$,

$$\lim_{i \to \infty} f(x_i) = f(\lim_{i \to \infty} x_i) = f(p)$$

Definitions 151

- Topological Definition:

 $f: X \to Y$ continuous $\iff U$ open in $\operatorname{im}(f) \subseteq Y \implies f^{-1}(U)$ open in X.

- Differentiability and the Derivative
 - For single variable functions:

$$f: \mathbb{R} \to \mathbb{R}$$
 differentiable at $p \iff \forall \{x_i\}_{i \in \mathbb{N}} \to p$,
$$f'(p) \coloneqq \lim_{i \to \infty} \frac{f(x_i) - f(p)}{x_i - p} < \infty$$

- For multivariable functions:

$$\mathbf{f}: \mathbb{R}^n \to \mathbb{R}^m$$
 differentiable at $\mathbf{p} \iff \exists$ a linear map $\mathbf{J}: \mathbb{R}^n \to \mathbb{R}^m$ such that:
$$\|\mathbf{f}(\mathbf{p} + \mathbf{h}) - \mathbf{f}(\mathbf{p}) - \mathbf{J}(\mathbf{h})\|_{\mathbb{R}^n}$$

$$\lim_{\mathbf{h}\to 0} \frac{\|\mathbf{f}\left(\mathbf{p}+\mathbf{h}\right) - \mathbf{f}\left(\mathbf{p}\right) - \mathbf{J}(\mathbf{h})\|_{\mathbb{R}^n}}{\|\mathbf{h}\|_{\mathbb{R}^m}} = 0$$

• Gradient

$$\nabla f = [f_x, f_y, f_z].$$

- Divergence
- Curl
- Taylor Series (at a point a)
 - Single Variable $\mathbb{R} \to \mathbb{R}$

$$T_a(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \cdots$$

$$\implies T_a(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n.$$

– Multivariable $\mathbb{R}^n \to \mathbb{R}$:

$$T_a(\mathbf{x}) = f(\mathbf{a}) + (\mathbf{x} - \mathbf{a})^T \nabla f(\mathbf{a}).$$

– Multivariable $\mathbb{R}^n \to \mathbb{R}^m$:

21.2 Calculus 152

$$T_{(a,b)}(x,y) = f(a,b) + (x-a)f_x(a,b) + (y-b)f_y(a,b) + \frac{1}{2!} \left((x-a)^2 f_{xx}(a,b) + 2(x-a)(y-b)f_{yy}(a,b) + (y-b)^2 f_{yx}(a,b) \right) + \cdots$$

$$T_a(\mathbf{x}) = f(\mathbf{a}) + (\mathbf{x} - \mathbf{a})^{\mathrm{T}} \mathbf{J}(\mathbf{a}) + \frac{1}{2!} (\mathbf{x} - \mathbf{a})^{\mathrm{T}} \mathbf{H}(\mathbf{a}) (\mathbf{x} - \mathbf{a}) + \cdots$$

$$\implies T_a(\mathbf{x}) = \sum_{|\alpha| \ge 0} \frac{(\mathbf{x} - \mathbf{a})^{\alpha}}{\alpha!} (\partial^{\alpha} f) (\mathbf{a})$$

21.3 Analysis

- Archimedean Property: $x \in \mathbb{R} \implies \exists n \in \mathbb{N}: \ x < n \text{ and } x > 0 \implies \exists n: \ \frac{1}{n} < x$
- Upper Bound (for S $\subseteq \mathbb{R}$) $\alpha \text{ is an upper bound for } S \iff s \in S \implies s < \alpha.$
- Triangle Inequality

$$- |a+b| \le |a| + |b| - |a-b| \le |a| + |b|$$

• Reverse Triangle Inequality

$$- ||a| - |b|| \le |a - b|$$

- Least Upper Bound / Supremum (for $S \subseteq \mathbb{R}$) $\alpha \text{ is a LUB for } S \iff s \in S \implies s < \alpha \text{ and } \forall t : (s \in S \implies s < t), \ \alpha < t.$
- Greatest Lower Bound / Infimum (for $S \subseteq \mathbb{R}$) $\alpha \text{ is a GLB for } S \iff s \in S \implies \alpha < s \text{ and } \forall t : (s \in S \implies t < s), \ t < \alpha.$
- Open Set
- Closed Set
- Limit Point

21.3 Analysis 153

- Interior Point
- Closure of a Set
- Boundary
- Metric
- Cauchy Sequence:

$$\{a_i\}$$
 is a cauchy sequence $\iff \forall \varepsilon \ \exists N \in \mathbb{N}: \ m,n > N \implies d(x_m,x_n) < \varepsilon.$

- Connected: S is connected $\iff \not\exists U, V \subset S$ nonempty, open, disjoint such that $S = U \cup V$
- Compact: Every open cover has a finite subcover:

$$X \subseteq \cup_{j \in J} V_j \implies \exists I \subseteq J : |I| < \infty \text{ and } X \subseteq \cup_{i \in I} V_i.$$

• Sequential Compactness Every sequence has a convergent subsequence:

$$\{x_i\}_{i\in I}\subseteq X\implies \exists J\subseteq I,\ \exists p\in X:\ \{x_j\}_{j\in J}\to p.$$

• Bounded (sequences, subsets, metric spaces)

$$U \subseteq X$$
 is bounded $\iff \exists x \in X, \exists M \in \mathbb{R}: u \in U \implies d(x, u) < M.$

• Totally Bounded

todo

• Pointwise Convergence

For
$$\{f_n : X \to Y\}_{n \in \mathbb{N}}$$
, $f_n \to f \iff \forall \varepsilon > 0, \ \forall x \in X, \ \exists N(x, \varepsilon) \in \mathbb{N} : n > N \implies d_Y(f_n(x), f(x)) < \varepsilon$

• Uniform Convergence

For
$$\{f_n: X \to Y\}_{n \in \mathbb{N}}$$
, $f_n \rightrightarrows f \iff \forall \varepsilon > 0, \ \exists N(\varepsilon) \in \mathbb{N}: \ \forall x \in X, \ n > N \implies d_Y(f_n(x), f(x)) < \varepsilon$

• Generalized Mean Value Theorem

$$(f(b) - f(a))g'(c) = (g(b) - g(a))f'(c).$$

21.3 Analysis 154

21.4 Linear Algebra

Convention: always over a field k, and $T: k^n \to k^m$ is a generic linear map (or $m \times n$ matrix).

• Consistent

A system of linear equations is *consistent* when it has at least one solution.

• Inconsistent

A system of linear equations is *inconsistent* when it has no solutions.

• Rank

The number of nonzero rows in RREF

- Elementary Matrix
- Row Equivalent
- Pivot
- Cofactor

$$\operatorname{cofactor}(A)_{i,j} = (-1)^{i+j} M_{i,j}$$

where $M_{i,j}$ is the minor obtained by deleting the *i*-th row and *j*-th column of A.

• Adjugate

$$\operatorname{adjugate}(A) = \operatorname{cofactor}(A)^T = (-1)^{i+j} M_{j,i}.$$

- Vector Space Axioms
 - Let k be a field and $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$ and $r, s, t \in k$. A vector space V over k satisfies:
 - 1. Closure under addition: $\mathbf{v} + \mathbf{w} \in V$
 - 2. Closure under scalar multiplication: $r\mathbf{v} \in V$
 - 3. Commutativity of addition: $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$
 - 4. Associativity of addition: $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
 - 5. Existence of an additive zero **0** satisfying $\mathbf{v} + 0 = 0 + \mathbf{v} = \mathbf{v}$
 - 6. Existence of additive inverse $-\mathbf{v}$ satisfying $v + (-\mathbf{v}) = 0$
 - 7. Unit property: $1\mathbf{v} = \mathbf{v}$
 - 8. Associativity of scalar multiplication: $(rs)\mathbf{v} = r(s\mathbf{v})$

- 9. Distribution of scalars multiplication over vector addition: $r(\mathbf{v} + \mathbf{w}) = r\mathbf{v} + r\mathbf{w}$
- 10. Distribution of scalar multiplication over scalar addition: $(r+s)\mathbf{v} = r\mathbf{v} + s\mathbf{v}$
- Subspace
 - A nonempty subset $W \subseteq V$ that is a vector space and satisfies

$$\left\{ \sum_{i} c_{i} \mathbf{x}_{i} \mid c_{i} \in \mathbb{F}, \ x_{i} \in W \right\} \subseteq W.$$

- Quick counter-check: find \mathbf{x}, \mathbf{y} such that $a\mathbf{x} + b\mathbf{y} \notin W$
- Span Given a set of n vectors $S = \{\mathbf{x}_i\}_{i=1}^n$, defined as

$$\operatorname{Span}(S) = \left\{ \sum_{i=1}^{n} c_i \mathbf{x}_i \mid c_i \in k \right\}.$$

- Row Space
 - The range of the linear map T.

- Given
$$T = \begin{bmatrix} \mathbf{x}_1 \to \\ \mathbf{x}_2 \to \\ \vdots \\ \mathbf{x}_m \to \end{bmatrix}$$
, defined as

$$\mathrm{Span}(\{\mathbf{x}_i\}_{i=1}^m) \subseteq k^m.$$

- $-\operatorname{rowspace}(T)^{\perp} = \operatorname{null}(T)$
- $|\operatorname{rowspace}(T)| = \operatorname{Rank}(T)$
- Column Space
- Null Space

– Defined as
$$\operatorname{null}(T) = \left\{ \mathbf{x} \in k^n \mid T(\mathbf{x}) = 0 \in k^m \right\}$$

– $\operatorname{null}(T)^{\perp} = \operatorname{rowspace}(T)$

- Eigenvalue
 - A value λ such that $Ax = \lambda x$
 - Invariant under similarity.

- Eigenspace
 - For a linear map T with eigenvalue λ , defined as $E_{\lambda} = \{ \mathbf{x} \in k^n \mid T(\mathbf{x}) = \lambda \mathbf{x} \}$
- Dimension
 - The cardinality of a basis of V
- Basis
 - A linearly independent set of vectors $S = \{\mathbf{x}_i\} \subset V$ such that $\mathrm{Span}(S) = V$
- Linear independence
 - A set of vectors $\{\mathbf{x}_i\}_{i=1}^n$ is linearly independent $\iff \sum_{i=1}^n c_i \mathbf{x}_i = 0 \implies c_i = 0$ for all i.
 - Can be detected by considering the matrix

$$T = [\mathbf{x}_1, \mathbf{x}_2, \cdots, \mathbf{x}_n]^T.$$

(linearly independent iff T is singular)

- Rank
 - Dimension of rowspace
- Rank-Nullity Theorem
 - |Nullspace(A)| + |Rank(A)| = |Codomain(A)|
- Nullspace
 - nullspace(A) = { $\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \mathbf{0}$ }
- Singular
 - A square $n \times n$ matrix T is singular iff Rank(T) < n
- Similarity
 - Two matrices A, B are similar iff there exists an invertible matrix S such that $B = SAS^{-1}$
- Diagonalizable

21.4 Linear Algebra 157

- A matrix X is diagonalizable if it can be written $X = EDE^{-1}$ where D is diagonal.
- If X is $n \times n$ and has n linearly independent eigenvectors λ_i , then $D_{ii} = \lambda_i$, and $E = \begin{bmatrix} \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \downarrow & \downarrow & \cdots & \downarrow \end{bmatrix}$
- Positive Definite
 - A matrix A is positive definite iff $\forall \mathbf{x} \in k^n$, we have the scalar inequality $\mathbf{x}^T A \mathbf{x} > 0$
- Projection
 - The projection of a vector \mathbf{v} onto \mathbf{u} is given by $P_{\mathbf{u}}(\mathbf{v}) = \langle \mathbf{u}, \mathbf{v} \rangle \, \hat{u}$
 - The projection of a vector \mathbf{v} onto a space $U = \mathrm{Span}(\{\mathbf{u}_i\})$ is given by

$$P_U(\mathbf{v}) = \sum_i P_{\mathbf{u}_i}(\mathbf{v}) = \sum_i \langle \mathbf{u}_i, \mathbf{v} \rangle \, \widehat{u}_i.$$

- Orthogonal Complement
 - Given a subspace $U \subseteq V$, defined as $U^{\perp} = \{ \mathbf{v} \in V \mid \forall \mathbf{u} \in U, \langle \mathbf{u}, \mathbf{v} \rangle = 0 \}$
- Determinant

$$\det(A) = \sum_{\tau \in S^n} \prod_{i=1}^n \sigma(\tau) a_{i,\tau(i)}.$$

• Trace

$$\operatorname{Tr}(A) = \sum_{i=1}^{n} A_{ii}.$$

- Characteristic Polynomial
 - $-p_A(x) = \det(xI A)$
 - Roots of p_A are eigenvalues of A
- Symmetric: $A = A^T$
- Skew-Symmetric: $A = -A^T$
- Inner Product

$$\begin{aligned}
& - \langle \mathbf{x}, \ \mathbf{x} \rangle \ge 0 \\
& - \langle \mathbf{x}, \ \mathbf{x} \rangle = 0 \iff \mathbf{x} = \mathbf{0} \\
& - \langle \mathbf{x}, \ \mathbf{y} \rangle = \overline{\langle \mathbf{y}, \ \mathbf{x} \rangle}
\end{aligned}$$

21.4 Linear Algebra 158


$$-\langle [, k\rangle \mathbf{x}]\mathbf{y} = k\langle \mathbf{x}, \mathbf{y}\rangle = \langle \mathbf{x}, k\mathbf{y}\rangle
-\langle \mathbf{x} + \mathbf{y}, \mathbf{z}\rangle = \langle \mathbf{x}, \mathbf{z}\rangle + \langle [, y\rangle]\mathbf{z}
-\langle [, a\rangle \mathbf{x}]b\mathbf{y} = \langle \mathbf{x}, \mathbf{x}\rangle + \langle a\mathbf{x}, y\rangle + \langle \mathbf{x}, b\mathbf{y}\rangle + \langle \mathbf{y}, \mathbf{y}\rangle
- Defines a norm: $\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x}\rangle} \implies \|\mathbf{x}\|^2 = \langle \mathbf{x}, \mathbf{x}\rangle$$$

- Cauchy-Schwarz Inequality: $|\langle \mathbf{x}, \mathbf{y} \rangle| \le ||\mathbf{x}|| ||\mathbf{y}||$
- Orthogonality:
 - For vectors: $\mathbf{x}^{\perp}\mathbf{y} \iff \langle \mathbf{x}, \ \mathbf{y} \rangle = 0$ - For matrices: A is orthogonal $\iff A^{-1} = A^T$
- Orthogonal Projection of \mathbf{x} onto \mathbf{y} :

$$P(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, \mathbf{y} \rangle \widehat{y} = \langle \mathbf{x}, \mathbf{y} \rangle \frac{\mathbf{y}}{\|\mathbf{y}\|^2}.$$

- Note $||P(\mathbf{x}, \mathbf{y})|| = ||\mathbf{x}|| \cos \theta_{x,y}$
- Defective: An $n \times n$ matrix A is defective \iff the number of linearly independent eigenvectors of A is less than n.

21.5 Differential Equations



$$f(x,y)$$
 homogeneous of degree $n \iff \exists n \in \mathbb{N} : f(tx,ty) = t^n f(x,y)...$

• Separable

$$p(y)\frac{dy}{dx} - q(x) = 0.$$

• Wronskian:

$$W[f_1, f_2, \dots, f_k](x) = \begin{vmatrix} f_1(x) & f_2(x) & \dots & f_k(x) \\ f'_1(x) & f'_2(x) & \dots & f'_k(x) \\ \vdots & \vdots & & \vdots \\ f_1^{(k-1)}(x) & f_2^{(k-1)}(x) & \dots & f_k^{(k-1)}(x) \end{vmatrix}$$

• Laplace Transform:

$$L_f(s) = \int_0^\infty e^{-st} f(t) dt.$$

21.6 Algebra



- Ring
- Group
- Subgroup
 - Two step subgroups test:
- Integral Domain
- Division Ring
- Principal Ideal Domain
- Tensor Product: #todo insert construction


21.7 Complex Analysis



- Analytic
- Harmonic
- Cauchy-Euler Equations
- Holomorphic
- The Complex Derivative
- Meromorphic
- The Gamma Function: Satisfies $\Gamma(p+1) + p\Gamma(p)$ and $\Gamma(1) = 1$, defined as

$$\Gamma(p) = \int_0^\infty t^{p-1} e^{-t} dt, \quad p > 0.$$

21.8 Algebra



- Let p be number of sign changes in f(x);
- Let q be number of sign changes in f(-x);
- Let n be the degree of f.
- Then p gives the maximum number of positive real roots, q gives the maximum number of negative real roots, and n p q gives the *minimum* number of complex roots.
- Rational Roots Theorem: If $p(x) = ax^n + \cdots + c$ and $r = \frac{p}{q}$ where p(r) = 0, then $p \mid c$ and $q \mid a$.
- Properties of logs:

$$-\ln(\prod) = \sum \ln \operatorname{but} \prod \ln \neq \ln \sum$$

21.6 Algebra 160

$$-\log_b x = \frac{\ln x}{\ln b}$$

Be careful! $\frac{\ln x}{\ln y} \neq \ln \frac{x}{y} = \ln x - \ln y$

• Completing the square:

$$-p(x) = ax^{2} + bx + c \implies p(x) = a(x + \frac{b}{2a})^{2} + -\frac{1}{2}\left(\frac{b^{2} - 4ac}{2a}\right)$$

21.9 Geometry

• Generic Conic Sections

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$

$$\frac{(x-x_0)^2}{w_0} \pm \frac{(y-y_0)^2}{h_0} = c$$

• Circles:

$$Ax^{2} + By^{2} + C = 0$$
 $(x - x_{0})^{2} + (y - y_{0})^{2} = r^{2}$

- Defining trait: locus of points at a constant distance from the **center**
- Center at (x_0, y_0)
- Parabolas:

$$Ax^2 + Bx + Cy + D = 0 y = ax^2$$

- Defining Trait:
 - ♦ Locus of points equidistant from the **focus** (a point) and the **directrix** (a line)
 - ♦ #todo add image
- **Focus** at $(0, \frac{1}{4a})$
- **Directrix** at the line $y = -\frac{1}{4a}$
 - \diamondsuit For an arbitrary quadratic: complete the square to write in the form $y = a(x w_0)^2 + h_0$, and translate points of interest by by $(x + w_0, y + h_0)$

• Ellipses:

$$\frac{x^2}{w^2} + \frac{y^2}{h^2} = 1$$

- Defining trait:
 - \Diamond The locus of points where the *sum* of distances to two **focii** are constant.
- Center at (0,0) (can translate easily)
- **Vertices** at $(\pm w, 0)$ and $(0, \pm h)$
- Focii at $F_1 = (\sqrt{w^2 h^2}, 0), F_2 = (-\sqrt{w^2 h^2}, 0)$
- Another useful shortcut form:
- Hyperbolas:

$$\frac{x^2}{w^2} - \frac{y^2}{h^2} = 1$$

- Defining trait:
 - \Diamond Locus of points where the difference between the distances to two **focii** are constant.
- Vertices at $(0, \pm h)$ and $(\pm w, 0)$
- Focii at $F_1 = (\sqrt{w^2 + h^2}, 0), F_2 = (-\sqrt{w^2 + h^2}, 0)$
- Summary of Traits:
 - One point p:
 - \Diamond Distance to p is constant: circle
 - Two points a, b:
 - \Diamond Distance to a equal to distance to b equals a constant: a line bisecting the midpoint of the line connecting them
 - ♦ Difference of distances constant: ellipse
 - \Diamond Sum of differences constant: hyperbola
 - Point p and a line l:
 - \Diamond Distance to p equals distance to l equals a constant: parabola
- Areas of certain figures:

21.9 Geometry 162

22 | Indices

List of Todos

Find examples	8
Todo	8
todo	8
Examples	10
For constants, this should allow differentiating under the integral when f, f_x are "jointly continuous"	15
Completing the square	16
Trig functions, double angle formulas.	16
Need lots of pictures	18
Distance from origin to plane	22
Distance from point to plane	22
todo	24
Add examples	29
Add examples	29
Need a better proof, not clear that this works	33
todo	33
todo	38
todo	38
Show how to compute basis of kernel	39
Show how to compute basis of row space (nonzero rows in $??A$)?)	39
Show how to compute basis of column space: leading ones	39

Indices 163

Proof of Cauchy-Schwarz: See Goode page 346	41
todo	44
Add examples	50
todo	51
todo	52
Check! Domain or range?	57
Derivation of normal equations	59
Explain shortcut for diagonal	61
todo	68
todo	68
Sufficient condition: each f_i is the solution to a linear homogeneous ODE $L(y)=0$	70
This section is very sketchy!	72
Ring Axioms	77
When is $\mathbb{Q}(\sqrt{d})$ a field?	78
Link to theorems	79
Check	79
Check	80
Example of Euclidean algorithm,	80
How to obtain solutions	81
todo	84
todo	84
todo	88
todo	93
todo	03

todo		97
todo		98
todo		.01
todo		.05
todo		.05
todo		.05
Bring in R	udin's list	.08
???		.11
todo		.54
Defin	itions	
4.2.1	Definition – Partial Derivative	
4.3.1	Definition – General definition of differentiability	
4.3.2	Definition – Jacobian	
4.3.3	Definition – Hessian	
4.6.1	Definition – Critical Points	
5.2.1	Definition – Directional Derivative	
6.4.1	Definition – Linear Transformation	
6.4.2	Definition – Kernel	
6.4.3	Definition – Basis and dimension	
6.4.4	Definition – The standard inner product	
6.4.5 $6.5.1$	7 1	40
6.5.1		44 44
6.5.2		44
6.5.4	•	45
6.5.5		45
6.5.6		48
6.5.7	Definition – Cofactors	48
6.5.8		51
6.5.9		51
9.7.1	9 , 9 ,	70
11.1.1	Definition – Multiplicative Functions	79
11.4.1	1	79
11.6.1	· ·	83
11.7.1	Definition – Quadratic Residue	84

22 Theorems

11.7.2	Definition – The Jacobi Symbols		 	•	•		•	84
11.7.3	Definition – The Legendre Symbol		 					84
12.2.1	Definition – Convergence of a Sequence		 					86
13.0.1	Definition – Series		 					87
13.2.1	Definition – Absolutely Convergent		 					88
14.1.1	Definition – Continuously Differentiable		 					92
14.6.1	Definition – Limit definition of continuity		 					97
14.6.2	Definition $-\varepsilon$ - δ definition of continuity		 					97
14.6.3	Definition – Uniform Continuity		 					98
14.6.4	Definition – Absolute Continuity		 					98
14.11.1	Definition – Convergence of an infinite series		 					101
14.11.2	Definition – Cauchy Sequence		 					105
	Definition – Metric							
	Definition – Complete							
	Definition – Bounded							
	Definition – Axioms for a Topology							
	Definition – Sequential Compactness							
16.1.1	Definition – Random Variable							
16.1.2	Definition – Probability Density Function (PDF)							
16.1.3	Definition – Cumulative Distribution Function (CDF)							
16.1.4	Definition – Mean/Expected Value							
16.1.5	Definition – Variance							
16.1.6	Definition – Covariance							
16.1.7	Definition – Entropy							
16.2.1	Definition – Axioms for a Probability Space							
16.2.2	Definition – Conditional Probability							
16.2.3	Definition – Odds							
16.2.4	Definition – Independence							
16.6.1	Definition – The Gamma Function							
16.6.2	Definition – Moment Generating Functions							
heo	rems							
1.1.1	$\label{eq:continuous} Proposition-Fundamental\ Theorem\ of\ Calculus\ I\ .\ .\ .\ .\ .$		 					7
1.1.2	Proposition – Generalized Fundamental Theorem of Calculus		 					7
1.1.3	Proposition – Extreme Value Theorem		 					8
1.1.4	Proposition – Mean Value Theorem		 					8
1.1.5	Proposition – Rolle's Theorem		 					8
1.1.6	Proposition – L'Hopital's Rule		 					9
1.1.7	Proposition – Taylor Expansion							9
1.6.1	Proposition – Chain Rule							11
1.6.2	Proposition – Product Rule							11
1.6.3	Proposition – Quotient Rule							11
1.6.4	Proposition – Inverse Rule							12
2.1.1	Proposition – Integral formula for average value				_			13

Theorems 166

22 Theorems

3.1.2	Proposition – Rotation matrices in \mathbb{R}^2	19
	T	
3.1.3	Proposition	19
3.3.1	Proposition – Equation for a line between two points	20
3.3.2	Proposition – Symmetric equation of a line	20
3.3.1	Theorem – The Tangent Plane Equation	21
3.4.1	Proposition – Writing equation from a point and a normal	21
3.4.2	Proposition – Writing equation from two vectors	21
4.0.1	Theorem – Key Theorem	25
4.6.1	Proposition – Second Derivative Test	28
5.2.1	Theorem – Stokes' Theorem	32
5.2.2	Theorem – Dot product expression of directional derivative	32
5.3.1	Proposition – Areas can be computed with Green's Theorem	35
5.4.1	Proposition – Sufficient Conditions	36
6.2.1	Theorem – Rank-Nullity	37
6.4.1	Proposition – Two-step vector subspace test	38
6.4.2	Proposition –?	39
6.4.3	Proposition – Inner products induce norms and angles	39
6.4.4	Proposition – Formula expanding a norm and 'Pythagorean theorem'	40
6.4.5	Proposition – Properties of the inner product	40
6.5.1	Proposition –?	44
6.5.2	Proposition – Formula for matrix multiplication	45
6.5.3	Proposition – Characterizing solutions to a system of linear equations	45
6.5.4	Proposition –?	46
6.5.5	Proposition – Inverse of a 2×2 matrix	46
6.5.6	Proposition – Properties of the determinant	46
6.5.7	Proposition – Characterizing singular matrices	48
6.5.8	Proposition – Laplace/Cofactor Expansion	48
6.5.9	Proposition – Computing determinant from RREF	49
6.5.10	Proposition – Cramer's Rule	49
6.5.11	Proposition – Gauss-Jordan Method for inverting a matrix	49
6.5.12	Proposition – Cofactor formula for inverse	49
6.5.13	Proposition	
6.5.14	Proposition – How to find eigenvectors	52
7.1.1	Proposition – Changing to the standard basis	57
7.3.1	Proposition – Projection onto range	57
7.3.2	Proposition – Normal Equations	59
9.6.1	Theorem – First Shifting Theorem	70
9.7.1	Proposition – Wronskian detects linear dependence of functions	70
11.3.1	Theorem – The fundamental theorem of arithmetic $\ldots \ldots \ldots$	79
11.4.1	Proposition – Relationship between gcd and lcm	80
11.4.2	Proposition –?	80
11.4.3	Proposition – Useful properties of gcd	80
11.5.1	Theorem – The Chinese Remainder Theorem	80
11.5.2	theorem-Euler's Theorem	81
11.5.3	Theorem – Fermat's Little Theorem	81
11.5.1	Proposition – Solutions to linear Diophantine equations	81
11.5.2	Proposition –?	81

Theorems 167

Theorems

11.5.3	Proposition – Repeated square/last exponentiation	82
11.5.4	Proposition – Reduction of modulus	82
11.5.5	Proposition – Characterization of invertibility	82
11.6.1	Proposition – Formulas involving the totient	83
11.7.1	Proposition –?	84
11.7.2	Proposition –?	
11.8.1	Proposition – Fermat Primality Test	85
11.8.2	Proposition – Miller-Rabin Primality Test	
11.9.1	Theorem – Bolzano-Weierstrass	
11.9.2	Theorem – Heine-Borel	
11.9.1	Proposition – Converse of Heine-Borel	
12.2.1	Theorem – Squeeze Theorem	
12.2.2	Theorem – Monotone Convergence Theorem for Sequences	
12.2.3	Theorem – Cauchy Criteria	
13.2.1	Proposition –?	
13.2.2	Proposition – The Cauchy Criterion	
13.2.1	Theorem – Comparison Test	
13.2.2	Theorem – Ratio Test	
13.2.3	Theorem – Root Test	
13.2.4	Theorem – Integral Test	
13.2.5	Theorem – Limit Test	
13.2.6	Theorem – Alternating Series Test	
13.2.7	Theorem – Weierstrass M -Test	
13.3.1	Proposition – Finding the radius of convergence	91
14.2.1	Theorem – Generalized Mean Value Theore	
14.6.1	Proposition – Thomae's function: the set of points of continuity and of disconti-	
	nuity can both be infinite	97
14.6.1	Theorem – Extreme Value Theorem	
14.9.1	Theorem – FTC for the Riemann Integral	
14.9.1	Proposition	
	Proposition – ?	
	Proposition –?	
	Proposition – ?	
	Proposition – A pointwise limit of continuous functions is not necessarily continuous.	
	Proposition – The limit of derivatives need not equal the derivative of the limit .	
	Proposition –?	
	Proposition – All subsequences of a convergent sequence share a limit	
	Proposition –?	
	Proposition – ?	
	Proposition – ?	
	Proposition –?	
	Proposition –?	
	Proposition –?	
	Proposition – Compactness and sequential compactness	
	Proposition – ?	
	Proposition – ?	
	Proposition – ?	
	Proposition – Linearity of Expectation	
	_ · · · · · · · · · · · · · · · · · · ·	

Theorems 168

16.1.2	Proposition – Properties of Variance
16.1.3	Proposition – Properties of Covariance
16.1.4	Proposition – Stirling's Approximation
16.1.5	Proposition – Markov Inequality
16.1.6	Proposition – Chebyshev's Inequality
16.1.1	Theorem – Central Limit Theorem
16.1.2	Theorem – Strong Large of Large Numbers
16.1.7	Proposition – Chernoff Bounds
16.1.8	Proposition – Jensen's Inequality
16.2.1	Proposition – Properties that follow from axioms
16.2.1	Theorem – Bayes' Rule
16.2.2	Proposition – Change of Variables for PDFs
16.2.3	Proposition – PDF for a sum of independent random variables
16.6.1	Proposition – Boole's Inequality

Exercises

List of Figures

1	Image	43
2	image	130
3	image	131
4	image	131
5	$z\mapsto \log z$	132
	$z \leftarrow \sin w$	
7	$z \mapsto z + z^{-1} \dots \dots$	133
8	image	133
	image	
10	image	135
11	image	135

Exercises 169