

# Field arithmetic and the complexity of algebraic objects

Danny Krashen, IAS/PCMI GSS 2021

*D. Zack Garza*  
*University of Georgia*  
[dzackgarza@gmail.com](mailto:dzackgarza@gmail.com)

*Last updated: 2021-07-15*

# Table of Contents

## Contents

<b>Table of Contents</b>	<b>2</b>
<b>1 Monday, July 12</b>	<b>3</b>
1.1 Intro . . . . .	3
1.2 Galois Cohomology . . . . .	3
1.3 Milnor K-Theory . . . . .	4
1.4 Witt Ring . . . . .	4
1.5 Motivic Cohomology . . . . .	5
1.6 Dimension . . . . .	6
1.7 Structural Problems in Galois Cohomology . . . . .	7
1.7.1 Period-Index Problems . . . . .	8
1.7.2 Symbol Length Problem . . . . .	8
<b>2 Tuesday, July 13</b>	<b>9</b>
2.1 Pfister Form . . . . .	11
2.2 Canonical Dimension . . . . .	12

# 1 | Monday, July 12

Talk: Danny Krashen

## 1.1 Intro

Missed first 13m

Fix a field  $k_0 \in \text{Field}$ , we'll consider extensions  $k \in \text{Field}/_{k_0}$ .

## 1.2 Galois Cohomology

**Definition 1.2.1** (Galois Cohomology)

For  $M \in \mathbf{G}_k\text{-Mod}$  for  $G_k$  the Galois group of  $k \in \text{Field}/_{k_0}$ , we can take invariants  $M^{G_k}$ . The functor  $-^{G_k}$  is left-exact, so we define

$$H_{\text{Gal}}^*(G_k; -) := \mathbb{R}^*(-)^{G_k}.$$

**Remark 1.2.2:** Note that the tensor product on  $\mathbf{G}_k\text{-Mod}$  induces a cup product on  $H_{\text{Gal}}^*$ . An important example of coefficients is  $M = \mu_\ell^{\otimes m}$ , where  $\mu_\ell^{\otimes 0} := \mathbb{Z}/n$ . It is known that  $H_{\text{Gal}}^*(G_k; \mu^{\otimes 0}) = \mathbb{Z}/n$ .

We'll define *symbols*

$$(a_1, \dots, a_n) := (a_1) \smile \dots \smile (a_n) \in H_{\text{Gal}}^*(k, \mu_\ell^{\otimes n}),$$

which are in fact generators. To remember the  $\ell$ , we write  $(a_1, a_2, \dots, a_n)_\ell$ .

**Remark 1.2.3:** Galois cohomology is a special case of étale cohomology, where for  $M \in \mathbf{G}_k\text{-Mod}$ ,

$$H_{\text{Gal}}^n(G_k; M) = H_{\text{ét}}^n(k; M) = H_{\text{ét}}^n(\text{Spec } k; M).$$

Étale cohomology works for schemes other than just  $\text{Spec } k$ .

## 1.3 Milnor K-Theory

**Definition 1.3.1** (?)

Given  $k \in \text{Field}$ , define

$$K_*^M(k) := \bigoplus_{i=1}^{\infty} K_i^M(k)$$

where

- $K_0^M(k) = \mathbb{Z}$
- $K_1^M(k) = k^m$ , written additively as elements  $\{a\}$  on the left-hand side, so  $\{a\} + \{b\} := \{ab\}$ .
- It's generated by  $K_1^M(k)$ , with products written by concatenation:

$$\{a_1, \dots, a_n\} = \{a_1\} \{a_2\} \cdots \{a_n\}.$$

- The only relations are  $\{a, b\} = 0$  when  $a + b = 1$ , motivated by

$$(a, b)_\ell = 0 \in H_{\text{Gal}}^2(k; \mu_\ell^{\otimes 2}) \iff a + b = 1.$$

- There is a map

$$\begin{aligned} K_0^M(k) &\rightarrow H_{\text{ét}}^*(k; \mu_\ell^{\otimes 0}) \\ \{a\} &\mapsto (a), \end{aligned}$$

and the **Norm-Residue isomorphism** (formerly the **Bloch-Kato conjecture**) states that this is an isomorphism after modding out by  $\ell$ , i.e.

$$K_0^M(k)/\ell \xrightarrow{\sim} H_{\text{ét}}^*(k; \mu_\ell^{\otimes 0}).$$

## 1.4 Witt Ring

**Remark 1.4.1:** Assume  $\text{ch } k \neq 2$ , so there is a correspondence between quadratic forms and symmetric bilinear forms given by polarization:

Quadratic forms  $\iff$  Symmetric bilinear forms

$$q_b(x) := b(x, x) \leftrightarrow b(x, y)$$

$$q \mapsto b_q(x, y) := \frac{1}{2} (q(x + y) - q(x) - q(y)).$$

So we'll identify these going forward and write  $q$  for an arbitrary symmetric bilinear form or a quadratic form. We say  $q$  is **nondegenerate** if there is an induced isomorphism:

$$\begin{aligned} V &\xrightarrow{\sim} V^\vee \\ v &\mapsto b_q(v, -). \end{aligned}$$

Note that a symmetric bilinear form  $q$  on  $V$  can be regarded as an element of  $\text{Sym}^2(V^\vee)$ .

**Definition 1.4.2** (The Witt Ring)

Let  $\text{QuadForm}/_k$  be the category of pairs  $(V, q)$  with  $V \in \text{Vect}/_k$  a  $k$ -vector space and  $q \in \text{Sym}^2(V^\vee)$  representing a quadratic form on  $V$ . The **Witt ring** is generated as a group by isomorphism representing a quadratic form on  $V$ .

$$W(k) = \frac{\mathbb{Z} \langle \{ [(V, q)] \in \text{QuadForm}/_k \} \rangle}{\langle q_{\text{hyp}}, (q_1 + q_2) - (q_1 \perp q_2) \rangle} \in \text{AbGrp.}$$

where the **hyperbolic form** is defined as  $q_{\text{hyp}}(x, y) = xy$ . The ring structure is given by the tensor product (a.k.a. Kronecker product of forms).

**Remark 1.4.3:** Noting that Galois cohomology lives mod  $\ell$  for various  $\ell$ , here  $K_0^M(k)$  lives over  $\mathbb{Z}$ . So Milnor K-theory relates all of the various mod  $\ell$  Galois cohomologies together.

**Definition 1.4.4** (Fundamental ideals and Pfister Forms)

The **fundamental ideal**  $I(k) \trianglelefteq W(k)$  is the ideal of even dimensional forms, and set  $I^n(k) := (I(k))^n$ . There is a map

$$\begin{aligned} K_n^M(k) &\rightarrow I^n(k)/I^{n+1}(k) \\ \{a_1, a_2, \dots, a_n\} &\mapsto \langle \langle a_1, a_2, \dots, a_n \rangle \rangle, \end{aligned}$$

which follows from Gram-Schmidt: any form can be diagonalized  $q \cong \sum a_i x_i^2$ , which we can write as  $\langle a_1, a_2, \dots, a_n \rangle$ . We can define the  **$n$ -fold Pfister forms**

$$\begin{aligned} \langle \langle a \rangle \rangle &:= \langle \langle 1, -a \rangle \rangle \\ \langle \langle a_1, a_2, \dots, a_n \rangle \rangle &:= \prod_{i=1}^n \langle \langle a_i \rangle \rangle. \end{aligned}$$

**Remark 1.4.5:** The **Milnor conjecture** (proved by Voevodsky et al) states that the above map is an isomorphism after modding out by 2, so

$$K_n^M(k)/2 \xrightarrow{\sim} I^n(k)/I^{n+1}(k).$$

Moreover, the LHS is isomorphic to  $H^n(k, \mu_2)$ . There are interesting maps going the other way

$$I^n(k) \rightarrow I^n(k)/I^{n+1}(k) \xrightarrow{\sim} H^n(k, \mu_2)$$

Upshot: this is surjective – any mod 2 cohomology class comes from a quadratic form, and thus we can reason about cohomology by reasoning about quadratic forms.

## 1.5 Motivic Cohomology

**Remark 1.5.1: Motivic cohomology** relates the various mod  $\ell$  cohomologies together much like  $K_*^M$ , but additionally relates different twists. In particular, it relates various  $H_{\text{ét}}^i(k; \mu_\ell^{\otimes j})$ , where Milnor K-theory interprets this “diagonally” when  $i = j$ . This works by constructing **motivic complexes**

$$\mathbb{Z}(m) \in \text{Ch}_{\text{pre}}(\text{ShsmSch}/k),$$

which are complexes of presheaves on smooth  $k$ -schemes, usually considered in the Zariski, étale, or Nisnevich topologies.

**Remark 1.5.2: Zariski hypercohomology** is defined as

$$\mathbb{H}^n(X; \mathbb{Z}(m)) = H^{n,m}(X; \mathbb{Z}) = H_{\text{mot}}^n(X; \mathbb{Z}(m)) \quad \text{for } X := \text{Spec } k.$$

These relate to Galois cohomology in the following ways:

- There is a quasi-isomorphism  $\mu_\ell^{\otimes m} \xrightarrow{\sim} \mathbb{Z}/\ell(n)$  in the étale topology.
- There is an isomorphism  $H_{\text{zar}}^n(k, \mathbb{Z}(n)) \xrightarrow{\sim} K_n^M(k)$ .
- Bloch-Kato identifies  $H_{\text{zar}}^*(X; \mathbb{Z}/\ell(n)) \xrightarrow{\sim} H_{\text{ét}}^n(X; \mathbb{Z}/\ell(n))$ .

## 1.6 Dimension

**Remark 1.6.1:** There are a number of competing notions for the “dimension” of a field.

**Definition 1.6.2** (Dimension of a field)

If  $k$  is finitely generated over either a prime field or an algebraically closed field, we say

$$\dim(k) = \begin{cases} [k : k_0]_{\text{tr}} & k_0 = \bar{k}_0 \\ [k : k_0]_{\text{tr}} + 1 & k_0 \text{ finite} \\ [k : k_0]_{\text{tr}} + 2 & k_0 = \mathbb{Q}. \end{cases}$$

**Definition 1.6.3** (Cohomological dimension)

We define its **cohomological dimension**  $\text{cohdim}(k)$ , which is at most  $n$  if  $H^m(G_k; M) = 0$  for all  $m > n$  and  $M$  torsion,

$$\text{cohdim}(k) := \min \left\{ n \mid \text{cohdim}(k) \leq n \right\}.$$

Equivalently,  $\text{cohdim}(k) = n \iff$  there exists a torsion  $M$  with  $H^n(G_k; M) \neq 0$  and  $H^m(G_k; M) = 0$  for all  $m > n$ .

**Remark 1.6.4:**  $\text{cohdim}(k) = \dim(k)$  if  $k$  is finitely generated or a finite extension of  $k_0 = \bar{k}_0$ , or if  $k$  is finitely generated over  $\mathbb{Q}$  and has no real orderings. So if  $k$  has orderings,  $\text{cohdim}(k) = \infty$ .

**Definition 1.6.5** (Diophantine Dimension)

We say  $k$  is  $C_n$  if for  $d > 0$  and  $m > d^n$ , then every homogeneous polynomial of degree  $d$  in  $m$  variables has a nontrivial root.

$$\text{ddim}(k) := \min \{n \mid k \text{ is } C_n\}.$$

**Example 1.6.6(?)**: If  $k$  is finitely generated or finite over  $k_0 = \bar{k}_0$ , then

$$\text{ddim}(k) = \dim(k) = \text{cohdim}(k).$$

**Definition 1.6.7** ( $T_n$ -rank)

We say  $k$  is  $T_n$  if for every  $d_1, d_2, \dots, d_r > 0$  and every system of polynomial equations  $f_1 = \dots = f_r = 0$  with  $\deg f_i = d_i$  in  $m$  variables, with  $m > \sum d_i^n$ . Then the  $T_n$ -rank is defined as

$$T_n\text{-rank}(k) := \min \{n \mid k \text{ is } T_n\}.$$

**Question 1.6.8**

Note that  $T_n \implies C_n$ , so  $T_n\text{-rank}(k) \geq \text{ddim}(k)$ , when are they equal? This is likely unknown.

**Remark 1.6.9**: There is a famous example of a field  $k$  with  $\text{cohdim}(k) = 1$  but  $\text{ddim}(k) = \infty$ .

**Question 1.6.10**

Is it true that  $\text{ddim}(k) \geq \text{cohdim}(k)$ ? Serre showed that this holds when  $\text{cohdim}$  is replaced by  $\text{cohdim}_2$ , the 2-primary part – does this hold for all  $p$ ? These are both open.

Why would one expect this to be true?

**Remark 1.6.11**: A recent result:  $\text{cohdim}_p$  grows at most linearly in  $\text{ddim}$ , with slope not 1 but rather  $\approx \log_2 p$ . These questions say that if an equation has enough variables then there is a solution, but why should this be reflected in cohomology? To show this bound, one would want to show that given some  $\alpha \in H^*(k)$ , there exists a polynomial  $f_\alpha$  where if  $f_\alpha$  has a root and  $\alpha = 0$  in homology. In special cases, we were able to come up with such polynomials. When  $\alpha$  is a symbol, this is closely related to *norm varieties* which have a point iff  $\alpha$  is split. One might optimistically hope these are described as hypersurfaces, from which answers to the above would follow, but they turn out to not have such a concrete realization.

## 1.7 Structural Problems in Galois Cohomology

**Remark 1.7.1:** Here we'll describe the problems we need help with! Perhaps insight from motivic cohomology will lend insight to them. We'll write  $H^i(k) := H^i(k; \mu_\ell^{\otimes j})$ .

### 1.7.1 Period-Index Problems

**Definition 1.7.2** (An extension splitting a cohomology class)

If  $\alpha \in H^i(k)$ , we say  $L/k$  **splits**  $\alpha$  if

$$\alpha|_L = 0 \in H^i(L).$$

**Definition 1.7.3** (?)

We define the **index**

$$\text{ind } \alpha := \gcd \{ [L : k] \mid L/k \text{ finite and splits } \alpha \}.$$

and the **period** of  $\alpha$  as its (group-theoretic) order  $H^i(k)$ . Note that  $\text{per } \alpha \leq \ell$ .

**Remark 1.7.4:** One can show that  $\text{per } \alpha \mid \text{ind } \alpha$ , and  $\text{ind } \alpha \mid (\text{per } \alpha)^m$  for some  $m$ .

#### Question 1.7.5

For a fixed  $k$  and  $i, j, \ell$ , which is the minimum  $m$  such that

$$\text{ind } \alpha \mid (\text{per } \alpha)^m?$$

Alternatively, what is the minimum  $m$  such that  $\text{ind } \alpha \mid \ell^m$ ?

#### Conjecture 1.7.6.

If  $\text{ddim}(k) = n$  (or  $\dim(k) = n$  since  $k$  is finitely generated) with  $\alpha \in H^2(k, \mu_\ell)$ , then

$$\text{ind } \alpha \mid (\text{per } \alpha)^{n-1}.$$

**Remark 1.7.7:** Even in this case, no known bound is known for  $k = \mathbb{Q}(t)$ , for any choice of  $\ell$ . How complicated can the cohomology class be? The rough idea is that for  $H^i(k)$  with  $i$  near  $\dim k$ , this should have a small index and if  $i = \dim k$  then  $\text{per } k = \text{ind } k$ .

**Remark 1.7.8:** We know  $\text{per} = \text{ind}$  for any number field for classes in  $H^2(\text{Spec } k; \mu_N)$ , with or without roots.

### 1.7.2 Symbol Length Problem

**Remark 1.7.9:** We know  $H^n(k, \mu_\ell^{\otimes n})$  is generated by symbols  $(a_1, a_2, \dots, a_n)$ . We can use symbol length to measure complexity, leading to the following:



**Question 1.7.10**

Given  $k, n$ , what is the minimal number  $m$  such that every  $\alpha \in H^n(k)$  is a sum of no more than  $m$  symbols. I.e. how easy is it to write  $\alpha$ ?

**Remark 1.7.11:** We'd like a bound in terms of  $\text{ddim}(k)$  and  $\text{dim}(k)$ . One can construct fields needing arbitrarily long symbols, but perhaps for finite dimensional fields, one feels there should be a bound. Danny feels that there may not be such a bound once  $n \geq 4$ .

**Remark 1.7.12:** What's known: for number fields (or global fields, i.e. a reasonable notion of dimension with  $\text{dim } k = 2$ ) which lie over finitely generated or prime fields and have a primitive  $\ell$ th root of unity, we know every class in  $H^2$  can be written as exactly one symbol.

**Remark 1.7.13:** A result of Malgri (?): assuming we have roots of unity, if  $\ell = p^t$ , then for  $H^2$  one needs at most  $t(p^{\text{ddim}(k)-1} - 1)$  symbols. If  $\text{ddim}(k) < \infty$  this yields a bound, and conjecturally this shouldn't depend on ???

For higher degree cohomology, we know almost nothing except for special cases of  $H^4$  for "3-dimensional"  $p$ -adic curves.

**Remark 1.7.14:** If one can bound the symbol length, one can uniformly write down a generic element in cohomology as a sum of at most  $n$  symbols. The inability to be able to write down a general form of a cohomology class for a given field is what makes this difficult – they have "complexity" that isn't necessarily bounded in a known way.

## 2 | Tuesday, July 13

**Remark 2.0.1:** Fix a  $k_0 \in \text{Field}$ .

### Outline

- Arithmetic problems: consider "complexity" of cohomology or algebraic structures (Witt group, symbol length, index of classes).
  - Examples were  $\text{ddim}$ ,  $\text{cohdim}$ , the period-index problem, the period-symbol length problem, which we saw last time.
- Algebraic structure problems: describe (algebraic) structural features of the class of all field extensions  $k \in \text{Field}/_{k_0}$ .

Today we'll describe a way to connect these using a notion of *essential dimension*. Computing this is difficult in general, but finding lower/upper bounds can be tractable. We'll get upper bounds from *canonical dimensions*, and lower bounds from cohomological invariants.

**Remark 2.0.2:** For a particularly concrete problem, consider

$$\alpha \in H^2(k; \mu_\ell) \subseteq H^2(k; \mathbb{G}_m)[\ell] := \text{Br}(k)[\ell],$$

i.e. this is a subgroup of the  $\ell$ -torsion of the **Brauer group**. Suppose we know

$$\text{ind } \alpha := \gcd \{ [L : k] \mid \alpha_L = 0 \} = \min \{ [L : K] \mid \alpha_L = 0 \},$$

where the last equality holds in the special case of  $\text{Br}(k)$ . If  $k$  contains a primitive  $\ell$ th root of unity, we can identify  $\mu_\ell = \mathbb{Z}/\ell = \mu_\ell^{\otimes 2}$ , and thus identify

$$H^2(k; \mu_\ell) = H^2(k; \mu_\ell^{\otimes 2}) = K_2^M(k)/\ell.$$

So we can write  $\alpha = \alpha_1 + \cdots + \alpha_r$  as a sum of symbols with  $\alpha_i = (b_i, c_i)_\ell$  with  $b_i, c_i \in k^\times$ .

### Question 2.0.3

How big does  $n$  have to be?

**Remark 2.0.4:** It follows from “the literature” (after stringing several results together) that there almost exists an absolute bounds depending only on  $\ell, n$  but not  $k$ . However, we do not know what this bound actually is. There are some known cases:

- $\ell = n = 2, 3$ :  $r \leq 1$ , so only one symbol is needed.
- $\ell = n = 4$ : probably  $r \leq 4$ .
- $\ell = 2, n = 4$ :  $r \leq 2$ , a classical results on central simple algebras.
- $\ell = 2, n = 8$ :  $r \leq 4$

**Remark 2.0.5:** It turns out that if  $k$  contains a field  $k_0$  with  $\text{ddim } k_0 < \infty$ , one can produce an explicit bound. Given some  $\alpha \in H^2(k; \mu_\ell)$  we can find some  $k_0 \subseteq L \subseteq k$  with  $L$  finitely generated over  $k_0$  and  $[L : k_0]_{\text{tr}}$  depending only on the period  $\ell$  and index  $n$ , such that  $\alpha \in \text{im} \left( H^2(L; \mu) \rightarrow H^2(k; \mu) \right)$ .

### Slogan 2.0.6

Central simple algebras of a given period and index have finite essential dimension.

An important property is that

$$\text{ddim } L \leq \text{ddim } k_0 + [L : k_0]_{\text{tr}}.$$

Recall that we can bound the symbol length in  $H^2(k; \mu_\ell)$  in terms of  $\text{ddim } L$ . The idea is that we can bound the transcendence degree in terms of  $\ell, n$ . This bound can be made very explicit, although it's not tight: for  $\ell = 2, n = 8$ , it's  $2^{17 + \text{ddim } k_0} - 1$ . This is an improvement over  $k_0 = \mathbb{Q}$  though, where it's known there's a bound but it can't be written down. The lower bound is *very* low: it is hard to show a symbol can not be written with very few symbols.

## 2.1 Pfister Form

**Remark 2.1.1:** Recall  $W(k)$ , whose elements are isometry classes of nondegenerate quadratic forms with addition given by perpendicular sum and the Kronecker product. There is a hyperbolic form  $xy$ , or  $x^2 - y^2$  in  $\text{ch } k \neq 2$ , which we can write as  $\langle 1, -1 \rangle$ , and a fundamental ideal of even-dimensional forms  $\langle 1, -a \rangle = \langle\langle a \rangle\rangle$ . We write

$$\langle\langle a_1, a_2, \dots, a_n \rangle\rangle := \langle\langle a_1 \rangle\rangle \langle\langle a_2 \rangle\rangle \cdots \langle\langle a_n \rangle\rangle \in I^n(k),$$

which in fact generate  $I^n(k)$ .

### Question 2.1.2

Given  $q \in I^n(k)$  of dimension  $d$ , we know we can write  $q \sim q_1 \perp \cdots \perp q_r$  where  $q_i$  are  $n$ -fold Pfister forms. How many are needed? Is this number even bounded?

### Theorem 2.1.3 (*Vishik*).

If  $d < 2^n + 2^{n-1}$  then  $r$  is bounded by some small number.

**Remark 2.1.4:** For  $d \geq 2^n + 2^{n-1}$ , it's not so clear, although it is bounded when  $n \geq 3$ . Why is  $n \leq 3$  easy and  $n \geq 4$  hard?

**Remark 2.1.5:** Consider the following objects:

- $H^2(k; \mu)$
- $\text{Br}(k)$
- $W(k)$
- $I^n(k)$
- $q \in I^n(k)$  with  $\dim q = d$

These can all be viewed as functors  $\text{Field}/k_0 \rightarrow \text{Set}$  taking field extensions to sets.


### Definition 2.1.6 (Essential dimension of a functor)

Given a functor  $f$  and  $\alpha \in F(k)$ , define

$$\begin{aligned} \text{essdim}(\alpha) &= \min \left\{ [L : k_0]_{\text{tr}} \mid \alpha \in \text{im}(F(L) \rightarrow F(k)) \right\} \\ \text{essdim}(F) &= \min \left\{ \text{essdim}(\alpha) \mid \alpha \in F(k) \forall k/k_0 \right\}. \end{aligned}$$

### Definition 2.1.7 (Versal)

Given a functor  $F : \text{Alg}/k_0 \rightarrow \text{Set}$ , we say  $\alpha \in F(R)$  is **versal** if for every  $\beta \in F(K)$ , for any  $k/k_0$ , there exists a morphism  $R \rightarrow k$  such that  $\beta$  is the image of  $\alpha$  under  $F(R) \rightarrow F(k)$ .

**Observation 2.1.8**

If there exists a versal  $\alpha \in F(R)$  then  $\text{krulldim } R \geq \text{essdim}(F)$ , so the essential dimension is bounded above by the Krull dimension.

**Example 2.1.9(?)**: Let  $F(k)$  be the set of quadratic forms of dimension  $n$  over  $k$ , then  $\text{essdim } F = n$ . Every such  $q$  can be diagonalized to yields  $q \simeq \langle a_1, a_2, \dots, a_n \rangle$  which is defined over  $L := k_0(a_1, a_2, \dots, a_n)$ . Alternatively,

$$q = \langle x_1, x_2, \dots, x_n \rangle / k_0[x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_n^{\pm 1}]$$

is versal. Thus every such quadratic form comes from “specializing”.

Considering now the fundamental ideals, the Milnor conjectures yield an isomorphism  $I^n/I^{n+1} \cong H^n(k; \mu_2)$ , so there is a SES

$$1 \rightarrow I^{n+1} \rightarrow I^n \xrightarrow{e_n} H^n(k; \mu_2) \rightarrow 1.$$

Thus a quadratic form  $q$  of dimension  $d$  in  $I^{n+1}$  is equivalent to  $q \in I^n$  such that  $e_n(q) = 0$ . ✍

## 2.2 Canonical Dimension

**Definition 2.2.1** (Canonical Dimension)

This is a generalization of  $\text{essdim}$ . Letting  $k/k_0$ , suppose  $F : \text{Field}/k \rightarrow \text{Set}_+$  is a functor now

from extensions of  $k$  (not  $k_0$ ) into pointed sets. For  $\alpha \in F(k)$ , define a new functor

$$\check{F}_\alpha(L) := \begin{cases} \emptyset & \alpha_L \neq \text{pt} \\ \{\text{pt}\} & \alpha_L = \text{pt}, \end{cases}$$

and define the **canonical dimension**

$$\text{candim}(\alpha) = \text{essdim}(\check{F}(\alpha)).$$

**Remark 2.2.2:** This measures how many parameters are needed to trivialize/split  $\alpha$ . To have  $\text{candim}(\alpha) \leq r$  means that if  $\alpha = \text{pt}$  means the following: if  $\alpha_L = \text{pt}$  then there exists an  $E$  with  $k \subseteq E \subseteq L$  with  $[E : k]_{\text{tr}} \leq r$  such that  $\alpha_E = \text{pt}$ .

**Definition 2.2.3** (Generic splitting scheme)

Given  $F$  as above and  $\alpha \in F(k)$ , we say an  $X \in \text{Sch}/_k$  is a **generic splitting scheme** for  $\alpha$  if

$$\alpha_L = 0 \iff X(L) \neq \emptyset.$$

**Remark 2.2.4:** So this encodes triviality of  $\alpha$  into polynomial equations.

**Example 2.2.5 (?)**: If  $X$  is a generic splitting scheme for  $\alpha$  finite type over  $L$  implies  $\text{candim}(\alpha) \leq \dim X$ .

**Question 2.2.6**

Does there exist a finite type generic splitting scheme for cohomology classes in  $H^i(k; \mu_\ell^{\otimes j})$ ?

**Remark 2.2.7:** We do know this in special cases:

- $i = 1$ : Yes, these are étale algebras, so finite schemes over  $k$ .
- $i = 2$ : Yes, Danny shows these exist for all twists.
  - $j = 1$ : Classical, these are Severi-Brauer varieties.
- For symbols,  $i = 3, j = 2, \ell$  a prime: see Merkurjev-Suslin
- For symbols,  $i = 4, j = 3, \ell = 3$ : see Albert algebras
- For symbols,  $\ell$  prime: this can be done up to prime-to- $\ell$  extensions, see [Rost's "Norm Varieties"](#). Related to Bloch-Kato conjecture.
- For symbols,  $\ell = 2$ : see Pfister quadrics.

**Remark 2.2.8:** Upshot: if there exist generic splitting schemes for classes in  $H^i(k; \mu_2)$  for  $i \geq 3$ , one could bound Pfister numbers and thus  $\text{essdim}$ . Write  $\mathcal{I}_d^n(k)$  to be the set of quadratic forms of dimension  $d$  in  $I^n$ , then  $\text{essdim}(\mathcal{I}_d^n) < \infty$  would imply that if  $q \in \mathcal{I}_d^n(k)$  for  $k \supseteq k_0$  then  $q$  would be defined over some  $L/k_0$  with  $[L : k_0]_{\text{tr}} < \infty$ .

If we knew that  $\text{ddim } k_0 < \infty$ , e.g. if  $k_0$  contains a finite field, this yields a bound on  $\text{ddim } L$  and thus on  $\text{cohdim } L$ . If there is a versal element in  $\alpha \in \mathcal{I}_d^n$ , then  $\alpha$  needs some finite number  $m$  of

Pfister forms to be written. Everything else is a specialization of  $\alpha$ , so the length  $m$  will almost give an upper bound.

 **Warning 2.2.9**

This may seem like a correct argument, but it is not! A problem arises where you may have denominators – specialization can get worse, but only a finite number of times, which is how the actual argument goes.

**Remark 2.2.10:** If you knew the essential dimensions were finite with some given bound, and some general period-index conjecture were known, these would give bounds on symbol length in  $H^i(L; \mu_2)$ . There's an argument pushing things into higher powers of the fundamental ideal, thus higher degree cohomology, which disappear at some point and yield a bound. Motives enter the picture in terms of the tools used to attack these problems.