Integral Affine Structures

Sterk's Conventions

○ : -4 Vectors

● : -2 Vectors

edges	g_{ij}	m_{ij}
● ●	0	2
●—●	½	3
●═●	$1/\sqrt{2}$	4
●≡●	$\sqrt{3}/2$	6
●—∞—●	1	∞
●—●	1	



IAS², Q = 3



IAS², Q = 3+2=5.



IAS², Q = 4.



IAS², Q = 4+2=6.

A full KPP model for $\tilde{A}_2$.

Step 1: Coxeter diagram



Step 2: Toric fan Σ whose T-invariant curves have intersection complex $\tilde{A}_2$



Cones

fan

Σ: $e_0, e_1, -e_0-e_1$.

Step 3: Polytope P_Σ (or P_Σ°?)



Orange just tracks interior verts.

Step 4: Glue $P \cup P^{op}$



12 Verts on equator

2·3 = 6 verts in hemispheres.

18 total verts

= S² "with a lattice"

This hides some details: really we need...

Step 3a: Complete to a triangulation by basis triangles.



I see...
2·6 + 4 = 16
triangles of area ½ each.

Step 4, redux:



A little complicated to draw precisely

Let's work w/ this instead

Step 5: The "tropical" surface (= union of (almost) toric sfcs = "broken toric sfc") = $\mathcal{X}_0$ for $\mathcal{X}$ a KPP model

$a_i = a_i'$
$b_i = b_i'$
$c_i = c_i'$

Some counts:



After identifications, we have a simplicial set with...

- 3+3+12 = 18 vertices
- 3 edges
- 2 faces

Stars at vertices:

Star @ p_1 = a fan.

Some interior:

p_1:



$= Bl_2(\mathbb{P}^1 \times \mathbb{P}^1)$ ⤳





---

p_4:



$= Bl_2(\mathbb{P}^1 \times \mathbb{P}^1)$



---

p_3:



$= Bl_2(\mathbb{P}^1 \times \mathbb{P}^1)$



---

p_6:



$= Bl_2(\mathbb{P}^1 \times \mathbb{P}^1)$



Some corners:



0: $a_1 = a_1'$ … $b_3 = b_3'$ = $\mathbb{P}^1$

∞: $b_1 = b_1'$ … $c_3 = c_3'$ = $\mathbb{P}^1$

1: $c_1 = c_1'$ … $a_3 = a_3'$ = $\mathbb{P}^1$

Not surfaces, what happened?? Gluing introduced an I_1 singularity, giving us an extra "monodromy" ray in the fan! We have a different procedure for sfcs at singular pts.



Some on the edges:

a_1:



$= Bl_1(\mathbb{P}^1 \times \mathbb{P}^1)$

$= Bl_1(\mathbb{F}_n)$



Note: no singularities on these vertices! So toric!

a_2:







c_2:





... etc!

So what is our broken sfc explicitly?

We interpret this picture

Some counts:



- 3+3+12 = 18 vertices
- 3 edges
- 2 faces

as...

$$\mathcal{X}_0 = \left(\bigcup_{i=0}^{6} V_{p_i}\right) \cup \left(\bigcup_{i=1}^{3} V_{a_i}\right) \cup \left(\bigcup_{i=1}^{3} V_{b_i}\right) \cup \left(\bigcup_{i=1}^{3} V_{c_i}\right)$$

$$\cup\, V_0 \cup V_1 \cup V_\infty$$

$$= 6+3+3+3+3 = 18 \text{ sfcs glued together}$$

Where V_0, V_1, V_∞ are semitoric sfcs with one I_1 singularity (equiv: Charge = 1, one nontoric blowup of some toric sfc.)

and the rest are honest toric sfcs.

How are they glued? E.g.



V_{p_1} $\xrightleftharpoons{\text{glue}}$ V_{a_1}

p_1:



a_1:





Glue red



⤳ yields

This describes V_{p_1} glued to V_{a_1} along a common curve, the edge $p_1 \to a_1$ in red

So $\mathcal{X}_0$ = (15 toric sfcs) ∪ (3 almost toric sfcs with Charge = 1)

Q: Can this be smoothed to a family $\mathcal{X}$?

A: When Q = 24, yes, we use a theorem of Freedman

(Note: $\mathcal{X}$ is a highly non-unique KPP model).

One last lie to clear up: what is the actual semitoric sfc at a node?

Each node p_j represents an ACP (V_j, D_j) where $D_j = \sum_i (D_j)_i$.

Set $\mathcal{F}(V_i, D_i) := \mathrm{Cone}(\underbrace{\Gamma(D_i)}_{\text{dual complex}})$, a triangulated IA sfc.

Dual complex =



Cone on dual complex:

Cone (



) =





The actual degeneration:

Blue = a cartoon "tropical divisor" (a weighted balanced subgraph)



Colored in 2-cells get collapsed. ⤳ Tiling of S²

This is the intersection complex, not the dual complex!!

This is $\mathcal{X}_0$ for a KPP degen.

Each tile corresponds to an irreducible component V_i of $\mathcal{X}_0 = \bigcup_i V_i$

I'm omitting necessary self-intersection numbers along edges, e.g.



Not accurate #s? they must satisfy a triple pt. formula!

Adjacent V_i are glued along an anticanonical cycle of curves.


∞
∞




P
P^opp
I_2